


13.KINETIC THEORY

Single Correct Answer Type

- The speeds of 5 molecules of a gas (in arbitrary units) are as follows: 2,3,4,5,6. The root mean square speed for these molecules is
 a) 2.91 b) 3.52 c) 4.00 d) 4.24
- The rate of cooling at 600 K, if surrounding temperature is 300 K is R . The rate of cooling at 900 K is
 a) $\frac{16}{3}R$ b) $2R$ c) $3R$ d) $\frac{2}{3}R$
- For a diatomic gas change in internal energy for unit change in temperature for constant volume is U_1 and U_2 respectively. $U_1:U_2$ is
 a) 5:3 b) 3:5 c) 1:1 d) 5:7
- The temperature of a piece of metal is increased from 27°C to 84°C . The rate at which energy is radiated is increased to
 a) Four times b) Two times c) Six times d) Eight times
- The kinetic energy of translation of 20 g of oxygen at 47°C is (molecular wt. of oxygen is 32 g/mol and $R = 8.3\text{ J/mol/K}$)
 a) 2490 joules b) 2490 ergs c) 830 joules d) 124.5 joules
- Two thermally insulated vessels 1 and 2 are filled with air at temperatures (T_1, T_2) volume (V_1, V_2) and pressure (P_1, P_2) respectively. If the valve joining the two vessels is opened, the temperature inside the vessel at equilibrium will be
 a) T_1+T_2 b) $(T_1+T_2)/2$ c) $\frac{T_1 T_2 (P_1 V_1 + P_2 V_2)}{P_1 V_1 T_2 + P_2 V_2 T_1}$ d) $\frac{T_1 T_2 (P_1 V_1 + P_2 V_2)}{P_1 V_1 T_1 + P_2 V_2 T_2}$
- The pressure and volume of saturated water vapour are P and V respectively. It is compressed isothermally thereby volume becomes $V/2$, the final pressure will be
 a) More than $2P$ b) P c) $2P$ d) $4P$
- At which temperature the velocity of O_2 molecules will be equal to the velocity of N_2 molecules at 0°C
 a) 40°C b) 93°C c) 39°C d) Cannot be calculated
- Kinetic theory of gases provide a base for
 a) Charle's law b) Boyle's law
 c) Charle's law and Boyle's law d) None of these
- The time average of the kinetic energy of one molecule of a gas taken over a long period of time
 a) Is proportional to the square root of the absolute temperature of the gas
 b) Is proportional to the absolute temperature of the gas
 c) Is proportional to the square of the absolute temperature of the gas
 d) Does not depend upon the absolute temperature of the gas
- Kinetic theory of gases was put forward by
 a) Einstein b) Newton c) Maxwell d) Raman
- In kinetic theory of gases, which of the following statements regarding elastic collisions of the molecules is wrong

- a) Kinetic energy is lost in collisions
 b) Kinetic energy remains constant in collision
 c) Momentum is conserved in collision
 d) Pressure of the gas remains constant in collisions
13. If γ is the ratio of specific heats and R is the universal gas constant, then the molar specific heat at constant volume C_v is given by
 a) γR b) $\frac{(\gamma-1)R}{\gamma}$ c) $\frac{R}{\gamma-1}$ d) $\frac{\gamma R}{\gamma-1}$
14. The vapour of a substance behaves as a gas
 a) Below critical temperature b) Above critical temperature
 c) At 100 °C d) At 1000 °C
15. If the temperature of an ideal gas increases three times, then its *rms* velocity will become
 a) $\sqrt{3}$ times b) 3 times c) One third d) Remains same
16. The relationship between pressure and the density of a gas expressed by Boyle's law, $P=KD$ holds true
 a) For any gas under any conditions b) For some gases under any conditions
 c) Only if the temperature is kept constant d) Only if the density is constant
17. If the ratio of vapour density for hydrogen and oxygen is $\frac{1}{16}$, then under constant pressure the ratio of their *rms* velocities will be
 a) $\frac{4}{1}$ b) $\frac{1}{4}$ c) $\frac{1}{16}$ d) $\frac{16}{1}$
18. The gases carbon-monoxide (CO) and nitrogen at the same temperature have kinetic energies E_1 and E_2 respectively. Then
 a) $E_1 = E_2$ b) $E_1 > E_2$
 c) $E_1 < E_2$ d) E_1 and E_2 cannot be compared
19. What is the mass of 2 L of nitrogen at 22.4 atm pressure and 273 K?
 a) 28 g b) 14×22.4 g c) 56 g d) None of these
20. The average kinetic energy of a gas molecules is
 a) Proportional to pressure of gas b) Inversely proportional to volume of gas
 c) Inversely proportional to absolute temperature of gas d) Directly proportional to absolute temperature of gas
21. The adjoining figure shows graph of pressure and volume of a gas at two temperatures T_1 and T_2 . Which of the following inferences is correct



- a) $T_1 > T_2$ b) $T_1 = T_2$
 c) $T_1 < T_2$ d) No inference can be drawn

22. At room temperature (27°C) the rms speed of the molecules of a certain diatomic gas is found to be 1920 m s^{-1} . The gas is
 a) Cl_2 b) O_2 c) N_2 d) H_2
23. At a given temperature, the pressure of an ideal gas of density ρ is proportional to
 a) $\frac{1}{\rho^2}$ b) $\frac{1}{\rho}$ c) ρ^2 d) ρ
24. Temperature remaining constant, the pressure of gas is decreased by 20%. The percentage change in volume
 a) Increases by 20% b) Decreases by 20% c) Increases by 25% d) decreases by 25%
25. The rms velocity of gas molecules is 300 m s^{-1} . The rms velocity of molecules of gas with twice the molecular weight and half the absolute temperature is
 a) 300 m s^{-1} b) 600 m s^{-1} c) 75 m s^{-1} d) 150 m s^{-1}
26. A jar contains a gas and few drops of water at $T\text{ K}$. The pressure in the jar is 830 mm of mercury. The temperature of jar is reduced by 1%. The saturated vapour pressure of water at the two temperatures are 30 mm and 25 mm of mercury. Then the new pressure in the jar will be
 a) 917 mm of Hg b) 717 mm of Hg c) 817 mm of Hg d) None of these
27. The gas equation $\frac{PV}{T} = \text{constant}$ is true for a constant mass of an ideal gas undergoing
 a) Isothermal change b) Adiabatic change c) Isobaric change d) Any type of change
28. The pressure and temperature of two different gases is P and T having the volume V for each. They are mixed keeping the same volume and temperature, the pressure of the mixture will be
 a) $P/2$ b) P c) $2P$ d) $4P$
29. Vessel A is filled with hydrogen while vessel B , whose volume is twice that of A , is filled with the same mass of oxygen at the same temperature. The ratio of the mean kinetic energies of hydrogen and oxygen is
 a) 16:1 b) 1:8 c) 8:1 d) 1:1
30. The root mean square speed of hydrogen molecules at 300 K is 1930 m/s . Then the root mean square speed of oxygen molecules at 900 K will be
 a) $1930\sqrt{3}\text{ m/s}$ b) 836 m/s c) 63 m/s d) $\frac{1930}{\sqrt{3}}\text{ m/s}$
31. A cylinder rolls without slipping down an inclined plane, the number of degrees of freedom it has, is
 a) 2 b) 3 c) 5 d) 1
32. Two spheres made of same material have radii in the ratio 1:2. Both are at same temperature. Ratio of heat radiation energy emitted per second by them is
 a) 1:2 b) 1:4 c) 1:8 d) 1:16
33. If r. m. s. velocity of a gas is $V_{rms} = 1840\text{ m/s}$ and its density $\rho = 8.99 \times 10^{-2}\text{ kg/m}^3$, the pressure of the gas will be
 a) 1.01 N/m^2 b) $1.01 \times 10^3\text{ N/m}^2$ c) $1.01 \times 10^5\text{ N/m}^2$ d) $1.01 \times 10^7\text{ N/m}^2$
34. An ideal gas ($\gamma = 1.5$) is expanded adiabatically. How many times has the gas to be expanded to reduce the root mean square velocity of molecules 2.0 times?
 a) 4 times b) 16 times c) 8 times d) 2 times
35. The quantity of heat required to raise one mole through one degree kelvin for a monoatomic gas at constant volume is
 a) $\frac{3}{2}R$ b) $\frac{5}{2}R$ c) $\frac{7}{2}R$ d) $4R$


36. Calculate the ratio of rms speeds of oxygen gas molecules to that of hydrogen gas molecules kept at the same temperature.

- a) 1:4 b) 1:8 c) 1:2 d) 1:6

37. At constant pressure, the ratio of increase in volume of an ideal gas per degree rise in kelvin temperature to its original volume is ($T = i$ absolute temperature of the gas)

- a) T^2 b) T c) $1/T$ d) $1/T^2$

38. Pressure versus temperature graphs of an ideal gas are as shown in figure. Choose the wrong statement



- a) Density of gas is increasing in graph (i) b) Density of gas is decreasing in graph (ii)
 c) Density of gas is constant in graph (iii) d) None of these

39. A body takes 10 min to cool from 60°C to 50°C . If the temperature of surroundings is 25°C and 527°C respectively. The ratio of energy radiated by $P \wedge Q$ is

- a) 48°C b) 46°C c) 49°C d) 42.85°C

40. A cylinder of radius r and thermal conductivity K_1 is surrounded by a cylindrical shell of linear radius r and outer radius $2r$, whose thermal conductivity is K_2 . There is no loss of heat across cylindrical surfaces, when the ends of the combined system are maintained at temperatures $T_1 \wedge T_2$. The effective thermal conductivity of the system, in the steady state is

- a) $\frac{K_1 K_2}{K_1 + K_2}$ b) $K_1 + K_2$ c) $\frac{K_1 + 3K_2}{4}$ d) $\frac{3K_1 + K_2}{4}$

41. A gaseous mixture consists of 16 g of helium and 16 g of oxygen. The ratio $\frac{C_p}{C_v}$ of the mixture is

- a) 1.4 b) 1.54 c) 1.59 d) 1.62

42. Mean free path of a gas molecule is

- a) Inversely proportional to number of molecules per unit volume
 b) Inversely proportional to diameter of the molecule
 c) Directly proportional to the square root of the absolute temperature
 d) Directly proportional to the molecular mass

43. The value of densities of two diatomic gases at constant temperature and pressure are d_1 and d_2 , then the ratio of speed of sound in these gases will be

- a) $d_1 d_2$ b) $\sqrt{d_2/d_1}$ c) $\sqrt{d_1/d_2}$ d) $\sqrt{d_1 d_2}$

44. If the internal energy of n_1 moles of He at temperature $10 T$ is equal to the internal energy of n_2 mole of hydrogen at temperature $6 T$. the ratio of $\frac{n_1}{n_2}$ is

- a) $\frac{3}{5}$ b) 2 c) 1 d) $\frac{5}{3}$

45. The heat capacity per mole of water is (R is universal gas constant)

- a) $9R$ b) $\frac{9}{2}R$ c) $6R$ d) $5R$

46. If number of molecules of H_2 are double than that of O_2 , then ratio of kinetic energy of hydrogen and that of

- oxygen at 300 K is
 a) 1:1 b) 1:2 c) 2:1 d) 1:16
47. According to the kinetic theory of gases, the temperature of a gas is a measure of average
 a) Velocities of its molecules b) Linear momenta of its molecules
 c) Kinetic energies of its molecules d) Angular momenta of its molecules
48. Air is filled in a bottle at atmospheric pressure and it is corked at 35 °C . If the cork can come out at 3 atmospheric pressure than upto what temperature should the bottle be heated in order to remove the cork
 a) 325.5 °C b) 851 °C c) 651 °C d) None of these
49. The temperature at which the average translational kinetic energy of a molecule is equal to the energy gained by an electron in accelerating from rest through a potential difference of 1 volt is
 a) $4.6 \times 10^3 K$ b) $11.6 \times 10^3 K$ c) $23.2 \times 10^3 K$ d) $7.7 \times 10^3 K$
50. The average momentum of a molecule in an ideal gas depends on
 a) Temperature b) Volume c) Molecular mass d) None of these
51. If pressure of CO_2 (real gas) in a container is given by $P = \frac{RT}{2V-b} - \frac{a}{4b^2}$, then mass of the gas in container is
 a) 11 g b) 22 g c) 33 g d) 44 g
52. For an ideal gas of diatomic molecules
 a) $C_p = \frac{5}{2}R$ b) $C_v = \frac{3}{2}R$ c) $C_p - C_v = 2R$ d) $C_p = \frac{7}{2}R$
53. What is the value of $\frac{R}{C_p}$ for diatomic gas
 a) 3/4 b) 3/5 c) 2/7 d) 5/7
54. When volume of system is increased two times and temperature is decreased half of its initial temperature, then pressure becomes
 a) 2 times b) 4 times c) $\frac{1}{4} \times i$ d) $\frac{1}{2} \times i$
55. A vessel of volume 4 L contains a mixture of 8 g of oxygen, 14 g of nitrogen and 22 g of carbon dioxide at 27°C. The pressure exerted by the mixture is
 a) $5.79 \times 10^5 N m^{-2}$ b) $6.79 \times 10^5 N m^{-2}$ c) $7.79 \times 10^3 N m^{-2}$ d) $7.79 \times 10^5 N m^{-2}$
56. 2 g of O_2 gas is taken at 27 °C and pressure 76 cm .Hg . Find out volume of gas (in litre)
 a) 1.53 b) 2.44 c) 3.08 d) 44.2
57. When an air bubble of radius 'r' rises from the bottom to the surface of a lake, its radius becomes 5r/4 (the pressure of the atmosphere is equal to the 10 m height of water column). If the temperature is constant and the surface tension is neglected, the depth of the lake is
 a) 3.53 m b) 6.53 m c) 9.53 m d) 12.53 m
58. At what temperature will the rms speed of air molecules be double than that at NTP?
 a) 519°C b) 619°C c) 719°C d) 819°C
59. The kinetic energy per g mol for a diatomic gas at room temperature is
 a) 3 RT b) $\frac{5}{2}RT$ c) $\frac{3}{2}RT$ d) $\frac{1}{2}RT$
60. The average kinetic energy of a gas at -23 °C and 75 cm pressure is $5 \times 10^{-14} \text{ erg}$ for H_2 . The mean kinetic energy of the O_2 at 227 °C and 150 cm pressure will be

- a) $80 \times 10^{-14} \text{ erg}$ b) $20 \times 10^{-14} \text{ erg}$ c) $40 \times 10^{-14} \text{ erg}$ d) $10 \times 10^{-14} \text{ erg}$


61. A monoatomic gas molecule has

- a) Three degrees of freedom b) Four degrees of freedom
c) Five degrees of freedom d) Six degrees of freedom

62. Considering the gases to be ideal, the value of $\gamma = \frac{C_P}{C_V}$ for a gaseous mixture consisting of 3 moles of carbon dioxide and 2 moles of oxygen will be ($\gamma_{O_2} = 1.4, \gamma_{CO_2} = 1.3$)

- a) 1.37 b) 1.34 c) 1.55 d) 1.63

63. The change in volume V with respect to an increase in pressure P has been shown in the figure for a non-ideal gas at four different temperatures T_1, T_2, T_3 and T_4 . The critical temperature of the gas is




- a) T_1 b) T_2 c) T_3 d) T_4

64. At a given temperature the ratio of *r.m.s.* velocities of hydrogen molecule and helium atom will be

- a) $\sqrt{2}:1$ b) $1:\sqrt{2}$ c) $1:2$ d) $2:1$

65. A vessel contains 14 g (7 moles) of hydrogen and 96 g (9 moles) of oxygen at STP. Chemical reaction is induced by passing electric spark in the vessel till one of the gases is consumed. The temperature is brought back to its starting value 273 K. The pressure in the vessel is



- a) 0.1 atm b) 0.2 atm c) 0.3 atm d) 0.4 atm

66. When the temperature of a gas is raised from 27°C to 90°C , the percentage increase in the *r.m.s.* velocity of the molecules will be

- a) 10% b) 15% c) 20% d) 17.5%


67. One litre of oxygen at a pressure of 1 atm and two litres of nitrogen at a pressure of 0.5 atm, are introduced into a vessel of volume 1 L. If there is no change in temperature, the final pressure of the mixture of gas (in atm) is

- a) 1.5 b) 2.5 c) 2 d) 4





68. The power radiated by a black body is P , and it radiates maximum energy around the wavelength λ_0 . If the temperature of black body is now changed so that it radiates maximum energy around a wavelength $\lambda_0/4$, the power radiated by it will increase by a factor of

- a) $\frac{4}{3}$ b) $\frac{16}{9}$ c) $\frac{64}{27}$ d) $\frac{256}{81}$


69. Figure shows two flasks connected to each other. The volume of the flask 1 is twice that of flask 2. The system is filled with an ideal gas at temperature 100 K and 200 K respectively. If the mass of the gas in 1 be m then what is the mass of the gas in flask 2



- 1) m b) $m/2$ c) $m/4$ d) $m/8$
70. Under constant temperature, graph between P and $1/V$ is
a) Parabola b) Hyperbola c) Straight line d) Circle
71. A gas mixture consists of molecules of type 1,2 and 3, with molar masses $m_1 > m_2 > m_3$. V_{rms} and \dot{K} are the *r. m. s.* speed and average kinetic energy of the gases. Which of the following is true
a) $(V_{rms})_1 < (V_{rms})_2 < (V_{rms})_3$ and $(\dot{K})_1 = (\dot{K})_2 = (\dot{K})_3$
b) $(V_{rms})_1 = (V_{rms})_2 \leq (V_{rms})_3$ and $(\dot{K})_1 = (\dot{K})_2 > (\dot{K})_3$
c) $(V_{rms})_1 > (V_{rms})_2 < (V_{rms})_3$ and $(\dot{K})_1 < (\dot{K})_2 > (\dot{K})_3$
d) $(V_{rms})_1 > (V_{rms})_2 > (V_{rms})_3$ and $(\dot{K})_1 < (\dot{K})_2 < (\dot{K})_3$
72. The ratio of mean kinetic energy of hydrogen and nitrogen at temperature 300 K and 450 K respectively is
a) 3:2 b) 2:3 c) 2:21 d) 4:9
73. Equation of gas in terms of pressure (P), absolute temperature (T) and density (d) is
a) $\frac{P_1}{T_1 d_1} = \frac{P_2}{T_2 d_2}$ b) $\frac{P_1 T_1}{d_1} = \frac{P_2 T_2}{d_2}$ c) $\frac{P_1 d_2}{T_1} = \frac{P_2 d_1}{T_2}$ d) $\frac{P_1 d_1}{T_1} = \frac{P_2 d_2}{T_2}$
74. On 0°C pressure measured by barometer is 760 mm. What will be pressure at 100°C
a) 760 mm b) 730 mm c) 780 mm d) None of these
75. The *r. m. s.* speed of the molecules of a gas in a vessel is 400 m s^{-1} . If half of the gas leaks out, at constant temperature, the *r. m. s.* speed of the remaining molecules will be
a) 800 m s^{-1} b) $400\sqrt{2}\text{ m s}^{-1}$ c) 400 m s^{-1} d) 200 m s^{-1}
76. Volume-temperature graph at atmospheric pressure for a monoatomic gas ($V \in \text{m}^3, T \in ^\circ\text{C}$) is



77. The temperature of argon, kept in a vessel, is raised by 1°C at a constant volume. The total heat supplied to the gas is a combination of translation and rotational energies. Their respective shares are
a) 60% and 40% b) 40% and 60% c) 50% and 50% d) 100% and 0%
78. The molar heat capacity at constant volume of oxygen gas at STP is nearly $\frac{5R}{2}$ and it approaches $\frac{7R}{2}$ as the temperature is increased. This happens because at higher temperature

- a) Oxygen becomes triatomic
 c) Oxygen molecules rotate more vigorously
- b) Oxygen does not behaves as an ideal gas
 d) Oxygen molecules start vibrating
79. Three containers of the same volume contain three different gases. The masses of the molecules are m_1, m_2 and m_3 and the number of molecules in their respective containers are N_1, N_2 and N_3 . The gas pressure in the containers are P_1, P_2 and P_3 respectively. All the gases are now mixed and put in one of the containers. The pressure P of mixture will be
 a) $P < (P_1 + P_2 + P_3)$ b) $P = \frac{P_1 + P_2 + P_3}{3}$ c) $P = P_1 + P_2 + P_3$ d) $P > (P_1 + P_2 + P_3)$
80. If temperature of gas increases from 27°C to 927°C the $K.E.$ will be
 a) Double b) Half c) One fourth d) Four times
81. A mixture of 2 moles of helium gas (atomic mass = 4 amu), and 1 mole of argon gas (atomic mass = 40 amu) is kept at 300 K in a container. The ratio of the rms speeds $\left[\frac{V_{rms}(\text{helium})}{V_{rms}(\text{argon})} \right]$ is
 a) 0.32 b) 0.45 c) 2.24 d) 3.16
82. The value of the gas constant (R) calculated from the perfect gas equation is $8.32\text{ joules/g mole K}$, whereas its value calculated from the knowledge of C_p and C_v of the gas is 1.98 cal/g mole K . From this data, the value of J is
 a) 4.16 J/cal b) 4.18 J/cal c) 4.20 J/cal d) 4.22 J/cal
83. S.I. unit of universal gas constant is
 a) $\text{cal}/^\circ\text{C}$ b) J/mol c) $\text{J mol}^{-1}\text{K}^{-1}$ d) J/kg
84. In Boyle's law what remains constant
 a) pV b) TV c) $\frac{V}{T}$ d) $\frac{P}{T}$
85. To what temperature should the hydrogen at 327°C be cooled at constant pressure, so that the root mean square velocity of its molecules becomes half of its previous value?
 a) -123°C b) 123°C c) -100°C d) 0°C
86. Two gases A and B having same pressure p , volume V and absolute temperature T are mixed. If the mixture has the volume and temperature as V and T respectively, then the pressure of the mixture is
 a) $2p$ b) p c) $\frac{p}{2}$ d) $4p$
87. The density (ρ) versus pressure (P) of a given mass of an ideal gas is shown at two temperatures T_1 and T_2
- 
- Then relation between T_1 and T_2 may be
 a) $T_1 > T_2$ b) $T_2 > T_1$
 c) $T_1 = T_2$ d) All the three are possible
88. The gas in vessel is subjected to a pressure of 20 atmosphere at a temperature 27°C . The pressure of the gas in a vessel after one half of the gas is released from the vessel and the temperature of the remainder is raised by 50°C is

- a) 8.5 atm b) 10.8 atm c) 11.7 atm d) 17 atm


89. On any planet, the presence of atmosphere implies (C_{rms} = root mean square velocity of molecules and V_e = escape velocity)

- a) $C_{rms} \ll V_e$ b) $C_{rms} > V_e$ c) $C_{rms} = V_e$ d) $C_{rms} = 0$

90. The degrees of freedom of a stationary rigid body about its axis will be

- a) One b) Two c) Three d) Four

91. From the following $V-T$ diagram we can conclude



- a) $P_1 = P_2$ b) $P_1 > P_2$ c) $P_1 < P_2$ d) None of these

92. An electron tube was sealed off during manufacture at a pressure of 1.2×10^{-7} mm of mercury at 27°C . Its volume is 100 cm^3 . The number of molecules that remain in the tube is

- a) 2×10^{16} b) 3×10^{15} c) 3.86×10^{11} d) 5×10^{11}

93. The average kinetic energy of hydrogen molecules at 300 K is E . At the same temperature, the average kinetic energy of oxygen molecules will be

- a) $E/4$ b) $E/16$ c) E d) $4E$

94. The temperature of an ideal gas is increased from 27°C to 927°C . The root mean square speed of its molecules becomes

- a) Twice b) Half c) Four times d) One-fourth

95. A given mass of a gas is allowed to expand freely until its volume becomes double. If C_b and C_a are the velocities of sound in this gas before and after expansion respectively, then C_a is equal to

- a) $2C_b$ b) $\sqrt{2}C_b$ c) C_b d) $\frac{1}{\sqrt{2}}C_b$

96. For a gas at a temperature T the root-mean-square velocity v_{rms} , the most probable speed v_{mp} , and the average speed v_{av} obey the relationship

- a) $v_{av} > v_{rms} > v_{mp}$ b) $v_{rms} > v_{av} > v_{mp}$ c) $v_{mp} > v_{av} > v_{rms}$ d) $v_{mp} > v_{rms} > v_{av}$

97. Two chambers containing m_1 and m_2 gram of a gas at pressures p_1 and p_2 respectively are put in communication with each other, temperature remaining constant. The common pressure reached will be

- a) $\frac{p_1 p_2 (m_1 + m_2)}{p_2 m_1 + p_1 m_2}$ b) $\frac{p_1 p_2 m_1}{p_2 m_1 + p_1 m_2}$ c) $\frac{m_1 m_2 (p_1 + p_2)}{p_2 m_1 + p_1 m_2}$ d) $\frac{m_1 m_2 p_2}{p_2 m_1 + p_1 m_2}$


98. The root mean square speed of the molecules of a diatomic gas is v . When the temperature is doubled, the molecules dissociate into two atoms. The new root mean square speed of the atom is

- a) $\sqrt{2}v$ b) v c) $2v$ d) $4v$

99. The ends of 2 different materials with their thermal conductivities, radii of cross section and length all in the ratio of 1 : 2 maintained at temperature difference. If the rate of the flow of heat in the longer rod is 4 cal s^{-1} , that in the shorter rod in cal s^{-1} will be

- a) 1 b) 2 c) 8 d) 6

100. An experiment is carried on a fixed amount of gas at different temperatures and at high pressure such that it deviates from the ideal gas behavior. The variation of $\frac{PV}{RT}$ with P is shown in the diagram. The correct variation will correspond to




- a) Curve A b) Curve B c) Curve C d) Curve D

101. A gas is filled in a cylinder, its temperature is increased by 20% on kelvin scale and volume is reduced by 10%. How much percentage of the gas will leak out
 a) 30% b) 40% c) 15% d) 25%

102. The degrees of freedom of a molecule of a triatomic gas are
 a) 2 b) 4 c) 6 d) 8

103. Six molecules speeds 2 unit, 5 unit, 3 unit, 6 unit, 3 unit, and 5 unit respectively. The rms speed is
 a) 4 unit b) 1.7 unit c) 4.2 unit d) 5 unit

104. Which one of the following graph is correct at constant pressure



105. The tyre of a motor car contains air at 15°C . If the temperature increases to 35°C , the approximate percentage increase is (ignore to expansion of tyre)
 a) 7 b) 9 c) 11 d) 13

106. The temperature of the hydrogen at which the average speed of its molecules is equal to that of oxygen molecules at a temperature of 31°C , is
 a) -216°C b) -235°C c) -254°C d) -264°C

107. The kinetic energy of one gram molecule of a gas at normal temperature and pressure is ($R=8.31\text{ J/mol-K}$)
 a) $0.56 \times 10^4\text{ J}$ b) $1.3 \times 10^2\text{ J}$ c) $2.7 \times 10^2\text{ J}$ d) $3.4 \times 10^3\text{ J}$

108. The temperature of a gas contained in a closed vessel of constant volume increases by 1°C when the pressure of the gas is increased by 1%. The initial temperature of the gas is
 a) 100 K b) 273°C c) 100°C d) 200 K

109. 70 cal of heat is required to raise the temperature of 2 moles of an ideal gas from 30°C to 35°C while the pressure of the gas is kept constant. The amount of the heat required to raise the temperature of the same gas through the same temperature range at constant volume is (gas constant $R=2\text{ cal mol}^{-1}\text{-K}^{-1}$)
 a) 70 cal b) 60 cal c) 50 cal d) 30 cal


110. A sample of gas is at 0°C . To what temperature it must be raised in order to double the r.m.s. speed of the molecule
 a) 270°C b) 819°C c) 1090°C d) 100°C


111. If p is the pressure, V the volume, R the ags constant, k the Boltzmann's constant and T the absolute temperature, then the number of molecules in the given mass of the gas is given by
 a) $\frac{pV}{RT}$ b) $\frac{pV}{kT}$ c) $\frac{pR}{T}$ d) pV

112. The pressure is P , volume V and temperature T of a gas in the jar A and the other gas in the jar B is at pressure $2P$, volume $V/4$ and temperature $2T$, then the ratio of the number of molecules in the jar A and B will be


- a) 1:1 b) 1:2 c) 2:1 d) 4:1
113. Suppose ideal gas equation follows $V P^3 = \text{constant}$. Initial temperature and volume of the gas are T and V respectively. If gas expands to $27V$ then its temperature will become
a) T b) $9T$ c) $27T$ d) $T/9$
114. For ideal gas, which statement is not true
a) It obeys Boyle's law b) It follows $PV = RT$
c) Internal energy depends on temperature only d) It follows Vander-Waal's equation
115. $\frac{1}{2}$ mole of helium gas is contained in a container at S.T.P. The heat energy needed to double the pressure of the gas, keeping the volume constant (specific heat of the gas $3 J g m^{-1} K^{-1}$) is
a) $3276 J$ b) $1638 J$ c) $819 J$ d) $409.5 J$
116. A vertical column 50 cm long at $50^\circ C$ balances another column of liquid 60 cm long at $100^\circ C$. The coefficient of absolute expansion of the liquid is
a) $0.005^\circ C^{-1}$ b) $0.0005^\circ C^{-1}$ c) $0.002^\circ C^{-1}$ d) $0.0002^\circ C^{-1}$
117. The diameter of oxygen molecule is $2.94 \times 10^{-10} m$. The Vander Waal's gas constant ' b ' in m^3/mol will be
a) 3.2 b) 16 c) 32×10^{-4} d) 32×10^{-6}
118. In a certain region of space there are only 5 molecules per cm^2 on an average. The temperature there is 3 K. The pressure of this dilute gas is ($k = 1.38 \times 10^{-23} J K^{-1}$)
a) $20.7 \times 10^{-17} N m^{-1}$ b) $15.3 \times 10^{-13} N m^{-1}$ c) $2.3 \times 10^{-10} N m^{-1}$ d) $5.3 \times 10^{-5} N m^{-1}$
119. The temperature at which the r. m. s. speed of hydrogen molecules is equal to escape velocity on earth surface, will be
a) 1060 K b) 5030 K c) 8270 K d) 10063 K
120. What is the velocity of wave in monoatomic gas having pressure 1 kilopascal and density $2.6 kg/m^3$
a) 3.6 m/s b) $8.9 \times 10^3 m/s$ c) Zero d) None of these
121. The temperature at which protons in proton gas would have enough energy to overcome. Coulomb barrier of $4.14 \times 10^{-14} J$ is (Boltzman constant $1.38 \times 10^{-23} J K^{-1}$)
a) $2 \times 10^9 K$ b) $10^9 K$ c) $6 \times 10^9 K$ d) $3 \times 10^9 K$
122. KE per unit volume is E . The pressure exerted by the gas is given by
a) $\frac{E}{3}$ b) $\frac{2E}{3}$ c) $\frac{3E}{2}$ d) $\frac{E}{2}$
123. Two cylindrical conductors $A \wedge B$ of same metallic material have their diameters in the ratio 1:2 and lengths in the ratio 2:1. If the temperature difference between their ends is same, the ratio of heat conducted respectively by $A \wedge B$ per second is
a) 1:2 b) 1:4 c) 1:16 d) 1:8
124. A gas is collected over the water at $25^\circ C$. The total pressure of moist gas was 735 mm of mercury. If the aqueous vapour pressure at $25^\circ C$ is 23.8 mm. Then the pressure of dry gas is
a) 760 mm b) 758.8 mm c) 710.8 mm d) 711.2 mm
125. Two moles of oxygen is mixed with eight moles of helium. The effective specific heat of the mixture at constant volume is
a) 1.3 R b) 1.4 R c) 1.7 R d) 1.9 R
126. Mean kinetic energy (or average energy) per g molecule of a monoatomic gas is given by

- a) $\frac{3}{2}RT$ b) $\frac{1}{2}kT$ c) $\frac{1}{2}RT$ d) $\frac{3}{2}kT$
127. A cylinder of fixed capacity 44.8 litre contains a monoatomic gas at standard temperature and pressure. The amount of heat required to cylinder by 10°C will be
(R = universal gas constant)
a) R b) $10R$ c) $20R$ d) $30R$
128. Air is pumped into an automobile tube upto a pressure of 200 kPa in the morning when the air temperature is 22°C . During the day, temperature rises to 42°C and the tube expands by 2%. The pressure of the air in the tube at this temperature, will be approximately
a) 212 kPa b) 209 kPa c) 206 kPa d) 200 kPa
129. The volume of a gas at pressure $21 \times 10^4 \text{ N/m}^2$ and temperature 27°C is 83 litres. If $R = 8.3 \text{ J/mol K}$, then the quantity of gas in $g\text{-mole}$ will be
a) 15 b) 42 c) 7 d) 14
130. What is an ideal gas?
a) One that consists of molecules b) A gas satisfying the assumptions of kinetic theory
c) A gas having Maxwellian distribution of speed d) A gas consisting of massless particles
131. The relation between the gas pressure P and average kinetic energy per unit volume E is
a) $P = \frac{1}{2}E$ b) $P = E$ c) $P = \frac{3}{2}E$ d) $P = \frac{2}{3}E$
132. For a gas $\gamma = 7/5$. The gas may probably be
a) Helium b) Hydrogen c) Argon d) Neon
133. When a vander waal's gas undergoes free expansion then its temperature
a) Decreases b) Increases
c) Does not change d) Depends upon the nature of the gas
134. If the oxygen (O_2) has root mean square velocity of $C \text{ m s}^{-1}$, then root mean square velocity of the hydrogen (H_2) will be
a) $C \text{ m s}^{-1}$ b) $\frac{1}{C} \text{ m s}^{-1}$ c) $4C \text{ m s}^{-1}$ d) $\frac{C}{4} \text{ m s}^{-1}$
135. A gas at the temperature 250 K is contained in a closed vessel. If the gas is heated through 1 K , then the percentage increase in its pressure will be
a) 0.4 % b) 0.2 % c) 0.1 % d) 0.8 %
136. To what temperature should the hydrogen at room temperature (27°C) be heated at constant pressure so that the R.M.S. velocity of its molecules becomes double of its previous value
a) 1200°C b) 927°C c) 600°C d) 108°C
137. Consider a collection of a large number of particles each with speed v . The direction of velocity is randomly distributed in the collection. What is the magnitude of the relative velocity between a pairs in the collection
a) $2V/\pi$ b) V/π c) $8V/\pi$ d) $4V/\pi$
138. A pressure cooker contains air at 1 atm and 30°C . If the safety value of the cooler blows when the inside pressure $\geq 3 \text{ atm}$, then the maximum temperature of the air, inside the cooker can be
a) 90°C b) 636°C c) 909°C d) 363°C
139. The value of $\frac{pV}{T}$ for one mole of an ideal gas is nearly equal to

- a) $2 J mol^{-1} K^{-1}$ b) $8.3 J mol^{-1} K^{-1}$ c) $4.2 J mol^{-1} K^{-1}$ d) $2 cal mol^{-1} K^{-1}$
140. $CO_2(O-C-O)$ is a triatomic gas. Mean kinetic energy of one gram gas will be (If N -Avogadro's number, k -Boltzmann's constant and molecular weight of $CO_2=44$)
 a) $(3/88) NkT$ b) $(5/88) NkT$ c) $(6/88) NkT$ d) $(7/88) NkT$
141. To double the volume of a given mass of an ideal gas at $27^\circ C$ keeping the pressure constant, one must raise the temperature in degree centigrade to
 a) 54° b) 270° c) 327° d) 600°
142. The following sets of values for C_V and C_P of a gas has been reported by different students. The units are $cal/g-mole-K$. Which of these sets is most reliable
 a) $C_V=3, C_P=5$ b) $C_V=4, C_P=6$ c) $C_V=3, C_P=2$ d) $C_V=3, C_P=4.2$
143. At what temperature is the root mean square velocity of gaseous hydrogen molecules equal to that of oxygen molecules at $47^\circ C$
 a) $20 K$ b) $80 K$ c) $-73 K$ d) $3 K$
144. Molecules of a gas behave like
 a) Inelastic rigid sphere b) Perfectly elastic non-rigid sphere
 c) Perfectly elastic rigid sphere d) Inelastic non-rigid sphere
145. A cylinder contains $10 kg$ of gas at pressure of $10^7 N/m^2$. The quantity of gas taken out of the cylinder, if final pressure is $2.5 \times 10^6 N/m^2$, will be (Temperature of gas is constant)
 a) $15.2 kg$ b) $3.7 kg$ c) Zero d) $7.5 kg$
146. In the adjoining figure, various isothermals are shown for a real gas. Then
- 
- a) EF represents liquification b) CB represents liquification
 c) HI represents the critical temperature d) AB represents gas at a high temperature
147. One mole of an ideal monoatomic gas requires $210 J$ heat to raise the temperature by $10 K$, when heated at constant temperature. If the same gas is heated at constant volume to raise the temperature by $10 K$ then heat required is
 a) $238 J$ b) $126 J$ c) $210 J$ d) $350 J$
148. The ratio of root mean square velocity of $O_3 \wedge O_2$ is
 a) $1:1$ b) $2:3$ c) $3:2$ d) $\sqrt{2}:\sqrt{3}$
149. At a given temperature the $r. m. s.$ velocity of molecules of the gas is
 a) Same
 b) Proportional to molecular weight
 c) Inversely proportional to molecular weight
 d) Inversely proportional to square root of molecular weight
150. Graph of specific heat at constant volume for a monoatomic gas is



151. PV versus T graph of equal masses of H_2 , He and O_2 is shown in fig. Choose the correct alternative



- a) C corresponds to H_2 , B to He and A to O_2 b) A corresponds to He , B to H_2 and C to O_2
 c) A corresponds to He , B to O_2 and C to H_2 d) A corresponds to O_2 , B to H_2 and C to He

152. Which of the following cylindrical rods will conduct maximum heat, when their ends are maintained at a constant temperature difference?

- a) $l=1\text{ m}, r=0.2\text{ m}$ b) $l=1\text{ m}, r=0.1\text{ m}$ c) $l=10\text{ m}, r=0.1\text{ m}$ d) $l=0.1\text{ m}, r=0.3\text{ m}$

153. A container with insulating walls is divided into two equal parts by a partition fitted with a valve. One part is filled with an ideal gas at a pressure p and temperature T , whereas the other part is completely evacuated. If the valve is suddenly opened, the pressure and temperature of the gas will be

- a) $\frac{p}{2}, T$ b) $\frac{p}{2}, \frac{T}{2}$ c) p, T d) $p, \frac{T}{2}$

154. Four molecules of a gas have speeds 1, 2, 3 and 4 km s^{-1} . The value of rms speed of the gas molecules is

- a) $\frac{1}{2}\sqrt{15}\text{ km s}^{-1}$ b) $\frac{1}{2}\sqrt{10}\text{ km s}^{-1}$ c) 2.5 km s^{-1} d) $\sqrt{\frac{15}{2}}\text{ km s}^{-1}$


155. A body cools from 50°C to 40°C in 5 min. Its temperature comes down to 33.33°C in next 5 min. The temperature of surroundings is

- a) 15°C b) 20°C c) 25°C d) 10°C

156. Which of the following statements is true

- a) Absolute zero degree temperature is not zero energy temperature
 b) Two different gases at the same temperature pressure have equal root mean square velocities
 c) The root mean square speed of the molecules of different ideal gases, maintained at the same temperature are the same
 d) Given sample of 1 cc of hydrogen and 1 cc of oxygen both at NTP; oxygen sample has a large number of molecules

157. The figure below shows the plot of $\frac{pV}{nT}$ versus p for oxygen gas at two different temperatures.



Read the following statements concerning the above curves.

I. The dotted line corresponds to the ideal gas behavior

II. $T_1 > T_2$

III. The value of $\frac{pV}{nT}$ at the point where the curves meet on the y -axis is the same for all gases.

a) (i) only b) (i) and (ii) only c) All of these d) None of these

158. The absolute temperature of a gas is determined by

a) The average momentum of the molecules b) The velocity of sound in the gas
c) The number of molecules in the gas d) The mean square velocity of the molecules

159. If V_H, V_N and V_O denote the root-mean square velocities of molecules of hydrogen, nitrogen and oxygen respectively at a given temperature, then

a) $V_N > V_O > V_H$ b) $V_H > V_N > V_O$ c) $V_O = V_N = V_H$ d) $V_O > V_H > V_N$

160. Air inside a closed container is saturated with water vapour. The air pressure is p and the saturated vapour pressure of water is \dot{p} . If the mixture is compressed to one half of its volume by maintaining temperature constant, the pressure becomes

a) $2(p + \dot{p})$ b) $(2p + \dot{p})$ c) \dot{p} d) $p + 2\dot{p}$

161. The average kinetic energy of a gas molecule can be determined by knowing

a) The number of molecules in the gas b) The pressure of the gas only
c) The temperature of the gas only d) None of the above is enough by itself


162. Volume, pressure and temperature of an ideal gas are V, P and T respectively. If mass of its molecule is m , then its density is [$k = \dot{c}$ boltzmann's constant]

a) mkT b) $\frac{P}{kT}$ c) $\frac{P}{kTV}$ d) $\frac{Pm}{kT}$

163. One kg of a diatomic gas is at a pressure of $8 \times 10^4 \text{ Nm}^{-2}$. The density of the gas is 4 kg m^{-3} . What is the energy of the gas due to its thermal motion?

a) $3 \times 10^4 \text{ J}$ b) $5 \times 10^4 \text{ J}$ c) $6 \times 10^4 \text{ J}$ d) $7 \times 10^4 \text{ J}$

164. Graph between volume and temperature for a gas is shown in figure. If $\alpha = \dot{c}$ volume coefficient of gas $\dot{c} \frac{1}{273} \text{ per } ^\circ\text{C}$, then what is the volume of gas at a temperature of 819°C



a) $1 \times 10^{-3} \text{ m}^3$ b) $2 \times 10^{-3} \text{ m}^3$ c) $3 \times 10^{-3} \text{ m}^3$ d) $4 \times 10^{-3} \text{ m}^3$

165. A lead bullet of 10 g travelling at 300 m s^{-1} strikes against a block of wood comes to rest. Assuming 50% of heat is absorbed by the bullet, the increase in its temperature is (Specific heat of lead = 150 J kg K^{-1})


a) 100°C b) 125°C c) 150°C d) 200°C

166. When the pressure on 1200 ml of a gas is increased from 70 cm to 120 cm of mercury at constant temperature, the new volume of the gas will be

a) 700 ml b) 600 ml c) 500 ml d) 400 ml

167. At constant temperature on increasing the pressure of a gas by 5% its volume will decrease by

a) 5% b) 5.26% c) 4.26% d) 4.76%

168. The average kinetic energy of a helium atom at 30°C is
- a) Less than 1 eV b) A few keV c) $50-60\text{ eV}$ d) 13.6 eV
169. A diatomic gas is heated at constant pressure. What fraction of the heat energy is used to increase the thermal energy
- a) $3/5$ b) $3/7$ c) $5/7$ d) $5/9$
170. The molecules of a given mass of a gas have a rms velocity of 200 m/s at 27°C and $1.0 \times 10^5\text{ N/m}^2$ pressure. When the temperature is 127°C and pressure is $0.5 \times 10^5\text{ N/m}^2$, the rms velocity in m/s will be
- a) $\frac{100\sqrt{2}}{3}$ b) $100\sqrt{2}$ c) $\frac{400}{\sqrt{3}}$ d) None of these
171. Three perfect gases at absolute temperature T_1, T_2, T_3 are mixed. The masses of molecules are m_1, m_2, m_3 and the number of molecules are n_1, n_2, n_3 respectively. Assuming no loss of energy, the final temperature of the mixture is
- a) $\frac{n_1T_1+n_2T_2+n_3T_3}{n_1+n_2+n_3}$ b) $\frac{n_1T_1^2+n_2T_2^2+n_3T_3^2}{n_1T_1+n_2T_2+n_3T_3}$
c) $\frac{n_1^2T_1^2+n_2^2T_2^2+n_3^2T_3^2}{n_1T_1+n_2T_2+n_3T_3}$ d) $\frac{T_1+T_2+T_3}{3}$
172. The density of a substance at 0°C is 10 g/cc and at 100°C , its density is 9.7 g/cc . The coefficient of linear expansion of the substance is
- a) $10^{-4}^\circ\text{C}^{-1}$ b) $10^{-2}^\circ\text{C}^{-1}$ c) $10^{-3}^\circ\text{C}^{-1}$ d) 10^2°C^{-1}
173. Molecular motion shows itself as
- a) Temperature b) Internal Energy c) Friction d) Viscosity
174. Three rods made of same material and having same cross-section have been joined as shown in figure. Each rod is of same length. The left and right ends are kept at 0°C and 90°C respectively. The temperature of the junction of the three rods will be
- 
- a) 45°C b) 60°C c) 30°C d) 20°C
175. An air bubble of volume 1.0 cm^3 rises from the bottom of a lake 40 m deep at a temperature of 12°C . The volume of the bubble when it reaches the surface, which is at a temperature of 35°C , will be
- a) 5.4 cm^3 b) 4.9 cm^3 c) 2.0 cm^3 d) 10.0 cm^3
176. The mean kinetic energy of a gas at 300 K is 100 J . The mean energy of the gas at 450 K is equal to
- a) 100 J b) 3000 J c) 450 J d) 150 J
177. Two identical vessels A and B with frictionless pistons contain the same ideal gas at the same temperature and the same volume V . The masses of gas in A and B are m_A, m_B respectively. The gases are allowed to expand isothermally to same final volume $2V$. The change in pressures of the gas in A and B are found to be $\Delta p, 1.5\Delta p$ respectively. Then
- a) $9m_A=4m_B$ b) $3m_A=2m_B$ c) $2m_A=3m_B$ d) $4m_A=9m_B$
178. The identical square rods of metal are welded end to end as shown in figure, $Q\text{ cal}$ of heat flow through this combination in 4 min . If the rods were welded as shown in figure, the same amount of heat will flow through the combination in



(a)



(b)

- a) 16 min b) 12 min c) 1 min d) 4 min

179. A steel ball of mass 0.1 kg falls freely from a height of 10 m and bounces to a height of 5.4 m from the ground. If the dissipated energy in this process is absorbed by the ball, the rise in its temperature is

- a) 0.01 °C b) 0.1 °C c) 1.1 °C d) 1 °C

180. The ratio of the vapour densities of two gases at a given temperature is 9:8. The ratio of the rms velocities of their molecules is

- a) $3:2\sqrt{2}$ b) $2\sqrt{2}:3$ c) 9:8 d) 8:9

181. The r. m. s. velocity of a gas at a certain temperature is $\sqrt{2}$ times than that of the oxygen molecules at that temperature. The gas can be

- a) H_2 b) He c) CH_4 d) SO_2

182. The equation of state for 5g of oxygen at a pressure p and temperature T , when occupying a volume V , will be

- a) $pV=(5/32)RT$ b) $pV=5RT$ c) $pV=(5/2)RT$ d) $pV=(5/16)RT$

183. At NTP, sample of equal volume of chlorine and oxygen is taken. Now ratio of no. of molecules is

- a) 1:1 b) 32:27 c) 2:1 d) 16:14

184. 125 ml of gas A at 0.60 atmosphere and 150 ml of gas B at 0.80 atmospheric pressure at same temperature is filled in a vessel of 1 litre volume. What will be the total pressure of mixture at the same temperature

- a) 0.140 atmosphere b) 0.120 atmosphere c) 0.195 atmosphere d) 0.212 atmosphere

185. The gas having average speed four times as that of SO_2 (molecular mass 64) is

- a) He (molecular mass 4) b) O_2 (molecular mass 32)
c) H_2 (molecular mass 2) d) CH_4 (molecular mass 16)

186. A bubble of 8 mole of helium is submerged at a certain depth in water. The temperature of water increases by 30 °C. How much heat is added approximately to helium during expansion?

- a) 4000 J b) 3000 J c) 3500 J d) 4500 J

187. In Vander Waal's equation a and b represent $\left(P + \frac{a}{V^2}\right)(V - b) = RT$

- a) Both a and b represent correction in volume
b) Both a and b represent adhesive force between molecules
c) a represents adhesive force between molecules and b correction in volume
d) a represents correction in volume and b represents adhesive force between molecules


188. The molar specific heat at constant pressure for a monoatomic gas is

- a) $\frac{3}{2}R$ b) $\frac{5}{2}R$ c) $\frac{7}{2}R$ d) $4R$

189. The rate of diffusion is

- a) Faster in solids than in liquids and gases b) Faster in liquids than in solids and gases
c) Equal to solids, liquids and gases d) Faster in gases than in liquids and solids





190. At what temperature the kinetic energy of gas molecule is half of the value at 27°C ?
- a) 13.5°C b) 150°C c) 75 K d) -123°C
191. A horizontal uniform glass tube of 100 cm length sealed at both ends contains 10 cm mercury column in the middle. The temperature and pressure of air on either side of mercury column are respectively 31°C and 76 cm of mercury. If the air column at one end is kept at 0°C and the other end at 273°C , the pressure of air which is at 0°C is (in cm of Hg)
- a) 76 b) 88.2 c) 102.4 d) 12.2
192. A pressure P -absolute temperature T diagram was obtained when a given mass of gas was heated. During the heating process from the state 1 to state 2 the volume



- a) Remained constant b) Decreased c) Increased d) Changed erratically
193. If mass of He atom is 4 times that of hydrogen atom then mean velocity of He is
- a) 2 times of H -mean value b) $1/2$ times of H -mean value
- c) 4 times of H -mean value d) Same as H -mean value
194. $r. m. s.$ velocity of nitrogen molecules at NTP is
- a) 492 m/s b) 517 m/s c) 546 m/s d) 33 m/s
195. Two gases of equal mass are in thermal equilibrium. If P_a, P_b and V_a and V_b are their respective pressure and volumes, then which relation is true
- a) $P_a \neq P_b; V_a = V_b$ b) $P_a = P_b; V_a \neq V_b$ c) $\frac{P_a}{V_a} = \frac{P_b}{V_b}$ d) $P_a V_a = P_b V_b$
196. The ratio of the molar heat capacities of a diatomic gas at constant pressure to that at constant volume is
- a) $\frac{7}{2}$ b) $\frac{3}{2}$ c) $\frac{3}{5}$ d) $\frac{7}{5}$
197. It is seen that in proper ventilation of building, windows must be opened near the bottom and the top of the walls, so as to let pass
- a) In hot near the roof and cool air out near the bottom b) Out hot air near the roof
- c) In cool air near the bottom and hot air our near the roof d) In more air
198. A vessel is partitioned in two equal halves by a fixed diathermic separator. Two different ideal gases are filled in left (L) and right (R) halves. The rms speed of the molecules in L part is equal to the mean speed of molecules in the R part. Then the ratio of the mass of a molecule in L part to that of a molecule in R part is



- a) $\sqrt{\frac{3}{2}}$ b) $\sqrt{\pi/4}$ c) $\sqrt{2/3}$ d) $3\pi/8$
199. An ideal gas is filled in a vessel, then
- a) If it is placed inside a moving train, its temperature increases
- b) Its centre of mass moves randomly

- c) Its temperature remains constant in a moving car
d) None of these
200. If one mole of a monoatomic gas ($\gamma = \frac{5}{3}$) is mixed with one mole of a diatomic gas ($\gamma = \frac{7}{5}$), the value of γ for the mixture is
a) 1.40 b) 1.50 c) 1.53 d) 3.07
201. The kinetic energy of one g-mole of a gas at normal temperature and pressure is ($R = 8.31 \text{ J/mol-K}$)
a) $0.56 \times 10^4 \text{ J}$ b) $1.3 \times 10^2 \text{ J}$ c) $2.7 \times 10^2 \text{ J}$ d) $3.4 \times 10^3 \text{ J}$
202. 1 mol of gas occupies a volume of 200 mL at 100 mm pressure. What is the volume occupied by two moles of gas at 400 mm pressure and at same temperature?
a) 50 mL b) 100 mL c) 200 mL d) 400 mL
203. The curve between absolute temperature and v_{rms}^2 is
a)  b)  c)  d) 
204. The temperature of the mixture of one mole of helium and one mole of hydrogen is increased from 0°C to 100°C at constant pressure. The amount of heat delivered will be
a) 600 cal b) 1200 cal c) 1800 cal d) 3600 cal
205. The velocity of 4 gas molecules are given by 1 km/s, 3 km/s, 5 km/s and 7 km/s. Calculate the difference between average and rms velocity.
a) 0.338 b) 0.438 c) 0.583 d) 0.683
206. A perfect gas at 27°C is heated at constant pressure to 327°C . If original volume of gas at 27°C is V then volume at 327°C is
a) V b) $3V$ c) $2V$ d) $V/2$
207. Two containers of equal volume contain the same gas at the pressure $p_1 \wedge p_2$ and absolute temperatures $T_1 \wedge T_2$ respectively. On joining the vessels, the gas reaches a common pressure p and a common temperature T . The ratio p/T is equal to
a) $\frac{p_1 T_2 + p_2 T_1}{T_1 \times T_2}$ b) $\frac{p_1 T_2 + p_2 T_1}{T_1 + T_2}$ c) $\frac{1}{2} \left[\frac{p_1 T_2 + p_2 T_1}{T_1 T_2} \right]$ d) $\frac{p_1 T_2 - p_2 T_1}{T_1 \times T_2}$
208. The kinetic energy, due to translation motion, of most of the molecules of an ideal gas at absolute temperature T is
a) kT b) k/T c) T/k d) $1/kT$
209. The latent heat of vaporization of water is 2240 J. If the work done in the process of vaporization of 1 g is 168 J, then increase in internal energy is
a) 2072 J b) 1904 J c) 2408 J d) 2240 J
210. At what temperature the rms velocity of helium molecules will be equal to that of hydrogen molecules at NTP?
a) 844 K b) 64 K c) 273°C d) 273 K
211. Which law states that effect of pressure is same for all portions
a) Pascal's law b) Gay Lussac's law c) Dalton's law d) None of these
212. A closed vessel is maintained at a constant temperature. It is first evacuated and then vapour is injected

into it continuously. The pressure of the vapour in the vessel

- a) Increases continuously
 b) First increases and then remains constant
 c) First increases and then decreases
 d) None of the above

213. An ideal gas is expanding such that $pT^2 = \text{constant}$. The coefficient of volume expansion of the gas is

- a) $\frac{1}{T}$
 b) $\frac{2}{T}$
 c) $\frac{3}{T}$
 d) $\frac{4}{T}$

214. Mean free path of gas molecule of constant temperature is inversely proportional to

- a) P
 b) V
 c) m
 d) n (number density)

215. A closed compartment containing gas is moving with some acceleration in horizontal direction. Neglect effect of gravity. Then the pressure in the compartment is

- a) Same everywhere
 b) Lower in the front side
 c) Lower in the rear side
 d) Lower in the upper side


216. At what temperature rms speed of air molecules is doubled of that at NTP?

- a) 819°C
 b) 719°C
 c) 909°C
 d) None of these

217. In the two vessels of same volume, atomic hydrogen and helium at pressure 1 atm and 2 atm are filled. If temperature of both the samples is same, then average speed of hydrogen atoms $\langle C_H \rangle$ will be related to that of helium $\langle C_{He} \rangle$ as

- a) $\langle C_H \rangle \geq \sqrt{2} \langle C_{He} \rangle$
 b) $\langle C_H \rangle \geq \langle C_{He} \rangle$
 c) $\langle C_H \rangle \geq 2 \langle C_{He} \rangle$
 d) $\langle C_H \rangle \geq \langle C_{He} \rangle > \frac{\langle C_{He} \rangle}{2}$

218. Two spherical vessel of equal volume, are connected by a narrow tube. The apparatus contains an ideal gas at one atmosphere and 300 K . Now if one vessel is immersed in a bath of constant temperature 600 K and the other in a bath of constant temperature 300 K . Then the common pressure will be



- a) 1 atm
 b) $\frac{4}{5} \text{ atm}$
 c) $\frac{4}{3} \text{ atm}$
 d) $\frac{3}{4} \text{ atm}$

219. At constant volume the specific heat of a gas is $\frac{3R}{2}$, then the value of ' γ ' will be

- a) $\frac{3}{2}$
 b) $\frac{5}{2}$
 c) $\frac{5}{3}$
 d) None of the above

220. Gas at a pressure P_0 in contained is a vessel. If the masses of all the molecules are halved and their speeds are doubled, the resulting pressure P will be equal to

- a) $4 P_0$
 b) $2 P_0$
 c) P_0
 d) $\frac{P_0}{2}$

221. The translational kinetic energy of gas molecule for one mole of the gas is equal to

- a) $\frac{3}{2} RT$
 b) $\frac{2}{3} RT$
 c) $\frac{1}{2} RT$
 d) $\frac{2}{3} KT$

222. The product of the pressure and volume of an ideal gas is

- a) A constant
 b) Approx. equal to the universal gas constant
 c) Directly proportional to its temperature
 d) Inversely proportional to its temperature

223. The diameter of oxygen atom is 3 \AA . The fraction of molecular volume to the actual volume occupied by oxygen


at STP is

- a) 6×10^{-28} b) 8×10^{-4} c) 4×10^{-10} d) 4×10^{-4}

224. A gas is allowed to expand isothermally. The root mean square velocity of the molecules

- a) Will increase b) Will decrease
c) Will remain unchanged d) Depends on the other factors

225. Two different isotherms representing the relationship between pressure p and volume V at a given temperature of the same ideal gas are shown for masses m_1 and m_2 of the gas respectively in the figure given, then



- a) $m_1 > m_2$ b) $m_1 = m_2$ c) $m_1 < m_2$ d) $m_1 = \frac{1}{2} m_2$

226. At 100 K and 0.1 atmospheric pressure, the volume of helium gas is 10 litres. If volume and pressure are doubled, its temperature will change to

- a) 400 K b) 127 K c) 200 K d) 25 K

227. Two balloons are filled, one with pure He gas and the other by air, respectively. If the pressure and temperature of these balloons are same, then the number of molecules per unit volume is

- a) More in the He filled balloon b) Same in both balloons
c) More in air filled balloon d) In the ratio of 1:4

228. If the rms velocity of a gas is v , then

- a) $v^2 T = \text{constant}$ b) $v^2 / T = \text{constant}$
c) $v T^2 = \text{constant}$ d) v is independent of T

229. The ratio of two specific heats $\frac{C_p}{C_v}$ of CO is

- a) 1.33 b) 1.40 c) 1.29 d) 1.66

230. A gas is filled in a closed container and its molecules are moving in horizontal direction with uniform acceleration. Neglecting acceleration due to gravity, the pressure inside the container is

- a) Uniform everywhere b) Less in the front
c) Less at the back d) Less at the top

231. A closed gas cylinder is divided into two parts by a piston held tight. The pressure and volume of gas in two parts respectively are $(P, 5V)$ and $(10P, V)$. If now the piston is left free and the system undergoes isothermal process, then the volume of the gas in two parts respectively are


- a) $2V, 4V$ b) $3V, 3V$ c) $5V, V$ d) $4V, 2V$

232. On colliding in a closed container the gas molecules

- a) Transfer momentum to the walls b) Momentum becomes zero
c) Move in opposite directions d) Perform Brownian motion

233. A sealed container with negligible coefficient of volumetric expansion contains helium (a monoatomic gas). When it is heated from 300 K to 600 K, the average K.E. of helium atoms is

- a) Halved b) Unchanged


- c) Doubled
d) Increased by factor $\sqrt{2}$
234. A monoatomic gas is kept at room temperature 300 K . Calculate the average kinetic energy of gas molecule (Use $k = 1.38 \times 10^{-23}\text{ MKS units}$)
a) 0.138 eV b) 0.062 eV c) 0.039 eV d) 0.013 eV
235. When the temperature of a gas increases by 1°C , its pressure increases 0.4% . What is its initial temperature?
a) 250 K b) 125 K c) 195 K d) 329 K
236. A bubble is at the bottom of the lake of depth h . As the bubble comes to sea level, its radius increases three times. If atmospheric pressure is equal to 1 metre of water column, then h is equal to
a) 26 l b) 1 c) 25 l d) 30 l
237. A diatomic gas molecule has translational, rotational and vibrational degrees of freedom. The C_p/C_v is
a) 1.67 b) 1.4 c) 1.29 d) 1.33
238. In the absence of intermolecular forces of attraction, the observed pressure p will be
a) p b) $i p$ c) $i p$ d) Zero
239. At 0 K which of the following properties of a gas will be zero
a) Kinetic energy b) Potential energy c) Vibrational energy d) Density
240. The equation for an ideal gas is $PV = RT$, where V represents the volume of
a) 1 g gas b) Any mass of the gas c) One g mol gas d) One litre gas
241. A gas at 27°C has a volume V and pressure P . On heating its pressure is doubled and volume becomes three times. The resulting temperature of the gas will be
a) 1800°C b) 162°C c) 1527°C d) 600°C
242. The figure shows the volume V versus temperature T graphs for a certain mass of a perfect gas at two constant pressures of P_1 and P_2 . What inference can you draw from the graphs
- 
- a) $P_1 > P_2$ b) $P_1 < P_2$
c) $P_1 = P_2$ d) No inference can be drawn due to insufficient information
243. For hydrogen gas $C_p - C_v = a$ and for oxygen gas $C_p - C_v = b$. So the relation between a and b is given by
a) $a = 16b$ b) $b = 16a$ c) $a = 4b$ d) $a = b$
244. For a real gas (van der Waal's gas)
a) Boyle temperature is a/Rb
b) Critical temperature is a/Rb
c) Triple temperature is $2a/Rb$
d) Inversion temperature is $a/2Rb$
245. According to the kinetic theory of gases the $r.m.s.$ velocity of gas molecules is directly proportional to

- a) T b) \sqrt{T} c) T^2 d) $1/\sqrt{T}$
246. Root mean square velocity of a particle is v at pressure P . If pressure is increased two times, then the *r. m. s.* velocity becomes
a) $2v$ b) $3v$ c) $0.5v$ d) v
247. The average translational kinetic energy of a hydrogen gas molecules at NTP will be [Boltzmann's constant $k_B = 1.38 \times 10^{-23} \text{ J/K}$]
a) $0.186 \times 10^{-20} \text{ Joule}$ b) $0.372 \times 10^{-20} \text{ Joule}$ c) $0.56 \times 10^{-20} \text{ Joule}$ d) $5.6 \times 10^{-20} \text{ Joule}$
248. The efficiency of a Carnot engine is 50% and temperature of sink is 500 K. If temperature of source is kept constant and its efficiency raised to 60%, then the required temperature of sink will be
a) 100 K b) 600 K c) 400 K d) 500 K
249. The temperature of a given mass is increased from 27°C to 327°C . The rms velocity of the molecules increases
a) $\sqrt{2} \times i$ b) 2 times c) $2\sqrt{2} \times i$ d) 4 times
250. A real gas behaves like an ideal gas if its
a) Pressure and temperature are both high b) Pressure and temperature are both low
c) Pressure is high and temperature is low d) Pressure is low and temperature is high
251. A gas mixture consists of 2 moles of oxygen and 4 moles of argon at temperature T . Neglecting all vibrational moles, the total internal energy of the system is
a) $4RT$ b) $15RT$ c) $9RT$ d) $11RT$
252. Six moles of O_2 gas is heated from 20°C to 35°C at constant volume. If specific heat capacity at constant pressure is $8 \text{ cal mol}^{-1} - \text{K}^{-1}$ and $R = 8.31 \text{ J mol}^{-1} - \text{K}^{-1}$, what is change in internal energy of gas?
a) 180 cal b) 300 cal c) 360 cal d) 540 cal
253. Read the given statements and decide which is/are correct on the basis of kinetic theory of gases
(I) Energy of one molecule at absolute temperature is zero
(II) *r. m. s.* speeds of different gases are same at same temperature
(III) For one gram of all ideal gas kinetic energy is same at same temperature
(IV) For one mole of all ideal gases mean kinetic energy is same at same temperature
a) All are correct b) I and IV are correct c) IV is correct d) None of these
254. A perfect gas at 27°C is heated at constant pressure so as to double its volume. The increase in temperature of the gas will be
a) 300°C b) 54°C c) 327°C d) 600°C
255. Cooking gas containers are kept in a lorry moving with uniform speed. The temperature of the gas molecules inside will
a) Increase b) Decrease
c) Remain same d) Decrease for some, while increase for others
256. The root mean square speed of the molecules of a gas is
a) Independent of its pressure but directly proportional to its Kelvin temperature
b) Directly proportional to the square roots of both its pressure and its Kelvin temperature
c) Independent of its pressure but directly proportional to the square root of its Kelvin temperature
d) Directly proportional to both its pressure and its kelvin temperature
257. The mean kinetic energy of one mole of gas per degree of freedom (on the basis of kinetic theory of

gases) is

- a) $\frac{1}{2}kT$ b) $\frac{3}{2}kT$ c) $\frac{3}{2}RT$ d) $\frac{1}{2}RT$

258. Two different masses m and $3m$ of an ideal gas are heated separately in a vessel of constant volume, the pressure P and absolute temperature T , graphs for these two cases are shown in the figure as A and B . The ratio of slopes of curves B to A is



- a) 3:1 b) 1:3 c) 9:1 d) 1:9

259. Mean kinetic energy per degree of freedom of gas molecules is

- a) $\frac{3}{2}kT$ b) kT c) $\frac{1}{2}kT$ d) $\frac{3}{2}RT$

260. 22 g of carbon dioxide at 27°C is mixed in a closed container with 16 g of oxygen at 37°C . If both gases are considered as ideal gases, then the temperature of the mixture is

- a) 24.2°C b) 28.5°C c) 31.5°C d) 33.5°C

261. 70 cal of heat are required to raise the temperature of 2 mole of an ideal gas at constant pressure from 30°C to 35°C . The amount of heat required to raise the temperature of the same sample of the gas through the same range at constant volume is nearly (Gas constant = $1.99 \text{ cal K}^{-1} - \text{mol}^{-1}$)

- a) 30 cal b) 50 cal c) 70 cal d) 90 cal

262. Which of the following formula is wrong

- a) $C_V = \frac{R}{\gamma - 1}$ b) $C_P = \frac{\gamma R}{\gamma - 1}$ c) $C_P / C_V = \gamma$ d) $C_P - C_V = 2R$

263. Ideal gas and real gas has major difference of

- a) Phase transition b) Temperature c) Pressure d) None of them

264. If mass of He is 4 times that of hydrogen, then mean velocity of He is

- a) 2 times of H-mean value
 b) $\frac{1}{2}$ times of H-mean value
 c) 4 times of H-mean value
 d) Same as H-mean value

265. Supposing the distance between the atoms of a diatomic gas to be constant, its specific heat at constant volume per mole (gram mole) is

- a) $\frac{5}{2}R$ b) $\frac{3}{2}R$ c) R d) $\frac{1}{2}R$

266. At what temperature is the kinetic energy of a gas molecule double that of its value of 27°C

- a) 54°C b) 300 K c) 327°C d) 108°C

267. A flask of volume 10^3 cc is completely filled with mercury at 0°C . The coefficient of cubical expansion of mercury is $180 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$ and that of glass is $40 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$.

If the flask is now placed in boiling water at 100°C , how much mercury will overflow?

- a) 7 cc b) 14 cc c) 21 cc d) 28 cc

268. The pressure is exerted by the gas on the walls of the container because

- a) It loses kinetic energy
 b) It sticks with the walls
 c) On collision with the walls there is a change in momentum
 d) It is accelerated towards the walls

269. A balloon contains 500 m^3 of helium at 27°C and 1 atmosphere pressure. The volume of the helium at -3°C temperature and 0.5 atmosphere pressure will be

- a) 500 m^3
 b) 700 m^3
 c) 900 m^3
 d) 1000 m^3

270. An ideal gas has an initial pressure of 3 pressure units and an initial volume of 4 volume units. The table gives the final the final pressure and volume of the gas (in those same units) in four, processes. Which processes start and end on the same isotherm

	A	B	C	D
P	5	4	12	6
V	7	6	1	3

- a) A
 b) B
 c) C
 d) D


271. Specific heats of monoatomic and diatomic gases are same and satisfy the relation which is

- a) $C_p(\text{mono}) = C_p(\text{dia})$
 b) $C_p(\text{mono}) = C_v(\text{dia})$
 c) $C_v(\text{mono}) = C_v(\text{dia})$
 d) $C_v(\text{mono}) = C_p(\text{dia})$

272. The root mean square velocity of gas molecules at 27°C is 1365 m s^{-1} . The gas is

- a) O_2
 b) He
 c) N_2
 d) CO_2

273. A volume V and pressure P diagram was obtained from state 1 to state 2 when a given mass of a gas is subjected to temperature changes. During this process the gas is



- a) Heated continuously
 b) Cooled continuously
 c) Heated in the beginning and cooled towards the end
 d) Cooled in the beginning and heated towards the end

274. At constant pressure, which of the following is true?

- a) $v \propto \sqrt{\rho}$
 b) $v \propto \frac{1}{\rho}$
 c) $v \propto \rho$
 d) $v \propto \frac{1}{\sqrt{\rho}}$

275. A vessel contains 32 g of O_2 at a temperature T . The pressure of the gas is p . An identical vessel containing 4 g of H_2 at a temperature $2T$ has a pressure of

- a) $8p$
 b) $4p$
 c) p
 d) $\frac{p}{8}$

276. Root mean square speed of the molecules of ideal gas is v . If pressure is increased two times at constant temperature, the rms speed will become

- a) $\frac{v}{2}$
 b) v
 c) $2v$
 d) $4v$

277. Relationship between P , V , and E for a gas is

- a) $P = \frac{3}{2}EV$
 b) $V = \frac{2}{3}EP$
 c) $PV = \frac{3}{2}E$
 d) $PV = \frac{2}{3}E$

278. The specific heat relation for ideal gas is

- a) $C_p + C_v = R$ b) $C_p - C_v = R$ c) $C_p / C_v = R$ d) $C_v / C_p = R$
279. The temperature of an ideal gas is increased from 27°C to 127°C , then percentage increase in V_{rms} is
a) 37% b) 11% c) 33% d) 15.5%
280. The coefficient of apparent expansion of a liquid when determined using two different vessels A and B are λ_1 and λ_2 , respectively. If the coefficient of linear expansion of the vessel A is α , the coefficient of linear expansion of vessel B is
a) $\frac{\alpha \lambda_1 \lambda_2}{\lambda_1 + \lambda_2}$ b) $\frac{\lambda_1 - \lambda_2}{2\alpha}$ c) $\frac{\lambda_1 - \lambda_2 + \alpha}{3\alpha}$ d) $\frac{\lambda_1 - \lambda_2}{3} + \alpha$
281. A steel tape measures the length of a copper rod as 90.0 cm, when both are at 10°C , the calibration temperature, for the tape. What would be tape read for the length of the rod when both are at 30°C . Given, α for steel $1.2 \times 10^{-5} \text{ }^\circ\text{C}^{-1}$ and α for copper is $1.7 \times 10^{-5} \text{ }^\circ\text{C}^{-1}$.
a) 90.01 cm b) 89.90 cm c) 90.22 cm d) 89.80 cm
282. According to the kinetic theory of gases, at absolute temperature
a) Water freezes b) Liquid helium freezes
c) Molecular motion stops d) Liquid hydrogen freezes
283. At the same temperature and pressure and volume of two gases, which of the following quantities is constant
a) Total number of molecules b) Average kinetic energy
c) Root mean square velocity d) Mean free path
284. A diatomic molecule has how many degrees of freedom
a) 3 b) 4 c) 5 d) 6
285. For a gas if $\gamma = 1.4$, then atomicity, C_p and C_v of the gas are respectively
a) Monoatomic, $\frac{5}{2}R, \frac{3}{2}R$ b) Monoatomic, $\frac{7}{2}R, \frac{5}{2}R$ c) Diatomic, $\frac{7}{2}R, \frac{3}{2}R$ d) Triatomic, $\frac{7}{2}R, \frac{5}{2}R$
286. Simple behaviour under all conditions of real gas is governed by the equation
a) $PV = \mu RT$ b) $\left(P + \frac{a}{V^2}\right)(V - b) = \mu RT$
c) $PV = \mu \text{ constant}$ d) $PV^\gamma = \mu \text{ constant}$
287. A metal ball immersed in water weighs w_1 at 0°C and w_2 at 50°C . The coefficient of cubical expansion of metal is less than that of water. Then
a) $w_1 < w_2$ b) $w_1 > w_2$ c) $w_1 = w_2$ d) Data is not sufficient
288. If the volume of the gas containing n number of molecules is V , then the pressure will decrease due to force of intermolecular attraction in the proportion
a) n/V b) n/V^2 c) $(n/V)^2$ d) $1/V^2$
289. The temperature of an ideal gas at atmospheric pressure is 300 K and volume 1 m^3 . If temperature and volume become double, then pressure will be
a) 10^5 N/m^2 b) $2 \times 10^5\text{ N/m}^2$ c) $0.5 \times 10^5\text{ N/m}^2$ d) $4 \times 10^5\text{ N/m}^2$
290. Volume of gas becomes four times if
a) Temperature becomes four times at constant pressure

- b) Temperature becomes one fourth at constant pressure
- c) Temperature becomes two times at constant pressure
- d) Temperature becomes half at constant pressure

291. The root mean square velocity of a gas molecule of mass m at a given temperature is proportional to

- a) m^0
- b) m
- c) \sqrt{m}
- d) $\frac{1}{\sqrt{m}}$

292. If universal gas constant is R , the essential heat to increase from 273 K to 473 K at constant volume for ideal gas of 4 mol is

- a) $200 R$
- b) $400 R$
- c) $800 R$
- d) $1200 R$

293. Universal gas constant is

- a) $\frac{C_p}{C_v}$
- b) $C_p - C_v$
- c) $C_p + C_v$
- d) $\frac{C_v}{C_p}$

294. 22 g of CO_2 at $27^\circ C$ is mixed with 16 g of oxygen at $37^\circ C$. The temperature of the mixture is

- a) $32^\circ C$
- b) $27^\circ C$
- c) $37^\circ C$
- d) $30^\circ C$

295. The molecular weight of a gas is 44. The volume occupied by 2.2 g of this gas at $0^\circ C$ and $2 atm$. pressure will be

- a) 0.56 litre
- b) 1.2 litres
- c) 2.4 litres
- d) 5.6 litres

296. A box contains n molecules of a gas. How will the pressure of the gas be effected, if the number of molecules is made $2n$

- a) Pressure will decrease
- b) Pressure will remain unchanged
- c) Pressure will be doubled
- d) Pressure will become three times

297. Consider a gas with density ρ and \dot{c} as the root mean square velocity of its molecules contained in a volume. If the system moves as whole with velocity v , then the pressure exerted by the gas is

- a) $\frac{1}{3} \rho \dot{c}^2$
- b) $\frac{1}{3} \rho (c+v)^2$
- c) $\frac{1}{3} \rho (\dot{c} - v)^2$
- d) $\frac{1}{3} \rho (c^2 - v)^2$

298. At constant volume, temperature is increased. Then

- a) Collision on walls will be less
- b) Number of collisions per unit time will increase
- c) Collisions will be in straight lines
- d) Collisions will not change


299. One mole of an ideal gas requires 207 J heat to raise the temperature by 10 K when heated at constant pressure. If the same gas is heated at constant volume to raise the temperature by the same 10 K, the heat required is (Given the gas constant $R=8.3 J/mol-K$)

- a) 198.7 J
- b) 29 J
- c) 215.3 J
- d) 124 J


300. At what temperature volume of an ideal gas at $0^\circ C$ becomes triple

- a) $546^\circ C$
- b) $182^\circ C$
- c) $819^\circ C$
- d) $646^\circ C$

301. An air bubble doubles its radius on raising from the bottom of water reservoir to be the surface of water in it. If the atmospheric pressure is equal to 10 m of water, the height of water in the reservoir is



- a) 10 m b) 20 m c) 70 m d) 80 m
302. A cylinder of 5 litre capacity, filled with air at N.T.P. is connected with another evacuated cylinder of 30 litres of capacity. The resultant air pressure in both the cylinders will be
a) 38.85 cm of Hg b) 21.85 cm of Hg c) 10.85 cm of Hg d) 14.85 cm of Hg
303. In gases of diatomic molecules, the ratio of the two specific heats of gases C_p/C_v is
a) 1.66 b) 1.40 c) 1.33 d) 1.00
304. Oxygen boils at (-183°C) . The temperature on the Fahrenheit scale is
a) -297.4°F b) -253.6°F c) -342.6°F d) -225.3°F
305. The specific heats at constant pressure is greater than that of the same gas at constant volume because
a) At constant pressure work is done in expanding the gas
b) At constant volume work is done in expanding the gas
c) The molecular attraction increases more at constant pressure
d) The molecular vibration increases more at constant pressure
306. A type kept outside in sunlight bursts off after sometime because of
a) Increases in pressure b) Increases in volume c) Both (a) and (b) d) None of these
307. 10 moles of an ideal monoatomic gas at 10°C is mixed with 20 moles of another monoatomic gas at 20°C . Then the temperature of the mixture is
a) 15.5°C b) 15°C c) 16°C d) 16.6°C
308. The number of translational degrees of freedom for a diatomic gas is
a) 2 b) 3 c) 5 d) 6
309. Let A and B the two gases and given $\frac{T_B}{M_A} = 4 \cdot \frac{T_B}{M_B}$; where T is the temperature and M is molecular mass. If C_A and C_B are the *r.m.s.* speed, then the ratio $\frac{C_A}{C_B}$ will be equal to
a) 2 b) 4 c) 1 d) 0.5
310. The value of C_v for one mole of neon gas is
a) $\frac{1}{2}R$ b) $\frac{3}{2}R$ c) $\frac{5}{2}R$ d) $\frac{7}{2}R$
311. Two spheres made of same substance have diameters in the ratio 1 : 2. Their thermal capacities are in the ratio of
a) 1 : 2 b) 1 : 8 c) 1 : 4 d) 2 : 1
312. For an ideal gas
a) C_p is less than C_v b) C_p is equal to C_v
c) C_p is greater than C_v d) $C_p = C_v = 0$
313. From the following $P-T$ graph what inference can be drawn



- a) $V_2 > V_1$ b) $V_2 < V_1$ c) $V_2 = V_1$ d) None of the above

314. Some gas at 300 K is enclosed in a container. Now, the container is placed on a fast moving train. While the train is in motion, the temperature of the gas

- a) Rises above 300 K b) Falls below 300 K
c) Remains unchanged d) Become unsteady

315. According to Maxwell's law of distribution of velocities of molecules, the most probable velocity is

- a) Greater than the mean velocity b) Equal to the mean velocity
c) Equal to the root mean square velocity d) Less than the root mean square velocity


316. If C_p and C_v denote the specific heats of nitrogen per unit mass at constant pressure and constant volume respectively, then

- a) $C_p - C_v = R/28$ b) $C_p - C_v = R/14$ c) $C_p - C_v = R$ d) $C_p - C_v = 28 R$

317. A cubical box with porous walls containing an equal number of O_2 and H_2 molecules is placed in a large evacuated chamber. The entire system is maintained at constant temperature T . The ratio of v_{rms} of O_2 molecules to that of the v_{rms} of H_2 molecules, found in the chamber outside the box after a short interval is

- a) $\frac{1}{2\sqrt{2}}$ b) $\frac{1}{4}$ c) $\frac{1}{\sqrt{2}}$ d) $\sqrt{2}$

318. The graph which represents the variation of mean kinetic energy of molecules with temperature $t^\circ C$ is



319. Boyle's law holds for an ideal gas during

- a) Isobaric changes b) Isothermal changes c) Isochoric changes d) Isotonic changes

320. The kinetic energy of one mole gas at 300 K temperature, is E . At 400 K temperature kinetic energy is E' . The value of E'/E is

- a) 1.33 b) $\sqrt{\left(\frac{4}{3}\right)}$ c) $\frac{16}{9}$ d) 2

321. Saturated vapour is compressed to half its volume without any change in temperature, then the pressure will be

- a) Doubled b) Halved c) The same d) Zero

322. The amount of heat required to convert 10 g of ice at $-10^\circ C$ into steam at $100^\circ C$ is (in calories)

- a) 6400 b) 5400 c) 7200 d) 7250

323. Inside a cylinder closed at both ends is a movable piston. On one side of the piston is a mass m of a gas, and on the other side a mass $2m$ of the same gas. What fraction of the volume of the cylinder will be occupied by the larger mass of the gas when the piston is in equilibrium? The temperature is the same throughout.

- a) $\frac{2}{3}$ b) $\frac{1}{3}$ c) $\frac{1}{2}$ d) $\frac{1}{4}$

324. O_2 gas is filled in a vessel. If pressure is doubled, temperature becomes four times, how many times its density

will become

- a) 2 b) 4 c) $\frac{1}{4}$ d) $\frac{1}{2}$


325. The ratio of mean kinetic energy of hydrogen and oxygen at a given temperature is

- a) 1:16 b) 1:8 c) 1:4 d) 1:1


326. For matter to exist simultaneously in gas and liquid phases

- a) The temperature must be 0 K
b) The temperature must be less than $0\text{ }^\circ\text{C}$
c) The temperature must be less than the critical temperature
d) The temperature must be less than the reduced temperature

327. Which one of the following graphs represents the behaviour of an ideal gas?



328. Pressure versus temperature graph of an ideal gas at constant volume V of an ideal gas is shown by the straight line A. Now mass of the gas is doubled and the volume is halved, then the corresponding pressure versus temperature graph will be shown by the line



- a) A b) B c) C d) None of these

329. If a Vander-Waal's gas expands freely, then final temperature is

- a) Less than the initial temperature
b) Equal to the initial temperature
c) More than the initial temperature
d) Less or more than the initial temperature depending on the nature of the gas

330. Oxygen and hydrogen are at the same temperature T . The ratio of the mean kinetic energy of oxygen molecules to that of the hydrogen molecules will be

- a) 16:1 b) 1:1 c) 4:1 d) 1:4

331. At temperature T , the *r.m.s.* speed of helium molecules is the same as *r.m.s.* speed of hydrogen molecules at normal temperature and pressure. The value of T is

- a) $273\text{ }^\circ\text{C}$ b) $546\text{ }^\circ\text{C}$ c) $0\text{ }^\circ\text{C}$ d) $136.5\text{ }^\circ\text{C}$

332. The pressure and temperature of an ideal gas in a closed vessel are 720 kPa and 40°C respectively. If $\frac{1}{4}$ th of the gas is released from the vessel and the temperature of the remaining gas is raised to 353°C , the final pressure of the gas is
 a) 1440 kPa b) 1080 kPa c) 720 kPa d) 540 kPa
333. A cylinder of fixed capacity (of 44.8 litres) contains 2 moles of helium gas at STP. What is the amount of heat needed to raise the temperature of the gas in the cylinder by 20°C (Use $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$)
 a) 996 J b) 831 J c) 498 J d) 374 J
334. A thin copper wire of length l increase in length by 1% , when heated from 0°C to 100°C . If a thin copper plate of area $2l \times l$ is heated from 0°C to 100°C , the percentage increase in its area would be
 a) 1% b) 4% c) 3% d) 2%
335. The *r.m.s.* speed of the molecules of a gas at a pressure 10^5 Pa and temperature 0°C is 0.5 km sec^{-1} . If the pressure is kept constant but temperature is raised to 819°C , the velocity will become
 a) 1.5 km s^{-1} b) 2 km s^{-1} c) 5 km s^{-1} d) 1 km s^{-1}
336. For one gram mol of a gas, the value of R in the equation $PV = RT$ is nearly
 a) 2 cal/K b) 10 cal/K c) 0.2 cal/K d) 200 cal/K
337. A solid whose volume does not change with temperature floats in liquid. For two different temperatures t_1 and t_2 , the fractions f_1 and f_2 of volume of solid remain submerged. What is the coefficient of volume expansion of liquid?
 a) $\frac{f_1 - f_2}{f_2 t_1 - f_1 t_2}$ b) $\frac{f_1 - f_2}{f_1 t_1 - f_2 t_2}$ c) $\frac{f_1 + f_2}{f_2 t_1 - f_1 t_2}$ d) $\frac{f_1 + f_2}{f_1 t_1 - f_2 t_2}$
338. Find the ratio of specific heat at constant pressure to the specific heat constant volume for NH_3
 a) 1.33 b) 1.44 c) 1.28 d) 1.67
339. What is the ratio of specific heats of constant pressure and constant volume for NH_3
 a) 1.33 b) 1.44 c) 1.28 d) 1.67
340. For a gas molecule with 6 degrees of freedom the law of equipartition of energy gives the following relation between the molecular specific heat (C_v) and gas constant (R)
 a) $C_v = \frac{R}{2}$ b) $C_v = R$ c) $C_v = 2R$ d) $C_v = 3R$
341. A polyatomic gas with n degrees of freedom has a mean energy per molecule given by (N is Avogadro's number)
 a) $\frac{nkT}{N}$ b) $\frac{nkT}{2N}$ c) $\frac{nkT}{2}$ d) $\frac{3kT}{2}$
342. If the mean free path of atoms is doubled then the pressure of gas will become
 a) $P/4$ b) $P/2$ c) $P/8$ d) P
343. The temperature of a gas is -68°C . At what temperature will the average kinetic energy of its molecules be twice that of at -68°C ?
 a) 137°C b) 127°C c) 100°C d) 105°C
344. For a gas $\frac{R}{C_v} = 0.67$. This gas is made up of molecules which are
 a) Diatomic b) Mixture of diatomic and polyatomic molecules
 c) Monoatomic d) Polyatomic
345. The specific heat of a gas

- d) The collisions amongst the molecules are of short duration
358. At what temperature, the mean kinetic energy of O_2 will be the same for H_2 molecules at $-73^\circ C$
- a) $127^\circ C$ b) $527^\circ C$ c) $-73^\circ C$ d) $-173^\circ C$
359. The relation between two specific heats of a gas is
- a) $C_p - C_v = \frac{R}{J}$ b) $C_v - C_p = \frac{R}{J}$ c) $C_p - C_v = J$ d) $C_v - C_p = J$
360. One mole of a monoatomic ideal gas is mixed with one mole of a diatomic ideal gas. The molar specific heat of the mixture at constant volume is
- a) $(3/2)R$ b) $(5/2)R$ c) $2R$ d) $4R$
361. Two moles of monoatomic gas is mixed with three moles of a diatomic gas. The molar specific heat of the mixture at constant volume is
- a) $1.55R$ b) $2.10R$ c) $1.63R$ d) $2.20R$
362. In the relation $n = \frac{PV}{RT}$, $n =$
- a) Number of molecules b) Atomic number c) Mass number d) Number of moles
363. The root mean square speed of hydrogen molecules of an ideal hydrogen gas kept in a gas chamber at $0^\circ C$ is $3180 \text{ metres/second}$. The pressure on the hydrogen gas is (Density of hydrogen gas is $8.99 \times 10^{-2} \text{ kg/m}^3$, $1 \text{ atmosphere} = 1.01 \times 10^5 \text{ N/m}^2$)
- a) 1.0 atm b) 1.5 atm c) 2.0 atm d) 3.0 atm
364. Pressure of an ideal gas is increased by keeping temperature constant. What is the effect on kinetic energy of molecules?
- a) Increases b) Decrease
- c) No change d) Can't be determined
365. The volume of a gas at $20^\circ C$ is 200 ml . If the temperature is reduced to $-20^\circ C$ at constant pressure, its volume will be
- a) 172.6 ml b) 17.26 ml c) 192.7 ml d) 19.27 ml
366. At $0^\circ C$ the density of a fixed mass of a gas divided by pressure is x . At $100^\circ C$, the ratio will be
- a) x b) $\frac{273}{373}x$ c) $\frac{373}{273}x$ d) $\frac{100}{273}x$
367. A wheel is 80.3 cm in circumference. An iron tyre measures 80.0 cm around its inner face. If the coefficient of linear expansion for iron is $12 \times 10^{-6} \text{ }^\circ C^{-1}$, the temperature of the tyre must be raised by
- a) $105^\circ C$ b) $417^\circ C$ c) $312^\circ C$ d) $223^\circ C$
368. Which one of the following is not an assumption of kinetic theory of gases?
- a) The volume occupied by the molecules of the gas is negligible
- b) The force of attraction between the molecules is negligible
- c) The collision between the molecules are elastic
- d) All molecules have same speed
369. The equation of state of a gas is given by $\left(P + \frac{aT^2}{V}\right)V^c = (RT + b)$, where a, b, c and R are constants. The isotherms can be represented by $P = AV^m - BV^n$, where A and B depend only on temperature and

- a) $m = -c$ and $n = -1$ b) $m = c$ and $n = 1$ c) $m = -c$ and $n = 1$ d) $m = c$ and $n = -1$
370. The temperature gradient in the earth's crust is 32°C km^{-1} and the mean conductivity of earth is $0.008 \text{ cal s}^{-1} \text{ cm}^{-1} \text{ }^\circ\text{C}^{-1}$. Considering earth to be a sphere of radius 6000 km loss of heat by earth everyday is about
- a) 10^{30} cal b) 10^{40} cal c) 10^{20} cal d) 10^{18} cal
371. For a gas, the *r.m.s.* speed at 800 K is
- a) Four times the value at 200 K b) Half the value at 200 K
c) Twice the value at 200 K d) Same as at 200 K
372. 8 g of O_2 , 14 g of N_2 and 22 g of CO_2 is mixed in a container of 10 L capacity at 27°C . The pressure exerted by the mixture in terms of atmospheric pressure is
($R = 0.082 \text{ L atm K}^{-1} \text{ mol}^{-1}$)
- a) 1.4 atm b) 2.5 atm c) 3.7 atm d) 8.7 atm
373. Vapour is injected at a uniform rate in a closed vessel which was initially evacuated. The pressure in the vessel
- a) Increases continuously b) Decreases continuously
c) First increases and then decreases d) First increases and then becomes constant
374. At what temperature the molecule of nitrogen will have same rms velocity as the molecule of oxygen at 127°C ?
- a) 457°C b) 273°C c) 350°C d) 77°C
375. The temperature of an ideal gas is reduced from 927°C to 27°C . The *r.m.s.* velocity of the molecules becomes
- a) Double the initial value b) Half of the initial value
c) Four times the initial value d) Ten times the initial value
376. At a given temperature the root mean square velocities of oxygen and hydrogen molecules are in the ratio
- a) 16:1 b) 1:16 c) 4:1 d) 1:4
377. The temperature of 5 moles of a gas at constant volume is changed from 100°C to 120°C . The change in internal energy is 80 J. the total heat capacity of the gas at constant volume will be in J K^{-1} is
- a) 8 b) 4 c) 0.8 d) 0.4
378. One mole of monoatomic gas and three moles of diatomic gas are put together in a container. The molar specific heat ($\text{J K}^{-1} \text{ mol}^{-1}$) at constant volume is ($R = 8.3 \text{ J K}^{-1} \text{ mol}^{-1}$)
- a) 18.7 b) 18.9 c) 19.2 d) None of these
379. If masses of all molecules of a gas are halved and their speeds are doubles, then the ratio of initial and final pressures is
- a) 1:2 b) 2:1 c) 4:1 d) 1:4
380. The molar specific heat at constant pressure of an ideal gas is $(7/2)R$. The ratio of specific heat at constant pressure to that at constant volume is
- a) $5/7$ b) $9/7$ c) $7/5$ d) $8/7$
381. The specific heat of an ideal gas is
- a) Proportional to T b) Proportional to T^2 c) Proportional to T^3 d) Independent of T
382. Speed of sound in a gas is v and *r.m.s.* velocity of the gas molecules is c . The ratio of v to c is

a) $\frac{3}{\gamma}$

b) $\frac{\gamma}{3}$

c) $\sqrt{\frac{3}{\gamma}}$

d) $\sqrt{\frac{\gamma}{3}}$

383. The molecular weights of O_2 and N_2 are 32 and 28 respectively. At $15^\circ C$, the pressure of $1g O_2$ will be the same as that of $1g N_2$ in the same bottle at the temperature

a) $-21^\circ C$

b) $13^\circ C$

c) $15^\circ C$

d) $56.4^\circ C$

384. On giving equal amount of heat at constant volume to $1 mole$ of a monoatomic and a diatomic gas the rise in temperature

a) Monoatomic

b) Diatomic

c) Same for both

d) Can not be predicted

385. The *r. m. s.* speed of gas molecules is given by

a) $2.5\sqrt{\frac{RT}{M}}$

b) $1.73\sqrt{\frac{RT}{M}}$

c) $2.5\sqrt{\frac{M}{RT}}$

d) $1.73\sqrt{\frac{M}{RT}}$

386. A sample of an ideal gas occupies a volume V at a pressure P and absolute temperature T , the mass of each molecule is m . The expression for the density of gas is ($k = \text{Boltzmann's constant}$)

a) $m k T$

b) P/kT

c) P/kTV

d) Pm/kT

387. A gaseous mixture contains equal number of hydrogen and nitrogen and nitrogen molecules. Specific heat measurements on this mixture at temperatures below $100 K$ would indicate that the of γ (ratio specific heats) for this mixture is

a) $3/2$

b) $4/3$

c) $5/3$

d) $7/5$

13.KINETIC THEORY

: ANSWER KEY :

1) d	2) a	3) c	4) b	169) c	170) c	171) a	172) a
5) a	6) c	7) b	8) c	173) a	174) b	175) a	176) d
9) c	10) b	11) c	12) a	177) a	178) c	179) b	180) b
13) c	14) b	15) a	16) c	181) c	182) a	183) a	184) c
17) a	18) a	19) a	20) d	185) a	186) b	187) c	188) b
21) c	22) d	23) d	24) c	189) d	190) d	191) c	192) c
25) d	26) c	27) d	28) c	193) b	194) b	195) d	196) d
29) d	30) b	31) a	32) b	197) c	198) d	199) c	200) c
33) c	34) b	35) a	36) a	201) d	202) b	203) b	204) b
37) c	38) c	39) d	40) c	205) c	206) c	207) c	208) a
41) d	42) a	43) b	44) c	209) a	210) c	211) a	212) b
45) a	46) a	47) c	48) c	213) c	214) d	215) b	216) a
49) d	50) d	51) b	52) d	217) c	218) c	219) c	220) b
53) c	54) c	55) d	56) a	221) a	222) c	223) b	224) c
57) c	58) d	59) b	60) d	225) c	226) a	227) b	228) b
61) a	62) b	63) b	64) a	229) b	230) a	231) a	232) a
65) a	66) a	67) c	68) d	233) c	234) c	235) a	236) a
69) c	70) c	71) a	72) b	237) d	238) c	239) a	240) c
73) a	74) d	75) c	76) c	241) c	242) a	243) d	244) b
77) d	78) d	79) c	80) d	245) b	246) d	247) c	248) c
81) d	82) c	83) c	84) a	249) a	250) c	251) d	252) d
85) a	86) a	87) b	88) c	253) d	254) a	255) c	256) c
89) a	90) c	91) b	92) c	257) d	258) a	259) c	260) c
93) c	94) a	95) c	96) b	261) b	262) d	263) a	264) b
97) a	98) c	99) a	100) b	265) a	266) c	267) b	268) c
101) d	102) c	103) c	104) a	269) c	270) c	271) b	272) b
105) a	106) c	107) d	108) a	273) c	274) d	275) b	276) b
109) c	110) b	111) b	112) d	277) d	278) b	279) d	280) d
113) b	114) d	115) b	116) a	281) a	282) c	283) b	284) c
117) d	118) a	119) d	120) d	285) c	286) b	287) a	288) c
121) a	122) b	123) d	124) d	289) a	290) a	291) d	292) d
125) c	126) a	127) d	128) b	293) b	294) a	295) a	296) c
129) c	130) b	131) d	132) b	297) a	298) b	299) d	300) a
133) a	134) c	135) a	136) b	301) c	302) c	303) b	304) a
137) d	138) b	139) d	140) d	305) a	306) a	307) d	308) b
141) c	142) a	143) a	144) c	309) a	310) b	311) b	312) c
145) d	146) b	147) b	148) d	313) a	314) a	315) d	316) a
149) d	150) c	151) a	152) d	317) b	318) c	319) b	320) a
153) a	154) d	155) b	156) a	321) c	322) d	323) a	324) d
157) c	158) d	159) b	160) b	325) d	326) c	327) d	328) b
161) c	162) d	163) b	164) b	329) a	330) b	331) a	332) b
165) c	166) a	167) d	168) a	333) c	334) d	335) d	336) a

337) a	338) c	339) a	340) d
341) c	342) b	343) a	344) c
345) c	346) b	347) d	348) a
349) a	350) c	351) a	352) d
353) a	354) a	355) b	356) b
357) b	358) c	359) a	360) c
361) b	362) d	363) d	364) c
365) a	366) b	367) c	368) d
369) a	370) d	371) c	372) c
373) c	374) d	375) b	376) d
377) b	378) a	379) b	380) c
381) d	382) d	383) a	384) a
385) b	386) d	387) c	

: HINTS AND SOLUTIONS :

1 (d)

$$v_{rms} = \sqrt{\frac{v_1^2 + v_2^2 + v_3^2 + v_4^2 + v_5^2}{5}} = 4.24$$

2 (a)

Rate of cooling proportional to $(T^4 - T_0^4)$, as per

Stefan's Law.

$$\therefore \frac{R'}{R} = \frac{(900)^4 - (300)^4}{(600)^4 - (300)^4}$$

$$\therefore \frac{9^4 - 3^4}{6^4 - 3^4} = \frac{3^4(3^4 - 1)}{3^4(2^4 - 1)}$$

$$= \frac{80}{15} = \frac{16}{3}$$

$$R' = \frac{16}{3}R$$

3 (c)

The temperature rises by the same amount in the two cases and the internal energy of an ideal gas depends only on its temperature

$$\text{Hence } \frac{U_1}{U_2} = \frac{1}{1}$$

4 (b)

$$\frac{E_2}{E_1} = \left(\frac{T_2}{T_1}\right)^4$$

$$= \left(\frac{273+84}{273+27}\right)^4 = \left(\frac{357}{300}\right)^4 = 2.0$$

5 (a)

$$\text{Kinetic energy for } mg \text{ gas } E = \frac{f}{2} mrT$$

If only translational degree of freedom is considered

$$\text{Then } f=3 \Rightarrow E_{trans} = \frac{3}{2} mrT = \frac{3}{2} m \left(\frac{R}{M}\right) T$$

$$\therefore \frac{3}{2} \times 20 \times \frac{8.3}{32} \times (273+47) = 2490 \text{ J}$$

6 (c)

The number of moles of the system remains same,

$$\frac{P_1 V_1}{RT_1} + \frac{P_2 V_2}{RT_2} = \frac{P(V_1+V_2)}{RT} \Rightarrow T = \frac{P(V_1+V_2)T_1 T_2}{(P_1 V_1 T_2 + P_2 V_2 T_1)}$$

According to Boyle's law,

$$P_1 V_1 + P_2 V_2 = P(V_1 + V_2) \therefore T = \frac{(P_1 V_1 + P_2 V_2) T_1 T_2}{(P_1 V_1 T_2 + P_2 V_2 T_1)}$$

7 (b)

Saturated water vapour do not obey gas laws

8 (c)

$$v_{rms} = \sqrt{\frac{3RT}{M}} = T \propto M [\because v_{rms}, R \rightarrow \text{constant}]$$

$$\Rightarrow \frac{T_{O_2}}{T_{N_2}} = \frac{M_{O_2}}{M_{N_2}} \Rightarrow \frac{T_{O_2}}{(273+0)} = \frac{32}{28} \Rightarrow T_{O_2} = 312 \text{ K} = 39^\circ \text{C}$$

9 (c)

Boyle's and Charle's law follow kinetic theory of gases

10 (b)

$$F = \frac{3}{2} kT \Rightarrow E \propto T$$

12 (a)

In elastic collision kinetic energy is conserved

13 (c)

From the Mayer's formula

$$C_p - C_v = R$$

... (i)

$$\text{and } \gamma = \frac{C_p}{C_v}$$

$$\Rightarrow \gamma C_v = C_p$$

... (ii)

Substituting Eq. (ii) in Eq. (i) we get

$$\Rightarrow \gamma C_v - C_v = R$$

$$C_v(\gamma - 1) = R$$

$$C_v = \frac{R}{\gamma - 1}$$

14 (b)

From Andrews curve

15 (a)

The rms velocity of an ideal gas is

$$v_{rms} = \sqrt{\frac{3RT}{M}}$$

Where T is the absolute temperature and M is the molar mass of an ideal gasSince M remains the same

$$\therefore v_{rms} \propto \sqrt{T}$$

$$\frac{v'_{rms}}{v_{rms}} = \sqrt{\frac{T'}{T}} = \sqrt{\frac{3T}{T}}$$

$$\Rightarrow v'_{rms} = \sqrt{3} v_{rms}$$

16 (c)

At constant temperature; $PV = \dot{i}$ constant

$$\Rightarrow P \times \left(\frac{m}{D}\right) = \text{constant}$$

$$\Rightarrow \frac{P}{D} = \text{constant} = K. [D = \dot{i} \text{ Density}]$$

17 (a)

$$v_{rms} = \sqrt{\frac{3p}{\rho}} \Rightarrow \frac{v_1}{v_2} = \sqrt{\frac{\rho_2}{\rho_1}} = \sqrt{\frac{16}{1}} = \frac{4}{1}$$

18 (a)

The gases carbon monoxide (CO) and nitrogen (N_2) are diatomic, so both have equal kinetic

energy $\frac{5}{2}kT$, i.e. $E_1 = E_2$.

19 (a)

From ideal gas equation, we have

$$pV = nRT$$

$$\therefore n = \frac{pV}{RT}$$

Given, $p = 22.4 \text{ atm pressure}$

$$\dot{i} 22.4 \times 1.01 \times 10^5 \text{ N m}^{-2},$$

$$V = 2L = 2 \times 10^{-3} \text{ m}^3,$$

$$R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1},$$

$$T = 273 \text{ K}$$

$$\therefore n = \frac{22.4 \times 1.01 \times 10^5 \times 2 \times 10^{-3}}{8.31 \times 273}$$

$$n = 1.99 \approx 2$$

Since, $n = \frac{\text{Mass}}{\text{Atomic weight}}$

We have,

$$\text{mass} = n \times \text{atomic weight} = 2 \times 14 = 28 \text{ g}$$

20 (d)


Average kinetic energy $E = \frac{3}{2}kT$

$$\Rightarrow E \propto T$$

Thus, average kinetic energy of a gas molecule is directly proportional to the absolute temperature of gas.

21 (c)

For a given pressure, volume will be more if temperature is more [Charle's law]



From the graph it is clear that $V_2 > V_1 \Rightarrow T_2 > T_1$

22 (d)

$$C_{rms} = \sqrt{\frac{3RT}{M}}$$

$$\text{Or } M = \frac{3RT}{C_{rms}^2} = \frac{3 \times 8.31 \times 300}{(1920)^2}$$

$$= 2 \times 10^{-3} \text{ kg} = 2 \text{ g}$$

Since, $M = 2$ for the hydrogen molecule. Hence, the gas is hydrogen.

23 (d)

$$v_{rms} = \sqrt{\frac{3P}{\rho}} = P \propto \rho [v_{rms} \text{ is constant for fixed temperature}]$$

24 (c)

According to Boyle's law

$$p_1 V_1 = p_2 V_2$$

As the pressure is decreased by 20%, so

$$p_2 = \frac{80}{100} p_1$$

$$p_1 V_1 = \frac{80}{100} p_1 V_2$$

$$V_1 = \frac{80}{100} V_2$$

Percentage increase in volume

$$\dot{i} \frac{V_2 - V_1}{V_1} \times 100$$

$$\dot{i} \frac{100 - 80}{80} \times 100 = 25\%$$

25 (d)

Root mean square velocity,

$$c = \sqrt{\frac{3pV}{M}} = \sqrt{\frac{3RT}{M}}$$

$$c_1 = \sqrt{\frac{3R(T/2)}{2M}} = \frac{1}{2} \sqrt{\frac{3RT}{M}}$$

$$\dot{i} \frac{c}{2} = \frac{300}{2} = 150 \text{ m s}^{-1}$$

26 (c)

At TK, pressure of gas (P) in the jar

= Total pressure – saturated vapour pressure

$$\Rightarrow P = (830 - 30) = 800 \text{ mm of Hg}$$

$$\text{New temperature } T' = \left(T - \frac{T}{100}\right) = \frac{99T}{100}$$

$$\text{Using Charles's law } \frac{P}{T} = \frac{P'}{T'} \Rightarrow P' = \frac{PT'}{T}$$

$$\therefore \frac{800 \times 99T}{100T} = 792 \text{ mm of Hg}$$

Saturated vapour pressure at $T' = 25 \text{ mm of Hg}$

\therefore Total pressure in the jar

\therefore Actual pressure of gas + \therefore Saturated vapour pressure

$$\therefore 792 + 25 = 817 \text{ mm of Hg}$$

28 (c)

$$\mu_1 = \frac{PV}{RT}, \mu_2 = \frac{PV}{RT}$$

$$P' = \frac{(\mu_1 + \mu_2)RT}{V} = \frac{2PV}{RT} \times \frac{RT}{V} = 2P$$

29 (d)

$$\text{Average kinetic energy } E = \frac{f}{2} kT$$

Since f and T are same for both the gases so they will have equal energies also

30 (b)

$$V_{rms} = \sqrt{\frac{3RT}{M}} \Rightarrow \frac{(V_{rms})_{O_2}}{(V_{rms})_{H_2}} = \sqrt{\frac{T_{O_2}}{T_{H_2}} \times \frac{M_{H_2}}{M_{O_2}}}$$

$$\Rightarrow \frac{(V_{rms})_{O_2}}{(V_{rms})_{H_2}} = \sqrt{\frac{900}{300} \times \frac{2}{32}} = \frac{\sqrt{3}}{4}$$

$$\Rightarrow (v_{rms})_{O_2} = 836 \text{ m/s}$$

31 (a)

As degree of freedom is defined as the number of independent variables required to define body's motion completely. Here $f = 2$ (1 Translational + 1 Rotational)

32 (b)

$$\frac{E_1}{E_2} = \frac{A_1}{A_2} \cdot \left(\frac{T_1}{T_2}\right)^4 = \frac{4\pi r_1^2}{4\pi r_2^2} \times 1 = \left(\frac{1}{2}\right)^2 = \frac{1}{4}$$

33 (c)

$$V_{rms} = \sqrt{\frac{3P}{\rho}} \Rightarrow P = \frac{\rho V_{rms}^2}{3}$$

$$\therefore \frac{8.99 \times 10^{-2} \times 1840 \times 1840}{3} = 1.01 \times 10^5 \text{ N/m}^2$$

34 (b)

$$v_{rms} = \sqrt{\frac{3RT}{M}} \Rightarrow v_{rms} \propto \sqrt{T}$$

v_{rms} is to reduce two times, i.e., the temperature of the gas will have to reduce four times or

$$\frac{T'}{T} = \frac{1}{4}$$

During adiabatic process,

$$TV^{\gamma-1} = T'V'^{\gamma-1}$$

$$\text{or } \frac{V'}{V} = \left(\frac{T}{T'}\right)^{\frac{1}{\gamma-1}}$$

$$\therefore (4)^{\frac{1}{1.5-1}} = 4^2 = 16$$

$$\therefore V' = 16V$$

35 (a)

$$(\Delta Q)_V = \mu C_V \Delta T \Rightarrow (\Delta Q)_V = 1 \times C_V \times 1 = C_V$$

$$\text{For monoatomic gas } C_V = \frac{3}{2}R$$

$$\therefore (\Delta Q)_V = \frac{3}{2}R$$

36 (a)

Root mean square velocity

$$v_{rms} \propto \frac{1}{\sqrt{M}}$$

$$\text{So } \frac{(v_{rms})_{O_2}}{(v_{rms})_{H_2}} = \sqrt{\frac{M_{H_2}}{M_{O_2}}}$$

$$\therefore \sqrt{\frac{2}{32}} = \frac{1}{4}$$

37 (c)

$$\text{At constant pressure } V \propto T \Rightarrow \frac{\Delta V}{V} = \frac{\Delta T}{T}$$

Hence ratio of increase in volume per degree rise in kelvin temperature to its original volume

$$\therefore \frac{(\Delta V / \Delta T)}{V} = \frac{1}{T}$$

38 (c)

$$\rho = \frac{PM}{RT}$$

Density ρ remains constant when P/T or volume remains constant.

In graph (i) Pressure is increasing at constant temperature hence volume is decreasing so density is increasing. Graphs (ii) and (iii) volume is increasing hence, density is decreasing. Note that volume would have been constant in case the straight line the graph (iii) had passed through origin

39 (d)

According to Newton's law

$$\frac{\theta_1 - \theta_2}{t} = K \left[\frac{\theta_1 + \theta_2}{2} - \theta_0 \right]$$

$$\therefore \frac{60-50}{10} = K \left[\frac{60+50}{2} - 25 \right] \dots(i)$$

Let θ be the temperature after another 10 min

$$\therefore \frac{50-\theta}{10} = K \left[\frac{\theta+50}{2} - 25 \right] \dots(ii)$$

Dividing Eq.(i) by Eq. (ii), we get

$$\frac{10}{50-\theta} = \frac{30 \times 2}{\theta} \therefore \theta = 42.85^\circ\text{C}$$

40 (c)

$$\left(\frac{\Delta Q}{\Delta t} \right)_{\text{inner}} + \left(\frac{\Delta Q}{\Delta t} \right)_{\text{outer}} = \left(\frac{\Delta Q}{\Delta t} \right)_{\text{total}}$$

$$\frac{K_1 \pi r^2 (T_2 - T_1)}{l} + \frac{K_2 \pi [(2r)^2 - r^2] (T_2 - T_1)}{l} = \frac{K \pi (2r)^2 (T_2 - T_1)}{l}$$

$$\text{or } (K_1 + 3K_2) \frac{\pi r^2 (T_2 - T_1)}{l} = \frac{K \pi 4r^2 (T_2 - T_1)}{l}$$

$$\therefore K = \frac{K_1 + 3K_2}{4}$$

41 (d)

$$Y_{\text{mixture}} = \frac{\frac{\mu_1 Y_1}{Y_1 - 1} + \frac{\mu_2 Y_2}{Y_2 - 1}}{\frac{\mu_1}{Y_1 - 1} + \frac{\mu_2}{Y_2 - 1}}$$

$$\mu_1 = 4 \text{ moles of helium } \therefore \frac{16}{4} = 4$$

$$\mu_2 = 1 \text{ moles of oxygen } \therefore \frac{16}{32} = \frac{1}{2}$$

$$\Rightarrow Y_{\text{mix}} = \frac{\frac{4 \times 5/3}{5/3 - 1} + \frac{1/2 \times 7/5}{7/5 - 1}}{\frac{4}{5/3 - 1} + \frac{1/2}{7/5 - 1}} = 1.62$$

42 (a)

$$\text{Mean free path, } \lambda = \frac{1}{\sqrt{2} \pi d^2 n}$$

Where, $n = \dot{i}$ Number of molecules per unit volume

$d = \dot{i}$ Diameter of the molecules

43 (b)

Speed of sound in gases is given by

$$v_{\text{sound}} = \sqrt{\frac{\gamma P}{\rho}} = \frac{v_1}{v_2} = \sqrt{\frac{\rho_2}{\rho_1}} = \sqrt{\frac{d_2}{d_1}}$$

44 (c)

$$n_1 C_{v1} \Delta T_1 = n_2 C_{v2} \Delta T_2$$

$$\Rightarrow n_1 \times \frac{3}{2} R \times 10 = n_2 \times \frac{5}{2} R \times 6 \Rightarrow \frac{n_1}{n_2} = 1$$

45 (a)

We treat water like a solid. For each atom average

energy is $3k_B T$. Water molecule has three atoms, two hydrogen and one oxygen. The total energy of one mole of water is

$$U = 3 \times 3k_B T \times N_A = 9RT \left[\because k_B = \frac{R}{N_A} \right]$$

\therefore Heat capacity per mole of water is

$$C = \frac{\Delta Q}{\Delta T} = \frac{\Delta U}{\Delta T} = 9R$$

46 (a)

K.E. is function of temperature. So $\frac{E_{H_2}}{E_{O_2}} = \frac{1}{1}$

47 (c)

According to kinetic theory of gases the temperature of a gas is a measure of the kinetic energies of the molecules of the gas.

48 (c)

At constant volume

$$\frac{P_1}{T_1} = \frac{P_2}{T_2} \Rightarrow T_2 = \left(\frac{P_2}{P_1} \right) T_1$$

$$\Rightarrow T_2 = \left(\frac{3P}{P} \right) \times (273 + 35) = 3 \times 308 = 924 \text{ K} = 651^\circ\text{C}$$

49 (d)

$$\frac{3}{2} kT = 1 \text{ eV} \Rightarrow T = \frac{2 \text{ eV}}{3 k} = \frac{\frac{2}{3} \times 1.6 \times 10^{-19}}{1.38 \times 10^{-23}} = 7.7 \times 10^3 \text{ K}$$

51 (b)

Vander Waal's gas equation for μ mole of real gas

$$\left(P + \frac{\mu^2 a}{V^2} \right) (V - \mu b) = \mu RT$$

$$P = \left(\frac{\mu RT}{V - \mu b} - \frac{\mu^2 a}{V^2} \right)$$

Given equation,

$$P = \left(\frac{RT}{2V - b} - \frac{a}{4b^2} \right)$$

On comparing the given equation with this standard equation, we get

$$\mu = \frac{1}{2}$$

$$\text{Hence, } \mu = \frac{m}{M}$$

$$\Rightarrow \text{mass of gas, } m = \mu M = \frac{1}{2} \times 44 = 22 \text{ g}$$

52 (d)

$$C_P = \left(\frac{f}{2} + 1 \right) R = \left(\frac{5}{2} + 1 \right) R = \frac{7}{2} R$$

53 (c)

$$\frac{R}{C_p} = \frac{R}{7/2R} = \frac{2}{7} \left[\because C_p = \frac{7}{2} R \right]$$

54 (c)

As temperature decreases to half and volume made twice, hence pressure becomes $\frac{1}{4}$ times

55 (d)

$$\begin{aligned} p &= p_1 + p_2 + p_3 \\ &= \left(\frac{nRT}{V} \right)_{O_2} + \left(\frac{nRT}{V} \right)_{N_2} + \left(\frac{nRT}{V} \right)_{CO_2} \\ &= (n_{O_2} + n_{N_2} + n_{CO_2}) \frac{RT}{V} \\ &= \frac{(0.25 + 0.5 + 0.5)(8.31) \times 300}{4 \times 10^{-3}} \\ &= 7.79 \times 10^5 \text{ N m}^{-2} \end{aligned}$$

56 (a)

$$\begin{aligned} PV &= \mu RT = \frac{m}{M} RT \Rightarrow V = \frac{mRT}{MP} \\ &= \frac{2 \times 10^{-3} \times 8.3 \times 300}{32 \times 10^{-3} \times 10^5} = 1.53 \times 10^{-3} \text{ m}^3 = 1.53 \text{ litre} \end{aligned}$$

57 (c)

According to Boyle's law

$$\begin{aligned} (P_1 V_1)_{\text{At top of the lake}} &= (P_2 V_2)_{\text{At the bottom of the lake}} \\ \Rightarrow P_1 V_1 &= (P_1 + h) V_2 \Rightarrow 10 \times \frac{4}{3} \pi \left(\frac{5r}{4} \right)^3 \\ \Rightarrow (10 + h) \times \frac{4}{3} \pi r^3 &\Rightarrow h = \frac{610}{64} = 9.53 \text{ m} \end{aligned}$$

58 (d)

We have $v_{rms} = \sqrt{\frac{3RT}{M}}$; at $T = T_0$ (NTP)

$$v_{rms} = \sqrt{\frac{3RT_0}{M}}$$

But at temperature T ,

$$v_{rms} = 2 \times \sqrt{\frac{3RT_0}{M}}$$

$$\Rightarrow \sqrt{\frac{3RT}{M}} = 2 \sqrt{\frac{3RT_0}{M}}$$

$$\Rightarrow \sqrt{T} = \sqrt{4T_0}$$

or $T = 4T_0$

$$T = 4 \times 273 \text{ K} = 1092 \text{ K}$$

$$\therefore T = 819^\circ \text{C}$$

59 (b)

$$E = \frac{f}{2} RT; f = 5 \text{ for diatomic gas} \Rightarrow E = \frac{5}{2} RT$$

60 (d)

Average kinetic energy

$$E = \frac{3}{2} kT \Rightarrow \frac{E_1}{E_2} = \frac{T_1}{T_2} = \frac{(273-23)}{(273+227)} = \frac{250}{500} = \frac{1}{2}$$

$$\Rightarrow E_2 = 2E_1 = 2 \times 5 \times 10^{-14} = 10 \times 10^{-14} \text{ erg}$$

61 (a)

A monoatomic gas molecule has only three translational degrees of freedom

62 (b)

$$\begin{aligned} Y_{mix} &= \frac{\frac{\mu_1 Y_1}{Y_1 - 1} + \frac{\mu_2 Y_2}{Y_2 - 1}}{\frac{\mu_1}{Y_1 - 1} + \frac{\mu_2}{Y_2 - 1}} = \frac{\frac{3 \times 1.3}{(1.3 - 1)} + \frac{2 \times 1.4}{(1.4 - 1)}}{\frac{3}{(1.3 - 1)} + \frac{2}{(1.4 - 1)}} = 1.33 \end{aligned}$$

63 (b)

At critical temperature the horizontal portion in $P-V$ curve almost vanishes as at temperature T_2 .

Hence the correct answer will be (b)

64 (a)

$$v_{rms} \propto \frac{1}{\sqrt{M}} \Rightarrow \frac{(v_{rms})_{H_2}}{(v_{rms})_{He}} = \sqrt{\frac{M_{He}}{M_{H_2}}} = \sqrt{\frac{4}{2}} = \frac{\sqrt{2}}{1}$$

65 (a)

When electric spark is passed, hydrogen reacts with oxygen to form water (H_2O). Each gram of hydrogen reacts with eight grams of oxygen. Thus 96 gm of oxygen will be totally consumed together with 12 gm of hydrogen. The gas left in the vessel will be 2 gm of hydrogen *i.e.*

$$\text{Number of moles } \mu = \frac{2}{2} = 1$$

$$\text{Using } PV = \mu RT \Rightarrow P \propto \mu \Rightarrow \frac{P_2}{P_1} = \frac{\mu_2}{\mu_1}$$

($\mu_1 = 7$ Initial number of moles $7 + 3 = 10$ and $\mu_2 = 1$ Final number of moles)

$$\Rightarrow \frac{P_2}{1} = \frac{1}{10} \Rightarrow P_2 = 0.1 \text{ atm}$$

66 (a)

$$v_{rms} = \sqrt{\frac{3RT}{M}} = \frac{v_2}{v_1} = \sqrt{\frac{T_2}{T_1}} = \sqrt{\frac{(273+90)}{(273+27)}} = 1.1$$

$$\% \text{ increase } = \left(\frac{v_2}{v_1} - 1 \right) \times 100 = 0.1 \times 100 = 10\%$$

67 (c)

Ideal gas equation is given by

$$pV = nRT$$

...(i)

For oxygen, $p = 1 \text{ atm}$, $V = 1 \text{ L}$, $n = n_{O_2}$

Therefore, Eq. (i) becomes

$$\therefore 1 \times 1 = n_{O_2} RT$$

$$\Rightarrow n_{O_2} = \frac{1}{RT}$$

For nitrogen $p = 0.5 \text{ atm}$, $V = 2 \text{ L}$, $n = n_N$
 $\therefore 0.5 \times 2 = n_{N_2} RT$

$$\Rightarrow n_{N_2} = \frac{1}{RT}$$

For mixture of gas

$$p_{\text{mix}} V_{\text{mix}} = n_{\text{mix}} RT$$

Here, $n_{\text{mix}} = n_{O_2} + n_{N_2}$

$$\therefore \frac{p_{\text{mix}} V_{\text{mix}}}{RT} = \frac{1}{RT} + \frac{1}{RT}$$

$$\Rightarrow p_{\text{mix}} V_{\text{mix}} = 2 \quad (V_{\text{mix}} = 1)$$

68 (d)

Let T_0 be the initial temperature of the black body

$\therefore \lambda_0 T_0 = b$ (Wien's law)

Power radiated, $P_0 = C T_0^4$, where, C is constant.

If T is new temperature of black body, then

$$\frac{3\lambda_0}{4} T = b = \lambda_0 T_0 \Rightarrow T = \frac{4}{3} T_0$$

$$\text{Power radiated, } P = C T^4 = C T_0^4 \left(\frac{4}{3}\right)^4$$

$$P = P_0 \times \frac{256}{81} \Rightarrow \frac{P}{P_0} = \frac{256}{81}$$

69 (c)

$$PV = \frac{m}{M} RT \Rightarrow V \propto mT \Rightarrow \frac{V_1}{V_2} = \frac{m_1}{m_2} \cdot \frac{T_1}{T_2}$$

$$\therefore \frac{2V}{V} = \frac{m}{m_2} \times \frac{100}{200} \Rightarrow m_2 = \frac{m}{4}$$

70 (c)

At constant temperature $PV = \text{constant} \Rightarrow P \propto \frac{1}{V}$

71 (a)

$$v_{\text{rms}} \propto \frac{1}{\sqrt{M}} \Rightarrow (v_{\text{rms}})_1 < (v_{\text{rms}})_2 < (v_{\text{rms}})_3 \text{ also in mixture}$$

temperature of each gas will be same, hence kinetic energy also remains same

72 (b)

$$\frac{E_1}{E_2} = \frac{T_1}{T_2} = \frac{300}{450} = \frac{2}{3}$$

73 (a)

$$PV = \mu RT = \frac{m}{M} RT \Rightarrow P = \frac{d}{M} RT \left[\text{Density } d = \frac{m}{V} \right]$$

$$\Rightarrow \frac{P}{dT} = \text{constant} \Rightarrow \frac{P_1}{d_1 T_1} = \frac{P_2}{d_2 T_2}$$

74 (d)

$$P \propto T \Rightarrow \frac{P_2}{P_1} = \frac{T_2}{T_1} = \frac{(273+100)}{(273+0)} = \frac{373}{273}$$


$$\Rightarrow P_2 = \frac{760 \times 373}{273} = 1038 \text{ mm}$$

75 (c)

Since temperature is constant, so v_{rms} remains same

76 (c)

At constant pressure, the volume of a given mass of a gas is directly proportional to its absolute temperature (T).



$$\text{ie., } \frac{V}{T} = \text{constant}$$

This is another form of Charles' law. Hence, variation of volume with temperature is as shown.

Hence, correct graph will be (C).

77 (d)

Argon is a monoatomic gas so it has only translational energy

79 (c)

According to the Dalton's law of partial pressure, the total pressure will be $P_1 + P_2 + P_3$

80 (d)

Kinetic energy \propto Temperature

$$\Rightarrow \frac{E_1}{E_2} = \frac{T_1}{T_2} \Rightarrow \frac{E_1}{E_2} = \frac{(273+27)}{(273+927)} = \frac{300}{1200} = \frac{1}{4}$$

$$\Rightarrow E_2 = 4 E_1$$

81 (d)

$$\frac{V_{\text{rmsHe}}}{V_{\text{rmsAr}}} = \frac{\sqrt{\frac{3RT}{m_{\text{He}}}}}{\sqrt{\frac{3RT}{m_{\text{Ar}}}}} = \sqrt{\frac{m_{\text{Ar}}}{m_{\text{He}}}} = \sqrt{\frac{40}{4}} = \sqrt{10} \approx 3.16$$

82 (c)

We know that $C_p - C_v = \frac{R}{J}$

$$\Rightarrow J = \frac{R}{C_p - C_v}$$

$$C_p - C_v = 1.98 \frac{\text{cal}}{\text{g-mol-K}}$$

$$R = 8.32 \frac{J}{g \cdot mol \cdot K}$$

$$\therefore J = \frac{8.32}{1.98} = 4.20 J/cal$$

83 (c)

S.I. unit of R is $J/mol \cdot K$

84 (a)

According to Boyle's law $PV = \text{constant}$

85 (a)

$$v_{rms} \propto \sqrt{\frac{3RT}{M}}$$

$$\Rightarrow T \propto v_{rms}^2$$

$$\Rightarrow \frac{T_2}{T_1} = \left[\frac{v_2}{v_1} \right]^2 = \frac{1}{4} \Rightarrow T_2 = \frac{T_1}{4} = \frac{273+327}{4}$$

$$\therefore 150 K = -123^\circ C$$

86 (a)

The total pressure exerted by a mixture of non-reacting gases occupying a vessel is equal to the sum of the individual pressure which each gas exert if it alone occupied the same volume at a given temperature.

For two gases,

$$p = p_1 + p_2 = p + p = 2p$$

87 (b)

According to ideal gas equation $PV = nRT$

$$PV = \frac{m}{M} RT, P = \frac{\rho}{M} RT \text{ or } \frac{\rho}{P} = \frac{M}{RT} \text{ or } \frac{\rho}{P} \propto \frac{1}{T}$$

Here, $\frac{\rho}{P}$ represent the slope of graph

Hence $T_2 > T_1$

88 (c)

$$PV = \mu RT = \frac{m}{M} RT \Rightarrow P \propto mT$$

$$\Rightarrow \frac{P_2}{P_1} = \frac{m_2 T_2}{m_1 T_1} = \frac{1}{2} \times \frac{(273+27+50)}{(273+27)} = \frac{7}{12}$$

$$\Rightarrow P_2 = \frac{7}{12} P_1 = \frac{7}{12} \times 20 = 11.67 \text{ atm} \approx 11.7 \text{ atm}$$

89 (a)

Since $c_{rms} \ll V_e$, hence molecules do not escape out

91 (b)

In case of given graph, V and T are related as $V = aT - b$, where a and b are constants.

From ideal gas equation, $PV = \mu RT$

$$\text{We find } P = \frac{\mu RT}{aT - b} = \frac{\mu R}{a - b/T}$$

Since $T_2 > T_1$, therefore $P_2 < P_1$

92 (c)

Gas equation for N molecules $PV = NkT$

$$\Rightarrow N = \frac{PV}{kT} = \frac{1.2 \times 10^{-10} \times 13.6 \times 10^3 \times 10 \times 10^{-4}}{1.38 \times 10^{-23} \times 300}$$

$$\therefore 3.86 \times 10^{11}$$

93 (c)

$$E \propto T$$

94 (a)

$$v_{rms} \propto \sqrt{T}, \frac{v_2}{v_1} = \sqrt{\frac{T_2}{T_1}} \Rightarrow v_2 = \sqrt{\frac{273+927}{273+27}} v_1 \Rightarrow v_2 = 2$$

95 (c)

For ideal gas, on free expansion there is no change in temperature. Hence $C_a = C_b$

96 (b)

$$v_{rms} > v_{av} > v_{mp}$$

97 (a)

According to Boyle's law, $pV = k$ (a constant)

$$\text{Or } p \frac{m}{p} = k \vee p = \frac{pm}{k}$$

$$\text{Or } p = \frac{p}{k} \text{ (a constant)}$$

$$\text{So, } \rho_1 = \frac{p_1}{k} \wedge V_1 \frac{p_1}{k} = \frac{m_1}{p_1} = \frac{m_1}{p_1/k} = \frac{k m_1}{p_1}$$

$$\text{Similarly, } V_2 = \frac{k m_2}{p_2}$$

$$\text{Total volume} = V_1 + V_2 = k \left(\frac{m_1}{p_1} + \frac{m_2}{p_2} \right)$$

Let p be the common pressure and ρ be the common density of mixture. Then

$$\rho = \frac{m_1 + m_2}{V_1 + V_2} = \frac{m_1 + m_2}{k \left(\frac{m_1}{p_1} + \frac{m_2}{p_2} \right)}$$

$$\therefore p = k\rho = \frac{m_1 + m_2}{\frac{m_1}{p_1} + \frac{m_2}{p_2}} = \frac{p_1 p_2 (m_1 + m_2)}{(m_1 p_2 + m_2 p_1)}$$

98 (c)

$$v_{rms} = \sqrt{\frac{3RT}{M}}. \text{ According to problem } T \text{ will become}$$

$2T$ and M will become $M/2$ so the value of v_{rms} will increase by $\sqrt{4} = 2 \times$, i.e., new root mean square velocity will be $2v$

99 (a)

$$\text{Here, } \frac{K_1}{K_2} = \frac{1}{2}, \frac{r_1}{r_2} = \frac{1}{2}$$

$$\therefore \frac{A_1}{A_2} = \frac{1}{4}$$

$$\frac{dx_1}{dx_2} = \frac{1}{2}, \frac{dQ_2}{dt} = 4 \text{ cal s}^{-1}, \frac{dQ_1}{dt} = ?$$

$$\frac{dQ_2/dt}{dQ_1/dt} = \frac{K_2 A_2 dT/dx_2}{K_1 A_1 dT/dx_1} = \frac{K_2 A_2 dx_1}{K_1 A_1 dx_2}$$

$$= 2 \times 4 \times \frac{1}{2} = 4$$

$$\frac{dQ_1}{dt} = \frac{dQ_2/dt}{4} = \frac{4}{4} = 1 \text{ cal s}^{-1}$$

100 (b)

At lower pressure we can assume that given gas behaves as ideal gas so $\frac{PV}{RT} = \dot{i}$ constant but when pressure increases, the decrease in volume will not take place in same proportion so $\frac{PV}{RT}$ will increase

101 (d)

Let initial conditions $\dot{i} V, T$
 And final conditions $\dot{i} V', T'$
 By Charle's law $V \propto T$ [P remains constant]

$$\frac{V}{T} = \frac{V'}{T'} \Rightarrow \frac{V}{T} = \frac{V'}{1.2T'} \Rightarrow V' = 1.2V$$


But as per question, volume is reduced by 10% means $V' = 0.9V$

So percentage of volume leaked out

$$\dot{i} \frac{(1.2 - 0.9)V}{1.2V} \times 100 = 25\%$$

102 (c)

As temperature requirement is not given so, the molecule of a triatomic gas has a tendency of rotating about any of three coordinate axes. So, it has 6 degrees of freedom; 3 translational and 3 rotational.



Thus,

(3 translational + 3 rotational) at room temperature.

103 (c)

$$\text{We have } v_{rms} = \sqrt{\frac{v_1^2 + v_2^2 + \dots + v_n^2}{n}}$$

$$= \sqrt{\frac{4 + 25 + 9 + 36 + 9 + 25}{6}}$$

$$=$$

$$\sqrt{\frac{108}{6}} = \sqrt{18} = 3\sqrt{2} = 3 \times 1.414 = 4.242 \text{ unit.}$$

104 (a)

According to ideal gas equation

$$PV = nRT \text{ or } \frac{V}{T} = \frac{nR}{P}$$

At constant pressure

$$\frac{V}{T} = \dot{i} \text{ constant}$$

Hence graph (a) is correct

105 (a)

Temperatures $T_1 = 15^\circ\text{C} = 15 + 273 = 288 \text{ K}$

$T_2 = 35^\circ\text{C} = 35 + 273 = 308 \text{ K}$

Volume remains constant.

$$\text{So, } \frac{p_1}{T_1} = \frac{p_2}{T_2}$$

$$\frac{p_1}{p_2} = \frac{T_1}{T_2} \Rightarrow \frac{p_1}{p_2} = \frac{288}{308}$$

$$\frac{p_2}{p_1} = \frac{308}{288}$$

$$\% \text{ increases in pressure} = \frac{p_2 - p_1}{p_1} \times 100$$

$$\dot{i} \frac{308 - 288}{288} \times 100 \approx 7\%$$

106 (c)

$$v_{av} = \sqrt{\frac{8RT}{\pi M}} \Rightarrow T \propto M [\because v_{av}, R \rightarrow \text{constant}]$$

$$\Rightarrow \frac{T_{H_2}}{T_{O_2}} = \frac{M_{H_2}}{M_{O_2}} \Rightarrow \frac{T_{H_2}}{(273 + 31)} = \frac{2}{32}$$

$$\Rightarrow T_{H_2} = 19 \text{ K} = -254^\circ\text{C}$$

107 (d)

$$\text{Kinetic energy per } g \text{ mole } E = \frac{f}{2} RT$$

If nothing is said about gas then we should calculate the translational kinetic energy

$$i.e., E_{trans} = \frac{3}{2} RT = \frac{3}{2} \times 8.31 \times (273+0) = 3.4 \times 10^3$$

108 (a)

According to Gay Lussac's law $p \propto T$

$$\therefore \frac{dp}{p} \times 100 = \frac{dT}{T} \times 100$$

$$1 = \frac{1}{T} \times 100$$

$$\Rightarrow T = 100 \text{ K}$$

109 (c)

Specific heat at constant pressure (C_p) is the amount of heat (Q) required to raise n moles of substance by $\Delta\theta$ when pressure is kept constant.

Then

$$C_p = \frac{Q}{n \Delta\theta}$$

$$\text{Given, } Q = 70 \text{ cal, } n = 2,$$

$$\Delta\theta = (35 - 35)^\circ\text{C} = 5^\circ\text{C}$$

$$\therefore C_p = \frac{70}{2 \times 5} = 7 \text{ cal mol}^{-1} - \text{K}^{-1}$$

From Mayer's formula $C_p - C_v = R$

where R is gas constant ($\approx 2 \text{ cal mol}^{-1}$)

$$\therefore 7 - C_v = 2$$

$$\Rightarrow C_v = 5 \text{ cal mol}^{-1} - \text{K}^{-1}$$

Hence, amount of heat required at constant volume (C_v) is

$$Q' = n C_v \Delta\theta$$

$$Q' = 2 \times 5 \times 5 = 50 \text{ cal}$$

110 (b)

$v_{rms} \propto \sqrt{T}$; To double the rms velocity temperature should be made four times, i.e.,

$$T_2 = 4 T_1 = 4(273+0) = 1092 \text{ K} = 819^\circ\text{C}$$

111 (b)

In a given mass of the gas

$$n = \frac{pV}{kT}$$

k being Boltzmann's constant.

112 (d)

$$PV = NkT \Rightarrow \frac{N_A}{N_B} = \frac{P_A V_A}{P_B V_B} \times \frac{T_B}{T_A}$$

$$\Rightarrow \frac{N_A}{N_B} = \frac{P \times V \times (2T)}{2P \times \frac{V}{4} \times T} = \frac{4}{1}$$

113 (b)

$$VP^3 = \text{constant } k \Rightarrow P = \frac{k}{V^{1/3}}$$

$$\text{Also } PV = \mu RT \Rightarrow \frac{k}{V^{1/3}} \cdot V = \mu RT \Rightarrow V^{2/3} = \frac{\mu RT}{k}$$

$$\text{Hence } \left(\frac{V_1}{V_2}\right)^{2/3} = \frac{T_1}{T_2} \Rightarrow \left(\frac{V}{27V}\right)^{2/3} = \frac{T}{T_2} \Rightarrow T_2 = 9T$$

114 (d)

Vander waal's equation is followed by real gases

115 (b)

Molecular mass of He; $M = 4 \text{ g}$

\Rightarrow Molar value of

$$C_v = M c_v = 4 \times 3 = 12 \frac{\text{J}}{\text{mole} - \text{kelvin}}$$

At constant volume $P \propto T$, therefore on doubling the pressure temperature also doubles

$$i.e., T_2 = 2 T_1 \Rightarrow \Delta T = T_2 - T_1 = 273 \text{ K}$$

$$\text{Also } (\Delta Q)_v = \mu C_v \Delta T = \frac{1}{2} \times 12 \times 273 = 1638 \text{ J}$$

116 (a)

$$\text{Here, } h_1 = 50 \text{ cm, } t_1 = 50^\circ\text{C}$$

$$h_2 = 60 \text{ cm, } t_2 = 100^\circ\text{C}$$

$$\text{Now, } \frac{h_1}{h_2} = \frac{d_2}{d_1} = \frac{d_0}{1 + \gamma t_2} \times \frac{1 + \gamma t_1}{d_0}$$

$$\frac{50}{60} = \frac{1 + \gamma \times 50}{1 + \gamma \times 100}$$

$$\therefore \gamma = \frac{1}{200} = 0.005^\circ\text{C}^{-1}$$

117 (d)

Vander Waal's gas constant $b = 4 \times$ total volume of all the molecules of the gas in the enclosure

$$\text{Or } b = 4 \times N \times \frac{4}{3} \pi \left(\frac{d}{2}\right)^3 = \frac{2}{3} \pi N d^3$$

$$\therefore \frac{2}{3} \times 3.14 \times 6.02 \times 10^{23} \times (2.94 \times 10^{-10})^3 = 32 \times 10^{-1}$$

118 (a)

From ideal gas equation

$$pV = nkT$$

$$p = \frac{n}{V} kT$$

$$\text{Here, } \frac{n}{V} = 5/c \text{ m}^3 = 5 \times 10^6 / \text{m}^3$$

$$\therefore p = k$$

$$p = 20.7 \times 10^{-17} \text{ N m}^{-2}$$

119 (d)

Escape velocity from the earth's surface is

$$11.2 \text{ km/sec}$$

$$\text{So, } v_{rms} = v_{escape} = \sqrt{\frac{3RT}{M}} \Rightarrow T = \frac{(v_{escape})^2 \times M}{3R}$$
$$\therefore \frac{(11.2 \times 10^3)^2 \times (2 \times 10^{-3})}{3 \times 8.31} = 10063 \text{ K}$$

120 (d)

$$v = \sqrt{\frac{\gamma P}{\rho}} = \sqrt{\frac{\frac{5}{3} \times 10^3}{2.6}} = 25 \text{ m/s}$$

121 (a)

The temperature at which protons in a proton gas would have enough energy to overcome Coulomb barrier between them is given by

$$\frac{3}{2}k_B T = K_{av} \dots (i)$$

Where k_{av} is the average kinetic energy of the proton, T is the temperature of the proton gas and k_B is the Boltzmann constant

$$\text{From (i), we get } T = \frac{2K_{av}}{3K_B}$$

Substituting the values, we get

$$T = \frac{2 \times 4.14 \times 10^{-14} \text{ J}}{3 \times 1.38 \times 10^{-23} \text{ J K}^{-1}} = 2 \times 10^9 \text{ K}$$

122 (b)

The pressure exerted by the gas,

$$p = \frac{1}{3} \rho c^2$$

$$\therefore \frac{1}{3} \frac{m}{V} c^2$$

$$\therefore \frac{2}{3} \left(\frac{1}{2} m c^2 \right)$$

$$\left(\because \frac{1}{2} m c^2 = \frac{E}{V} = \text{energy per unit volume, } V = 1 \right)$$

$$p = \frac{2}{3} E$$

123 (d)

$$\text{Here, } \frac{D_1}{D_2} = \frac{1}{2}$$

$$\frac{A_1}{A_2} = \frac{D_1^2}{D_2^2} = \frac{1}{4}$$

$$\frac{dx_1}{dx_2} = \frac{2}{1}$$

$$\frac{dQ_1}{dt} = K A_1 \frac{dT}{dx_1} ; \frac{dQ_2}{dt} = K A_2 \frac{dT}{dx_2}$$

$$\frac{dQ_1/dt}{dQ_2/dt} = \frac{A_1}{A_2} \cdot \frac{dx_2}{dx_1} = \frac{1}{4} \times \frac{1}{2} = \frac{1}{8}$$

124 (d)

Total pressure (P) of gas = Actual pressure of gas

$P_a + i$ aqueous vapour pressure (P_v)

$$\Rightarrow P_a = P - P_v = 735 - 23.8 = 711.2 \text{ mm}$$

125 (c)

Let for mixture of gases, specific heat at constant volume be C_v

$$C_v = \frac{n_1(C_v)_1 + n_2(C_v)_2}{n_1 + n_2}$$

where for oxygen; $C_{v1} = \frac{5R}{2}, n_1 = 2 \text{ mol}$

For helium; $C_{v2} = \frac{3R}{2}, n_2 = 8 \text{ mol}$

$$\text{Therefore, } C_v = \frac{\frac{2 \times 5R}{2} + 8 \times \frac{3R}{2}}{2 + 8} = \frac{17R}{10} = 1.7R$$

126 (a)

For one $g \text{ mole}$; average kinetic energy $\therefore \frac{3}{2} RT$

127 (d)

As we know 1 mol of any ideal gas at STP occupies a volume of 22.4 litres .

$$\text{Hence number of moles of gas } \mu = \frac{44.8}{22.4} = 2$$

Since the volume of cylinder is fixed,

$$\text{Hence } (\Delta Q)_V = \mu C_v \Delta T$$

$$\therefore 2 \times \frac{3}{2} R \times 10 = 30R \left[\because (C_v)_{\text{mono}} = \frac{3}{2} R \right]$$

128 (b)

The ideal gas law is the equation of state of an ideal gas. The state of an amount of gas is determined by its pressure, volume and temperature. The equation has the form

$$pV = nRT$$

where, p is pressure, V the volume, n the number of moles, R the gas constant and T the temperature.

$$\therefore \frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2}$$

Given,

$$p_1 = 200 \text{ kPa}, V_1 = V, T_1 = 273 + 22 = 295 \text{ K}, V_2 = V$$

$$T_2 = 273 + 42 = 315 \text{ K}$$

$$\frac{200 \times V}{295} = \frac{p_2 \times 1.02 V}{315}$$

$$\Rightarrow p_2 = \frac{200 \times 315}{295 \times 1.02}$$

$$p_2 = 209 \text{ kPa}$$

129 (c)

$$PV = \mu RT \Rightarrow \mu = \frac{PV}{RT} = \frac{21 \times 10^4 \times 83 \times 10^{-3}}{8.3 \times 300} = 7$$

130 (b)

An ideal gas is a gas which satisfying the assumptions of the kinetic energy.

131 (d)

$$P = \frac{2}{3} E$$

132 (b)

$\gamma = 7/5$ for a diatomic gas

134 (c)

$$v_{rms} \propto \frac{1}{\sqrt{M}} \Rightarrow \frac{v_{O_2}}{v_{H_2}} = \sqrt{\frac{M_{H_2}}{M_{O_2}}} \Rightarrow \frac{C}{v_{H_2}} = \sqrt{\frac{2}{32}} = \frac{1}{4}$$

$$\Rightarrow v_{H_2} = 4C \text{ cm/s}$$

135 (a)

$$P \propto T \Rightarrow \frac{P_1}{P_2} = \frac{T_1}{T_2} \Rightarrow \frac{P_2 - P_1}{P_1} = \frac{T_2 - T_1}{T_1}$$

$$\Rightarrow \left(\frac{\Delta P}{P}\right)\% = \left(\frac{251 - 250}{250}\right) \times 100 = 0.4\%$$


136 (b)

$$v_{rms} \propto \sqrt{T} \Rightarrow \frac{(v_{rms})_2}{(v_{rms})_1} = \sqrt{\frac{T_2}{T_1}}$$

$$\Rightarrow 2 = \sqrt{\frac{T_2}{300}} \Rightarrow T_2 = 1200 \text{ K} = 927^\circ \text{C}$$

137 (d)

Figure shows the particles each moving with same speed v but in different directions. Consider any two particles having angle θ between directions of their velocities



$$\text{Then, } \vec{v}_{rel} = \vec{v}_B - \vec{v}_A$$

$$i.e., v_{rel} = \sqrt{v^2 + v^2 - 2vv \cos \theta}$$

$$\Rightarrow v_{rel} = \sqrt{2v^2(1 - \cos \theta)} = 2v \sin(\theta/2)$$

So averaging v_{rel} over all pairs

$$\dot{v}_{rel} = \frac{\int_0^{2\pi} v_{rel} d\theta}{\int_0^{2\pi} d\theta} = \frac{\int_0^{2\pi} 2v \sin(\theta/2) d\theta}{\int_0^{2\pi} d\theta} = \frac{2v \times 2[-\cos(\theta/2)]_0^{2\pi}}{2\pi}$$

$$\Rightarrow \dot{v}_{rel} = (4v/\pi) > v \quad [\text{as } 4/\pi > 1]$$

138 (b)

Since volume is constant,

$$\text{Hence } \frac{P_1}{P_2} = \frac{T_1}{T_2} \Rightarrow \frac{1}{3} = \frac{(273+30)}{T_2}$$

$$\Rightarrow T_2 = 909 \text{ K} = 636^\circ \text{C}$$

139 (d)

The value of $\frac{pV}{T}$ for one mole of an ideal gas

$$= \text{gas constant}$$

$$= 2 \text{ cal mol}^{-1} \text{K}^{-1}$$

140 (d)

Mean kinetic energy for μ mole gas $\dot{\mu} \cdot \frac{f}{2} RT$

$$\therefore E = \mu \frac{7}{2} RT = \left(\frac{m}{M}\right) \frac{7}{2} NkT = \frac{1}{44} \left(\frac{7}{2}\right) NkT$$

$$\dot{\mu} \frac{7}{88} NkT \quad [\text{As } f = 7 \wedge M = 44 \text{ for } CO_2]$$

141 (c)

$$V \propto T \Rightarrow \frac{V_1}{V_2} = \frac{T_1}{T_2} \Rightarrow \frac{V}{2V} = \frac{(273+27)}{T_2} = \frac{300}{T_2}$$

$$\Rightarrow T_2 = 600 \text{ K} = 327^\circ \text{C}$$

142 (a)

$$C_p - C_v = R = 2 \cdot \frac{\text{cal}}{\text{g-mol-K}}$$

Which is correct for option (a) and (b). Further the

ratio $\frac{C_p}{C_v} (\dot{\gamma})$ should be equal to some standard value

corresponding to that of either, mono, di, or triatomic gases. From this point of view option (a) is correct

$$\text{because } \left(\frac{C_p}{C_v}\right)_{\text{mono}} = \frac{5}{3}$$

143 (a)

$$v_{rms} = \sqrt{\frac{3RT}{M}} \Rightarrow T \propto M \quad [\because v_{rms}, R \rightarrow \text{constant}]$$

$$\frac{T_{H_2}}{T_{O_2}} = \frac{M_{H_2}}{M_{O_2}} = \frac{T_{H_2}}{(273+47)} = \frac{2}{32} \Rightarrow T_{H_2} = 20 \text{ K}$$

144 (c)

Molecules of ideal gas behaves like perfectly elastic rigid sphere

145 (d)

$$PV = nrT \Rightarrow P \propto m \dot{\mu} \text{ constant}$$

$$\Rightarrow \frac{m_1}{m_2} = \frac{P_1}{P_2} = \frac{10}{m_2} = \frac{10^7}{2.5 \times 10^6} \Rightarrow m_2 = 2.5 \text{ kg}$$

Hence mass of the gas taken out of the cylinder

$$\dot{\mu} 10 - 2.5 = 7.5 \text{ kg}$$

147 (b)

$$(\Delta Q)_p = \mu C_p \Delta T \text{ and } (\Delta Q)_v = \mu C_v \Delta T$$

$$\Rightarrow \frac{(\Delta Q)_v}{(\Delta Q)_p} = \frac{C_v}{C_p} = \frac{\frac{3}{2}R}{\frac{5}{2}R} = 3/5$$

$$\left[\because (C_v)_{\text{mono}} = \frac{3}{2}R, (C_p)_{\text{mono}} = \frac{5}{2}R \right]$$

$$\Rightarrow (\Delta Q)_v = \frac{3}{5} \times (\Delta Q)_p = \frac{3}{5} \times 210 = 126 \text{ J}$$

148 (d)

Root mean square velocity of gas molecules

$$v_{rms} = \sqrt{\frac{3RT}{M}}$$

$$v_{rms} \propto \frac{1}{\sqrt{M}}$$

$$\frac{v_{O3}}{v_{O2}} = \sqrt{\frac{M_{O2}}{M_{O3}}}$$

Here, $M_{O2} = 32, M_{O3} = 48$

$$\therefore \frac{v_{O3}}{v_{O2}} = \sqrt{\frac{32}{48}} = \frac{\sqrt{2}}{\sqrt{3}}$$

149 (d)

$$v_{rms} = \sqrt{\frac{3RT}{M}} \Rightarrow v_{rms} \propto \frac{1}{\sqrt{M}}$$

150 (c)

For mono atomic gas, C_V is constant $\left(\frac{3}{2}R\right)$. It

doesn't vary with temperature

151 (a)

$$PV = \mu RT = \frac{m}{M} RT$$

$$\Rightarrow \frac{PV}{T} \propto \frac{1}{M} \text{ [molecule mass]}$$

$$\text{From graph } \left(\frac{PV}{T}\right)_A < \left(\frac{PV}{T}\right)_B < \left(\frac{PV}{T}\right)_C$$

$$\Rightarrow M_A > M_B > M_C$$

152 (d)

$$\frac{\Delta Q}{\Delta t} = KA \left(\frac{\Delta T}{\Delta x}\right) = K\pi r^2 \left(\frac{\Delta T}{l}\right) \propto \frac{r^2}{l}$$

As $\frac{r^2}{l}$ is maximum for (d), it is the correct choice.

153 (a)

Internal energy of the gas remains constant, hence

$$T_2 = T$$

$$\text{Using } p_1 V_1 = p_2 V_2$$

$$p \cdot \frac{V}{2} = p_2 V_2$$

$$p_2 = \frac{p}{2}$$

154 (d)

The square root of \dot{v}^2 is called the root mean square velocity (rms) speed of the molecules.

$$v_{rms} = \sqrt{\dot{v}^2} = \sqrt{\frac{v_1^2 + v_2^2 + v_3^2 + v_4^2}{4}}$$

$$\dot{v} = \sqrt{\frac{(1)^2 + (2)^2 + (3)^2 + (4)^2}{4}}$$

$$\dot{v} = \sqrt{\frac{1+4+9+16}{4}} = \sqrt{\frac{30}{4}} = \sqrt{\frac{15}{2}} \text{ km s}^{-1}$$

155 (b)

Using Newton's law of cooling,

$$\log \frac{\theta_2 - \theta_0}{\theta_1 - \theta_0} = -Kt$$

$$\text{Log } \frac{40 - \theta_0}{50 - \theta_0} = -K \times 5 \quad \dots (i)$$

$$\text{Log } \frac{33.33 - \theta_0}{40 - \theta_0} = -K \times 5 \quad \dots (ii)$$

From Eqs.(i) and (ii),

$$\frac{40 - \theta_0}{50 - \theta_0} = \frac{33.33 - \theta_0}{40 - \theta_0}$$

On solving, we get

$$\theta_0 = 19.95^\circ\text{C} \approx 20^\circ\text{C}$$

157 (c)

- The dotted line in the diagram shows that there is no derivation in the value of $\frac{pV}{nT}$ for different temperature $T_1 \wedge T_2$ for increasing pressure so, this gas behaves ideally. Hence, dotted line corresponds to 'ideal' gas behavior.
- At high temperature, the derivation of the gas is less and at low temperature the derivation of gas is more. In the graph, derivation for T_2 is greater than for T_1 . Thus, $T_1 > T_2$
- Since, the two curves intersect at dotted line so, the value of $\frac{pV}{nT}$ at that point on the y-axis is same for all gases.

158 (d)

Since $v_{rms} \propto \sqrt{T}$. Also mean square velocity $\dot{v}^2 = v_{rms}^2$

159 (b)

$$v_{rms} \propto \frac{1}{\sqrt{M}} \Rightarrow V_H > V_N > V_O [\because M_H < M_N < M_O]$$

160 (b)

$$P_f = 2p + \dot{p}$$

Saturated vapour pressure will not change if temperature remains constant.

161 (c)

Kinetic energy \propto Temperature

162 (d)

$$PV = nRT$$

$$\Rightarrow PV = \frac{\omega}{M} RT$$

$$\frac{PM}{RT} = \frac{\omega}{V} = e$$

$$\Rightarrow e = \frac{PM}{RT} = \frac{P \times m \times N_A}{RT} = \frac{Pm}{\left(\frac{R}{N_A}\right)T} = \frac{Pm}{kT}$$

163 (b)

Thermal energy corresponds to internal energy

$$\text{Mass} = 1 \text{ kg}$$

$$\text{Density} = 4 \text{ kg m}^{-3}$$

$$\text{Volume} = \frac{\text{Mass}}{\text{Density}} = \frac{1}{4} \text{ m}^3$$

$$\text{Pressure} = 8 \times 10^4 \text{ N m}^{-2}$$

$$\therefore \text{Internal energy} = \frac{5}{2} p \times V = 5 \times 10^4 \text{ J}$$

164 (b)

$$V_t = V_0(1 + \alpha t) = 0.5 \left(1 + \frac{1}{273} \times 819 \right) = 2 \text{ litre} = 2 \times 10^{-3} \text{ m}^3$$

165 (c)

$$\text{Here, } m = 10 \text{ g} = 10^{-2} \text{ kg}$$

$$v = 300 \text{ m s}^{-1}, \theta = ? \text{ } ^\circ\text{C}, = 150 \text{ J} \cdot \text{kg}^{-1} \text{ K}^{-1}$$

$$Q = \frac{50}{100} \left(\frac{1}{2} m v^2 \right) = \frac{1}{4} \times 10^{-2} (300)^2 = 225 \text{ J}$$

$$\text{From } Q = cm\theta$$

$$\theta = \frac{Q}{cm} = \frac{225}{150 \times 10^{-2}} = 150 \text{ } ^\circ\text{C}$$

166 (a)

At constant temperature

$$PV = \text{constant}$$

$$\Rightarrow \frac{P_1}{P_2} = \frac{V_2}{V_1} = \frac{70}{120} = \frac{V_2}{1200} \Rightarrow V_2 = 700 \text{ ml}$$

167 (d)

$$P \propto \frac{1}{V} \Rightarrow \frac{V_2}{V_1} = \frac{P_1}{P_2} = \frac{100}{105} \Rightarrow V_2 = \frac{100}{105} V_1 = 0.953 V_1$$

$$\% \text{ change in volume} = \frac{V_1 - V_2}{V_1} \times 100$$

$$= \frac{V_1 - 0.953 V_1}{V_1} \times 100 = 4.76 \%$$

168 (a)

$$\text{Average kinetic energy } E = \frac{f}{2} kT = \frac{3}{2} kT$$

$$\Rightarrow E = \frac{3}{2} \times (1.38 \times 10^{-23}) (273 + 30) = 6.27 \times 10^{-21} \text{ J}$$

$$= 0.039 \text{ eV} < 1 \text{ eV}$$

169 (c)

$$\therefore C_p - C_v = R$$

$$\text{Fractional part of heat energy} = \frac{C_p - R}{C_p}$$

$$= \frac{\frac{7}{2} R - R}{\frac{7}{2} R} = \frac{5}{7}$$

170 (c)

RMS velocity doesn't depend upon pressure, it depends upon temperature only,

$$\text{ie., } v_{\text{rms}} \propto \sqrt{T}$$

$$\Rightarrow \frac{v_1}{v_2} = \sqrt{\frac{T_1}{T_2}} \Rightarrow \frac{200}{v_2} = \sqrt{\frac{(273+27)}{(273+127)}} = \sqrt{\frac{300}{400}}$$

$$\Rightarrow v_2 = \frac{400}{\sqrt{3}} \text{ m/s}$$

171 (a)

$$\frac{F}{2} n_1 k T_1 + \frac{F}{2} n_2 k T_2 + \frac{F}{2} n_3 k T_3$$

$$= \frac{F}{2} (n_1 + n_2 + n_3) k T$$

$$T = \frac{n_1 T_1 + n_2 T_2 + n_3 T_3}{n_1 + n_2 + n_3}$$

172 (a)

$$\text{As } \rho - \rho_0(1 - \gamma \Delta T)$$

$$\therefore 9.7 = 10(1 - \gamma \times 100)$$

$$\frac{9.7}{10} = 1 - \gamma \times 100$$

$$\gamma \times 100 = 1 - \frac{9.7}{10} = \frac{0.3}{10} = 3 \times 10^{-2}$$

$$\gamma = 3 \times 10^{-4} \therefore \alpha = \frac{1}{3} \gamma = 10^{-4} \text{ } ^\circ\text{C}^{-1}$$

174 (b)

Let the temperature of junction be Q . In equilibrium, rate of flow of heat through rod 1 = sum of rate of flow of heat through rods 2 and 3.

$$\left(\frac{dQ}{dt} \right)_1 = \left(\frac{dQ}{dt} \right)_2 + \left(\frac{dQ}{dt} \right)_3$$

$$KA \frac{(\theta - 0)}{l} = \frac{KA(90^\circ - \theta)}{l} + \frac{KA(90^\circ - \theta)}{l}$$

$$\theta = 2(90^\circ - \theta)$$

$$3\theta = 180^\circ, \theta = \frac{180^\circ}{3} = 60^\circ$$

175 (a)

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$\frac{(P+h\rho g)1.0}{273+12} = \frac{P \cdot V_2}{273+35}$$

$$V_2 = 5.4 \text{ cm}^3$$

176 (d)

Average kinetic energy \propto Temperature

$$\Rightarrow \frac{E_1}{E_2} = \frac{T_1}{T_2} \Rightarrow \frac{100}{E_2} = \frac{300}{450} \Rightarrow E_2 = 150 \text{ J}$$

177 (a)

Let p_1 and p_2 are the initial and final pressures of the gas filled in A. Then

$$p_1 = \frac{n_A RT}{V} \wedge p_2 = \frac{n_A RT}{2V}$$

$$\Delta p = p_2 - p_1 = \frac{-n_A RT}{2V}$$

$$\therefore -\left(\frac{m_A}{M}\right) \frac{RT}{2V} \quad \dots(i)$$

where M is the atomic weight of the gas.

$$\text{Similarly, } 1.5 \Delta p = -\left(\frac{m_B}{M}\right) \frac{RT}{2V}$$

... (ii)

Dividing Eq. (ii) by Eq. (i), we get

$$1.5 = \frac{m_B}{m_A} \sqrt{\frac{3}{2}} = \frac{m_B}{m_A}$$

or

$$3m_A = 2m_B$$

178 (c)

$$\text{From } \frac{\Delta Q}{\Delta t} = KA \left(\frac{\Delta T}{\Delta x} \right)$$

$$\Delta t = \frac{\Delta Q \Delta x}{KA (\Delta T)}$$

In arrangement (b), A is doubled and Δx is halved.

$$\therefore \Delta t \rightarrow \frac{1/2}{2} \rightarrow \frac{1}{4} \text{ time}$$

$$\text{ie, } \frac{1}{4} \times 4 \text{ min} = 1 \text{ min}$$

179 (b)

Here, $m = 0.1 \text{ kg}$, $h_1 = 10 \text{ m}$, $h_2 = 5.4 \text{ m}$

$c = 460 \text{ J-kg}^{-1} \text{ }^\circ\text{C}^{-1}$, $g = 10 \text{ m s}^{-2}$, $\theta = ?$

Energy dissipated, $Q = mg(h_1 - h_2)$

$$= 0.1 \times 10(10 - 5.4) = 4.6 \text{ J}$$

From $Q = cm\theta$

$$\theta = \frac{Q}{cm} = \frac{4.6}{460 \times 0.1} = 0.1 \text{ }^\circ\text{C}$$

180 (b)

Root mean square speed

$$v_{rms} \propto \frac{1}{\sqrt{\rho}}$$

$$\therefore \frac{v_{rms1}}{v_{rms2}} = \sqrt{\frac{\rho_2}{\rho_1}}$$

$$\text{Given, } \frac{\rho_1}{\rho_2} = \frac{9}{8}$$

$$\Rightarrow \frac{v_{rms1}}{v_{rms2}} = \sqrt{\frac{8}{9}} = \frac{2\sqrt{2}}{3}$$

181 (c)

$$v_{rms} \propto \frac{1}{\sqrt{M}} \Rightarrow \frac{v_1}{v_2} = \sqrt{\frac{M_2}{M_1}}$$

$$\therefore \frac{1}{\sqrt{2}} = \sqrt{\frac{M_2}{32}} \Rightarrow M_2 = 16. \text{ Hence the gas is } CH_4$$

182 (a)

$$\text{No. of moles } n = \frac{m}{\text{molecular weight}} = \frac{5}{32}$$

So, from ideal gas equation

$$pV = nRT$$

$$\Rightarrow pV = \frac{5}{32} RT$$

183 (a)

According to Avogadro's hypothesis

184 (c)

$$\text{Pressure of gas A, } P_A = \frac{125 \times 0.6}{1000} = 0.075 \text{ atm}$$

$$\text{Pressure of gas B, } P_B = \frac{150 \times 0.8}{100} = 0.120 \text{ atm}$$

Hence, by using Dalton's law of pressure

$$P_{\text{mixture}} = P_A + P_B = 0.075 + 0.120 = 0.195 \text{ atm}$$

185 (a)

Average speed (v_{av}) of gas molecules is

$$v_{av} = \sqrt{\frac{8RT}{\pi M}}$$

where R is gas constant and M the molecular weight.

Given, $v_1 = v$, $M_1 = 64$, $v_2 = 4v$

$$\therefore \frac{v_1}{v_2} = \sqrt{\frac{M_2}{M_1}}$$

$$\frac{v}{4v} = \sqrt{\frac{M_2}{64}}$$

$$\Rightarrow M_2 = \frac{64}{16} = 4$$

Hence, the gas is helium (molecular mass 4).

186 (b)

Heat added to helium during expansion

$$H = nC_v \Delta T = 8 \times \frac{3}{2} R \times 30 \left(C_v \text{ for monoatomic gas} \right)$$

$$= 360 R$$

$$= 360 \times 8.31 \text{ J} \quad (R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1})$$

$$\approx 3000 \text{ J}$$

187 (c)

In Vander Waal's equation $\left(P + \frac{a}{V^2} \right) (V - b) = RT$

a represents intermolecular attractive force and b represents volume correction

188 (b)

$$C_p - C_v = R \Rightarrow C_p = R + C_v = R + \frac{f}{2} R$$

$$R + \frac{3}{2} R = \frac{5}{2} R$$

189 (d)

It is because of their low densities

190 (d)

Kinetic energy of a gas molecule

$$E = \frac{3}{2} kT$$

where k is Boltzmann's constant.

$\therefore E \propto T$



$$\text{or } \frac{E_1}{E_2} = \frac{T_1}{T_2} \Rightarrow \frac{E}{(E/2)} = \frac{300}{T_2}$$

$$\text{or } T_2 = 150 \text{ K}$$

$$T_2 = 150 - 273 = -123 \text{ }^\circ\text{C}$$

191 (c)

On keeping the temperature of the ends of tube at 0°C and 273°C .



Applying ideal gas equation

$$\frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2} = \frac{p_3 V_3}{T_3}$$

$$\frac{76 \times 45}{(273+31)} = \frac{p_2 \times l}{(273+0)} = \frac{p_3(90-l)}{273+273}$$

$$\frac{76 \times 45}{304} = \frac{p_2 \times l}{273} = \frac{p_3(90-l)}{546}$$

I II III

From II and III

$$\frac{p_2 \times l}{273} = \frac{p_3(90-l)}{546}$$

(Mercury column is at rest, so pressure difference $p_2 - p_3 = 0 \Rightarrow p_2 = p_3$)

$$\therefore \frac{p_2 \times l}{273} = \frac{p_2(90-l)}{546}$$

$$\Rightarrow 2l = 90 - l \Rightarrow l = 30 \text{ cm}$$


From I and II

$$\frac{76 \times 45}{304} = \frac{p_2 \times 30}{273}$$

$$\Rightarrow p_2 = \frac{76 \times 45 \times 273}{30 \times 304}$$

$$p_2 = 102.4$$

192 (c)



$$PV = \mu RT$$

$$\Rightarrow V \propto \frac{T}{P} \text{ and } R \text{ are fixed}$$

Since, T increases rapidly and P increases slowly

thus volume of the gas increases

193 (b)

$$v_{av} \propto \frac{1}{\sqrt{M}} \Rightarrow \frac{v_{He}}{v_H} = \sqrt{\frac{M_H}{M_{He}}} = \sqrt{\frac{1}{4}} = \frac{1}{2} \Rightarrow v_{He} = \frac{v_H}{2}$$

194 (b)

$$v_{rms} = \sqrt{\frac{3RT}{M}} = \sqrt{\frac{3 \times 8.3 \times 300}{28 \times 10^{-3}}} = 517 \text{ m/s}$$

195 (d)

Thermal equilibrium implies that the temperature of gases is same. Hence Boyle's law is applicable i.e.

$$P_a V_a = P_b V_b$$

196 (d)

$$C_v = \frac{5}{2} R \wedge C_p = \frac{7}{2} R$$

$$\therefore \gamma = \frac{C_p}{C_v} = \frac{7}{5}$$

197 (c)

Moist and hot air being lighter rises up and leaves the room through the ventilator near the roof and fresh air rushes into the room through the doors.

198 (d)

Root mean square velocity of molecule in left part

$$v_{rms} = \sqrt{\frac{3KT}{m_L}}$$

Mean or average speed of molecule in right part


$$v_{av} = \sqrt{\frac{8KT}{\pi m_R}}$$

$$\text{According to problem } \sqrt{\frac{3KT}{m_L}} = \sqrt{\frac{8KT}{\pi m_R}}$$

$$\Rightarrow \frac{3}{m_L} = \frac{8}{\pi m_R} \Rightarrow \frac{m_L}{m_R} = \frac{3\pi}{8}$$

199 (c)

Temperature of the gas is concerned only with its disordered motion. It is not concerned with its ordered motion



200 (c)

$$Y_{max} = \frac{\frac{\mu_1 \gamma_1}{\gamma_1 - 1} + \frac{\mu_2 \gamma_2}{\gamma_2 - 1}}{\frac{\mu_1}{\gamma_1 - 1} + \frac{\mu_2}{\gamma_2 - 1}}$$

$$\frac{1 \times \frac{5}{3} + 1 \times \frac{7}{5}}{\frac{\frac{5}{3} - 1}{3} + \frac{\frac{7}{5} - 1}{5}} = \frac{3}{2} = 1.5$$

201 (d)

$$E = \frac{3}{2} RT = \frac{3}{2} \times 8.31 \times 273 = 3.4 \times 10^3 \text{ J}$$

202 (b)

Given, $p_1 = 100 \text{ mm}$, $V_1 = 200 \text{ mL}$ \wedge $p_2 = 400 \text{ mm}$

From Boyle's Law

$$p_1 V_1 = p_2 V_2$$

$$V_2 = \frac{p_1 V_1}{p_2}$$

$$\therefore \frac{100 \times 200}{400}$$

$$V_2 = 50 \text{ mL}$$

Volume of 2 mol gas = $2 \times 50 = 100 \text{ mL}$

203 (b)

$$v_{rms} = \sqrt{\frac{3RT}{M}} \Rightarrow v_{rms}^2 \propto T$$

204 (b)

$$(C_p)_{mix} = \frac{\mu_1 C_{p_1} + \mu_2 C_{p_2}}{\mu_1 + \mu_2} (C_{p_1}(\text{He}) = \frac{5}{2} R \wedge C_{p_2}(\text{H}_2) =$$

$$1 \times \frac{5}{2} R + 1 \times \frac{7}{2} R$$

$$\therefore \frac{1 \times \frac{5}{2} R + 1 \times \frac{7}{2} R}{1 + 1} = 3R = 3 \times 2 = 6 \text{ cal/mol} \cdot ^\circ\text{C}$$

\therefore Amount of heat needed to raise the temperature from 0°C to 100°C

$$(\Delta Q)_p = \mu C_p \Delta T = 2 \times 6 \times 100 = 1200 \text{ cal}$$

205 (c)

The average velocity

$$v_{av} = \frac{v_1 + v_2 + v_3 + \dots + v_n}{N}$$

$$\therefore \frac{1 + 3 + 5 + 7}{4} = 4 \text{ km/s}$$

Root mean square velocity

$$v_{rms} = \sqrt{\frac{v_1^2 + v_2^2 + v_3^2 + \dots + v_n^2}{N}}$$

$$\therefore \sqrt{\frac{1 + (3)^2 + (5)^2 + (7)^2}{4}}$$

$$\therefore \sqrt{21} = 4.583 \text{ km/s}$$

Difference between average velocity and root

mean square velocity
 $= 4.583 \times 10^4$
 $= 0.583 \text{ km/s}$

206 (c)

$$V \propto T \Rightarrow \frac{V_1}{V_2} = \frac{T_1}{T_2}$$

$$\Rightarrow \frac{V}{V_2} = \frac{(273+27)}{(273+327)} = \frac{300}{600} = \frac{1}{2} \Rightarrow V_2 = 2V$$

207 (c)

For a closed system, the total number of moles remains constant. So

$$p_1 V = n_1 R T_1 \wedge p_2 V = n_2 R T_2$$

$$\therefore p(2V) = (n_1 + n_2) R T$$

$$\therefore \frac{p}{T} = \frac{(n_1 + n_2)}{2} R = \frac{1}{2} \left[\frac{P_1}{T_1} + \frac{P_2}{T_2} \right]$$

$$= \frac{1}{2} \left[\frac{p_1 T_2 + p_2 T_1}{T_1 T_2} \right]$$

208 (a)

$$\text{Most probable speed } v_{mp} = \sqrt{\frac{2kT}{m}} \Rightarrow \frac{1}{2} m v_{mp}^2 = kT$$

209 (a)

$$\text{As } dQ = dU + dW$$

$$\therefore dU = dQ - dW = 2240 - 168$$

$$= 2072 \text{ J}$$

210 (c)

The root mean square velocity

$$v_{rms} = \sqrt{\frac{3RT}{M}}$$

where R is gas constant, T the temperature and M the molecular weight.

$$\text{Given, } v_{He} = v_H, T_H = 273 \text{ K}, M_H = 2, M_{He} = 4$$

$$\therefore \frac{v_H}{v_{He}} = \sqrt{\frac{T_H}{T_{He}} \times \frac{M_{He}}{M_H}}$$

$$\therefore 1 = \sqrt{\frac{273}{T_{He}} \times \frac{4}{2}}$$

$$\Rightarrow T_{He} = 546 \text{ K}$$

$$\text{In } ^\circ\text{C, } T_{He} = (546 - 273) ^\circ\text{C} = 273 ^\circ\text{C}$$

212 (b)

The molecules of a gas are in a state of random motion. They continuously collide against the walls of the container. Even at ordinary temperature and pressure, the number of molecular collisions with walls is very large. During each collision, certain momentum is

transferred to the walls of the container. The pressure exerted by the gas is due to continuous bombardment of gas molecules against the walls of the container. Due to this continuous bombardment, the walls of the container experience a continuous force which is equal to the total momentum imparted to the walls per second. The average force experienced per unit area of the walls container determines the pressure exerted by the gas. This should be clear from the fact that although the molecular collisions are random the pressure remains constant.

213 (c)

$$\text{Given, } p T^2 = \text{constant}$$

$$\therefore \left(\frac{nRT}{V} \right) T^2 = \text{constant}$$

$$\text{or } T^3 V^{-1} = \text{constant}$$

Differentiating the equation, we get

$$\frac{3T^2}{V} \cdot dT - \frac{T^3}{V^2} \cdot dV = 0$$

$$\text{or } 3 \cdot dT = \frac{T}{V} \cdot dV$$


$$\text{From the equation, } dV = V \cdot \gamma \cdot dT$$

$\gamma = \text{coefficient of volume expansion of gas}$

$$\therefore \gamma = \frac{dV}{V \cdot dT} = \frac{3}{T}$$

215 (b)

Pressure will be less in front portion of the compartment because in accelerated frame molecules will feel pseudo force in backward direction. Also density of gas will be more in the back portion



216 (a)

$$v_{rms} \propto \sqrt{T}$$

$$\Rightarrow \frac{v_1^2}{v_2^2} = \frac{T_1}{T_2}$$

$$\Rightarrow \frac{v^2}{2v^2} = \frac{273}{T_2}$$

$$\Rightarrow T_2 = 1092 \text{ K}$$

$$= 819^\circ\text{C}$$

217 (c)

Average velocity of gas molecule is

$$v_{av} = \sqrt{\frac{8RT}{\pi M}} = v_{av} \times \frac{1}{\sqrt{M}}$$

$$\Rightarrow \sqrt{C_H} > \frac{1}{\sqrt{M_{He}}} > \frac{1}{\sqrt{M_H}} = \sqrt{\frac{4}{1}} = 2 \Rightarrow \sqrt{C_H} \geq 2 < C_H$$

218 (c)

$$\mu = \mu_1 + \mu_2$$

$$\frac{P(2V)}{RT_1} = \frac{P'V}{RT_1} + \frac{P'V}{RT_2} \Rightarrow \frac{2P}{RT_1} = \frac{P'}{R} \left[\frac{T_2 + T_1}{T_1 T_2} \right]$$

$$P' = \frac{2PT_2}{(T_1 + T_2)} = \frac{2 \times 1 \times 600}{(300 + 600)} = \frac{4}{3} \text{ atm}$$

219 (c)

$$C_V = \frac{R}{(\gamma - 1)} \Rightarrow \gamma = 1 + \frac{R}{C_V} = 1 + \frac{R}{\frac{3}{2}R} = \frac{5}{3}$$

220 (b)

$$v_{rms} = \sqrt{\frac{3P}{\rho}} = \sqrt{\frac{3PV}{m}} \Rightarrow v_{rms} \propto \sqrt{\frac{P}{m}}$$

$$\Rightarrow \frac{v_1}{v_2} = \sqrt{\frac{P_1}{P_2} \times \frac{m_2}{m_1}} = \frac{v}{2v} = \sqrt{\frac{P_0}{P_2} \times \frac{m/2}{m}} \Rightarrow P_2 = 2P_0$$

221 (a)

$$\text{Kinetic energy for 1 mole gas } E = \frac{f}{2} RT$$

$$\Rightarrow E_{\text{Translation}} = \frac{3}{2} RT$$

[∵ For all gases translational degree of freedom $f = 3$]

222 (c)

$$PV = \mu RT \text{ [Gas equation]} \Rightarrow PV \propto T$$

223 (b)

Neglecting bond length, the volume of an oxygen molecule has been taken as 2 times that of one oxygen atom.

In 22.4 litres i.e., $22.4 \times 10^{-3} \text{ m}^3$, there are $N_A = 6.23 \times 10^{23}$ molecules

Total volume of oxygen molecules $2 \times \frac{4}{3} \pi r^3 \times N_A$

$22.4 \times 10^{-3} \text{ m}^3$ is occupied by N_A molecules


∴ Fraction of volume occupied

$$\frac{2 \times \frac{4}{3} \times \pi \times (1.5 \times 10^{-10})^3 \times 6.2 \times 10^{23}}{(22.4 \times 10^{-3})} = 8 \times 10^{-4}$$

224 (c)

No change, because rms velocity of gas depends upon temperature only

225 (c)



$$PV = \mu RT = \frac{m}{M} RT$$

For 1st graph,

$$P = \frac{m_1}{M} \frac{RT}{V_1} \dots (i)$$

For 2nd graph,

$$P = \frac{m_2}{M} \frac{RT}{V_2} \dots (ii)$$

Equating the two, we get, $\frac{m_1}{m_2} = \frac{V_1}{V_2} \Rightarrow m \propto V$

As $V_2 > V_1 \Rightarrow m_1 < m_2$

226 (a)

$$PV = \mu RT \Rightarrow PV \propto T$$

If P and V are doubled then T becomes four times, i.e.,

$$T_2 = 4T_1 = 4 \times 100 = 400 \text{ K}$$

227 (b)

Ideal gas equation can be written as

$$pV = nRT$$

...(i)

From Eq. (i), we have

$$\frac{n}{V} = \frac{p}{RT} = \text{constant}$$

So, at constant pressure and temperature, all gases will contain equal number of molecules per unit volume.

228 (b)

RMS velocity is given by

$$v = \sqrt{\frac{3kT}{m}} \Rightarrow v^2 = \frac{3kT}{m}$$

For a gas, k and m are constants.

$$\therefore \frac{v^2}{T} = \text{constant}$$

229 (b)


CO is diatomic gas, for diatomic gas

$$C_p = \frac{7}{2}R \text{ and } C_v = \frac{5}{2}R \Rightarrow \gamma_{di} = \frac{C_p}{C_v} = \frac{7R/2}{5R/2} = 1.4$$

230 (a)

When gas is filled in a closed container, it exerts pressure on the walls of the vessel.

According to kinetic theory this pressure is developed due to the collisions of the moving molecules on the walls of the vessels. Whenever a molecule collides with the wall, it returns with changed momentum and an equal momentum is transferred to the wall. According to Newton's law of motion, the rate of change of momentum of the ball is equal to the force exerted on the wall. Since, the gas contains a large number of molecules which are colliding with the walls of the vessel, they exert a steady force on the walls. This force measured per unit area gives pressure, which is same as the molecules are moving in horizontal direction with constant acceleration.



231 (a)

Part (i)	Part (ii)
$P, 5V$	$10P, V$

When the piston is allowed to move the gases are kept separated but the pressure has to be equal.

($P_1 = P_2$) and final volume x and $(6V - x)$, the no. of moles are same in initial and final position at each part.

$$\therefore P_1 = P_2 P_v = n_1 RT$$

$$\frac{n_1 RT}{x} = \frac{n_2 RT}{6V - x} \Rightarrow n_1 = \frac{5PV}{RT}$$

$$\frac{n_1}{x} = \frac{n_2}{6V - x} \Rightarrow n_2 = \frac{10PV}{RT}$$

$$\Rightarrow \frac{5PV}{xRT} = \frac{10PV}{(6V - x)RT} \Rightarrow \frac{1}{x} = \frac{2}{6V - x}$$

$$\Rightarrow 6V - x = 2x \Rightarrow x = 2V \text{ and}$$

$$6V - x = 6V - 2V = 4V$$

$$\therefore (2V, 4V)$$

233 (c)

Kinetic energy \propto Temperature. Hence if temperature is doubled, kinetic energy will also be doubled

234 (c)

The average kinetic energy of monoatomic gas

$$\text{molecule is } K = \frac{3}{2} k_B T$$

Where k_B is the Boltzmann constant and T is the temperature of the gas in kelvin

$$K = \frac{3}{2} \times (1.38 \times 10^{-23} \text{ J K}^{-1}) \times (300 \text{ K})$$

$$\therefore \frac{3 \times (1.38 \times 10^{-23} \text{ J K}^{-1}) \times (300 \text{ K})}{2 \times (1.6 \times 10^{-19} \text{ J/eV})}$$

$$\therefore 3.9 \times 10^{-2} \text{ eV} = 0.039 \text{ eV}$$

235 (a)

If the volume remains constant, then

$$\frac{p_1}{p_2} = \frac{T_1}{T_2}$$

$$\Rightarrow \frac{p}{p + \frac{0.4}{100}p} = \frac{T}{T + 1}$$

$$\text{or } T = 250 \text{ K}$$

236 (a)

From Boyle's law

$$pV = \text{constant}$$

$$\therefore p_1 V_1 = p_2 V_2$$


$$\text{Here, } p_1 = (h+l), V_1 = \frac{4}{3} \pi r^3$$

$$p_2 = l, V_2 = \frac{4}{3} \pi l^3$$

$$\therefore (h+l) \frac{4}{3} \pi r^3 = l \times \frac{4}{3} \pi (3r)^3$$

$$\text{or } h+l = 27l$$

$$\therefore h = 26l$$



237 (d)

Degree of freedom $f = 3$ (Translatory) + 2 (rotatory) + 1 (vibratory) = 6

$$\Rightarrow \frac{C_p}{C_v} = \gamma = 1 + \frac{2}{f} = 1 + \frac{2}{6} = \frac{4}{3} = 1.33$$

238 (c)

In the absence of intermolecular forces, there

will be no stickiness of molecules. Hence, pressure will increase.

239 (a)

At $T = 0 \text{ K}$, $v_{rms} = 0$

240 (c)

The given equation is for 1 g mol gas

241 (c)

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$T_2 = \frac{P_2 V_2}{P_1 V_1} T_1 = \frac{2}{1} \times \frac{3}{1} \times 300 = 1800 \text{ K} = 1527^\circ \text{C}$$

242 (a)

$$\because \theta_1 < \theta_2 \Rightarrow \tan \theta_1 < \tan \theta_2 \Rightarrow \left(\frac{V}{T}\right)_1 < \left(\frac{V}{T}\right)_2$$

$$\text{Form } PV = \mu RT; \frac{V}{T} \propto \frac{1}{P}$$

$$\text{Hence } \left(\frac{1}{P}\right)_1 < \left(\frac{1}{P}\right)_2 \Rightarrow P_1 > P_2$$

243 (d)

$C_p - C_v = R$ and R is constant for all gases

244 (b)

For a real gas the two van der Waal's constants and Boyle's temperature (T_B) are related as

$$T_B = \frac{a}{bR}$$

245 (b)

$$v_{rms} \propto \sqrt{T}$$

246 (d)

r. m. s. velocity does not depend upon pressure

247 (c)

$$E_{av} = \frac{f}{2} kT = \frac{3}{2} \times 1.38 \times 10^{-23} \times 273 = 0.56 \times 10^{-20} \text{ J}$$

248 (c)

$$\text{As } \eta = 1 = \frac{T_2}{T_1}$$

$$\therefore \frac{50}{100} = 1 = \frac{500}{T_1} \Rightarrow T_1 = 1000 \text{ K}$$

$$\text{Again, } \frac{60}{100} = 1 - \frac{T_2}{1000}$$

$$\text{Or } T_2 = 400 \text{ K}$$

249 (a)

Root mean square velocity (v_{rms}), given by

$$v_{rms} = \sqrt{\frac{3RT}{M}}$$

where R is gas constant, T the temperature and M molecular weight.

$$\text{Given, } T_1 = 27^\circ \text{C} = 273 + 27 = 300 \text{ K},$$

$$T_2 = 327^\circ \text{C} = 273 + 327 = 600 \text{ K}$$

$$\therefore \frac{(v_{rms})_1}{(v_{rms})_2} = \sqrt{\frac{300}{600}} = \sqrt{\frac{1}{2}}$$

$$\Rightarrow (v_{rms})_2 = \sqrt{2} (v_{rms})_1$$

Hence, rms speed increases $\sqrt{2}$ times.

251 (d)

Oxygen being a diatomic gas possesses 5 degrees of freedom, 3 translational and 2 rotational.

Argon being monoatomic has 3 translational degrees of freedom.

Total energy of the system

$$\downarrow E_{\text{oxygen}} + E_{\text{argon}}$$

$$\downarrow n_1 f_1 \left(\frac{1}{2} RT\right) + n_2 f_2 \left(\frac{1}{2} RT\right)$$

$$\downarrow 2 \times 5 \times \frac{1}{2} RT + 4 \times 3 \times \frac{1}{2} RT$$

$$\downarrow 5 RT + 6 RT = 11 RT$$

252 (d)

Consider n moles of a gas which undergo isochoric process, *ie*, $V = \text{constant}$. From first law of thermodynamics,

$$\Delta Q = \Delta W + \Delta U$$

...(i)

Here, $\Delta W = 0$ as $V = \text{constant}$

$$\Delta Q = n C_v \Delta T$$

Substituting in Eq. (i), we get

$$\Delta U = n C_v \Delta T$$

...(ii)

Mayer's relation can be written as

$$C_p - C_v = R$$

$$\Rightarrow C_v = C_p - R$$

...(iii)

From Eqs. (ii) and (iii), we have

$$\Delta U = n(C_p - R) \Delta T$$

$$\text{Given, } n = 6, C_p = 8 \text{ cal mol}^{-1} - \text{K}^{-1},$$

$$R = 8.31 \text{ J mol}^{-1} - \text{K}^{-1}$$

$$\approx 2 \text{ cal mol}^{-1} - \text{K}^{-1}$$

$$\text{Hence, } \Delta U = 6(8 - 2)(35 - 20)$$

$$\downarrow 6 \times 6 \times 15 = 540 \text{ cal}$$

253 (d)

Mean kinetic energy of any ideal gas is given by

$$E = \frac{f}{2} RT \text{ which is different gases. (} f \text{ is not same for}$$

all gases)

254 (a)

$$\frac{V_1}{V_2} = \frac{T_1}{T_2}$$

$$\frac{1}{2} = \frac{300}{T_2}$$

$$T_2 = 600 \text{ K} = 600 - 273 = 327^\circ\text{C}$$

$$\Delta t = 327 - 27 = 300^\circ\text{C}$$

255 (c)

Since P and V are not changing, so temperature remains same

256 (c)

$v_{r.m.s.}$ is independent of pressure but depends upon temperature as $v_{r.m.s.} \propto \sqrt{T}$

257 (d)

The main kinetic energy of one mole of gas n degree of freedom.


$$E = \frac{n}{2} RT$$

The mean kinetic energy of one mole of gas per degree of freedom.

$$E' = \frac{E}{n} = \frac{\frac{n}{2} RT}{n}$$

$$E' = \frac{1}{2} RT$$

258 (a)



$$\text{For a gas, } PV = \mu RT = \frac{m}{M} RT$$

$$\text{For graph A, } PV = \frac{m}{M} RT$$

Slope of graph A,

$$\left(\frac{P}{T}\right) = \frac{m}{M} \frac{R}{V} \quad \dots(i)$$

$$\text{For graph B, } PV = \frac{3m}{M} RT$$

Slope of graph B,

$$\left(\frac{P}{T}\right) = \frac{3m}{M} \frac{R}{V} \quad \dots(ii)$$

$$\frac{\text{Slope of curve B}}{\text{Slope of curve A}} = \frac{\frac{3m}{M} \frac{R}{V}}{\frac{m}{M} \frac{R}{V}} = \frac{3}{1}$$

259 (c)

According to law of equipartition of energy, kinetic energy per degree of freedom of a gas molecule is

$$\frac{1}{2}kT$$

260 (c)

For carbon dioxide, number of moles $(n_1) = \frac{22}{44} = \frac{1}{2}$;

molar specific heat of CO_2 at constant volume $C_{V1} = 3R$

For oxygen, number of moles $(n_2) = \frac{16}{32} = \frac{1}{2}$;

molar specific heat of O_2 at constant volume

$$C_{V2} = \frac{5R}{2}$$

Let TK be the temperature of mixture.

Heat lost by $O_2 =$ Heat gained by CO_2 .

$$n_2 C_{V2} \Delta T_2 = n_1 C_{V1} \Delta T_1$$

$$\frac{1}{2} \left(\frac{5}{2} R \right) (310 - T) = \frac{1}{2} \times (3R) (T - 300)$$

$$\text{Or } 1550 - 5T = 6T - 1800$$

$$\text{Or } T = 304.54 \text{ K} = 31.5^\circ\text{C}$$

261 (b)

$$\text{As } dQ = C_p m \Delta T$$

$$\therefore 70 = C_p \times 2(35 - 30)$$

$$C_v = C_p - R$$

$$= 7 - 1.99 = 5.01 \text{ cal mol}^{-1} \text{ } ^\circ\text{C}^{-1}$$

$$\therefore dQ = C_v m \Delta T$$

$$= 5.01 \times 2 \times (35 - 30) = 50.1 \text{ cal}$$

262 (d)

The difference of C_p and C_v is equal to R , not $2R$

264 (b)

Average speed or mean speed of gas molecules

$$\dot{v} = \sqrt{\frac{8RT}{\pi M}} \quad \dot{v} \propto \frac{1}{\sqrt{M}}$$

$$\text{or } \frac{\dot{v}_H}{\dot{v}_{He}} = \sqrt{\frac{M_{He}}{M_H}}$$

$$\text{Here, } M_{He} = 4M_H$$

$$\therefore \frac{\dot{v}_H}{\dot{v}_{He}} = \sqrt{\frac{4}{1}} = 2 \dot{v}_{He} = \frac{1}{2} \dot{v}_H$$

265 (a)

$$C_v = \frac{f}{2} R$$

For diatomic gas $f = 5$

$$\therefore C_v = \frac{5}{2} R$$

266 (c)

$$\frac{E_1}{E_2} = \frac{T_1}{T_2} \Rightarrow \frac{E}{2E} = \frac{(273+27)}{T_2} \Rightarrow T_2 = 600 \text{ K} = 327^\circ\text{C}$$

267 (b)

Here, $V_0 = 10^3 \text{ cc}$

$$\gamma_r = 180 \times 10^{-6} \text{ } ^\circ\text{C}^{-1}$$

$$g = 40 \times 10^{-6} \text{ } ^\circ\text{C}^{-1}, t = 100^\circ\text{C}$$

$$\gamma_a = \gamma_r - g = (180 - 40) 10^{-6}$$

$$V_t = V_0 (1 + 140 \times 10^{-6} \times 10^2)$$

$$= (10^3 + 14) \text{ cc}$$

\therefore Volume of mercury that will overflow

$$= V_t - V_0 = 14 \text{ cc}$$

268 (c)

$$\text{Pressure, } P = \frac{F}{A} = \frac{1}{A} \cdot \frac{\Delta p}{\Delta t} \quad \dot{\text{change in momentum}}$$

269 (c)

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow \frac{1 \times 500}{300} = \frac{0.5 \times V_2}{270} \Rightarrow V_2 = 900 \text{ m}^3$$

270 (c)

For same isotherm; $T \rightarrow$ constant

$$\therefore P \propto \frac{1}{V} \Rightarrow P_1 V_1 = P_2 V_2$$

272 (b)

Given that, $T = 27^\circ\text{C} = 300\text{K}$

$$v_{rms} = 1365 \text{ m s}^{-1}$$

We know that

$$v_{rms} = \sqrt{\frac{3RT}{M}}$$

$$\dot{\text{ }} v_{rms}^2 = \frac{3RT}{M}$$


$$\dot{\text{ }} M = \frac{3RT}{v_{rms}^2}$$

$$\Rightarrow M = \frac{3 \times 8.31 \times 300}{1365 \times 1365} \text{ kg}$$

$$\dot{\text{ }} \frac{3 \times 8.31 \times 300}{1365 \times 1365} \times 1000 \text{ g} = 4 \text{ g}$$

The molecular weight of helium is 4.

273 (c)



Draw two isothermals one passing through points 1 and 2 the other through mid point of straight line

joining 1 and 2

$T_2 > T_1$, at point 1 temperature is T_1 and that at mid point is T_2 and then at point 2 again it is T_1

\therefore The gas is first heated and then cooled towards end

274 (d)

Pressure due to an ideal gas is given by

$$p = \frac{M}{3V} v^2$$

Putting $\frac{M}{V} = \rho$, the density of gas

$$p = \frac{1}{3} \rho v^2$$

$$\Rightarrow v = \sqrt{\left(\frac{3p}{\rho}\right)}$$

$$\therefore v \propto \frac{1}{\sqrt{\rho}}$$

275 (b)

For first vessel, number of moles

$$n_1 = \frac{m_1}{M_1} = \frac{32}{32} = 1$$

Volume = V , Temperature = T

$$\therefore p_1 V = RT \quad \dots(i)$$

For second vessel number of moles

$$= n_2 = \frac{m_2}{M_2} = \frac{4}{2} = 2$$

Volume = V , Temperature = $2T$

$$\therefore p_2 V = 2R(2T) \quad \dots(ii)$$

From Eqs. (i) and (ii),

$$p_2 = 4 p_1 = 4 p$$

276 (b)

RMS speed of gas molecules does not depend on the pressure of gas (if temperature remains constant) because $p \propto \rho$. If pressure is increased n times density will also increase by n times but v_{rms} remains constant.

277 (d)

$$P = \frac{2}{3} \times (\text{Energy per unit volume})$$

$$\therefore \frac{2}{3} \frac{E}{V} \Rightarrow PV = \frac{2}{3} E$$

278 (b)

$C_p - C_v = R = \dot{i}$ Universal gas constant

279 (d)

$$V_{rms} = \sqrt{\frac{3RT}{M}}$$

$$\% \text{ increase in } V_{rms} = \frac{\sqrt{\frac{3RT_2}{M}} - \sqrt{\frac{3RT_1}{M}}}{\sqrt{\frac{3RT_1}{M}}} \times 100\%$$

$$\therefore \frac{20 - 17.32}{17.32} \times 100 = 15.5\%$$

280 (d)

Using $\gamma_r = \gamma_a + \dot{i} g$, we get

$$\gamma_r = \gamma_1 + 3\alpha = \gamma_2 + 3\beta$$

$$\therefore \beta = \frac{\gamma_1 - \gamma_2}{3} + \alpha$$

281 (a)

As the steel tape is calibrated at 10°C , therefore, adjacent centimeter marks on the steel tape will be separated by a distance of

$$l_t = l_{10}(1 + \alpha_s \Delta T) = (1 + \alpha_s 20) \text{ cm}$$

Length of copper rod at 30°C

$$= 90(1 + \alpha_c 20) \text{ cm}$$

Therefore, number of centimeters read on the tape will be

$$\begin{aligned} &= \frac{90(1 + \alpha_c 20)}{1(1 + \alpha_s 20)} = \frac{90(1 + 1.7 \times 10^{-5} \times 20)}{1(1 + 1.2 \times 10^{-5} \times 20)} \\ &= \frac{90 \times 1.00034}{1.00024} = 90.01 \text{ cm} \end{aligned}$$

282 (c)

$$\text{At absolute temperature } T = 0 \Rightarrow v_{rms} = \sqrt{\frac{3RT}{M}} = 0$$

Therefore, there is no motion of gas molecules at this temperature

283 (b)

Average kinetic energy \propto Temperature

284 (c)

A diatomic molecule has three translational and two rotational degrees of freedom

Hence total degrees of freedom $f = 3 + 2 = 5$

285 (c)

$$\gamma = 1 + \frac{2}{f} \Rightarrow 1.4 = 1 + \frac{2}{f} \Rightarrow \text{Degree of freedom } f = 5$$

\Rightarrow Degree of freedom of diatomic gas is 5 and it's

$$C_p = \frac{7}{2} R \text{ and } C_v = \frac{5}{2} R$$

287 (a)

Apparent weight (w_a) = Actual weight (w)

– upthrust (F), where upthrust = weight of water displaced = $V \rho_0 g$

$$\text{Now, } F_0 = V_0 \rho_0 g \text{ and } F_{50} = V_{50} \rho_{50} g$$

$$\therefore \frac{F_{50}}{F_0} = \frac{V_{50} \rho_{50} g}{V_0 \rho_0 g} = \frac{1 + \gamma_m \times 50}{1 + \gamma_w \times 50}$$

As $\gamma_m < \gamma_w$, therefore, $F_{50} < F_0$

Hence, $(w_a)_{50} (w_a)_0 \vee w_2 > w_1 \vee w_1 < w_2$

288 (c)

For intermolecular attraction is considered in real gas and for real gases pressure is given by

$$P = \frac{nRT}{V - nb} - \frac{n^2 a}{V^2}. \text{ Here } \left(\frac{n}{V}\right)^2 \text{ represents the}$$

reduction in pressure due to intermolecular attraction

289 (a)

$$PV = \mu RT \Rightarrow P \propto \frac{T}{V}. \text{ If } T \text{ and } V \text{ both doubled then}$$

pressure remains same,

$$i.e., P_2 = P_1 = 1 \text{ atm} = 1 \times 10^5 \text{ N/m}^2$$

290 (a)

$$V \propto T \text{ [as constant pressure]}$$

291 (d)

$$v_{rms} = \sqrt{\frac{3kT}{m}} = v_{rms} \propto \frac{1}{\sqrt{m}}$$

292 (d)

Specific heat for a monoatomic gas

$$C_V = \frac{3}{2} R$$

$$\therefore \text{Heat } dQ = \mu C_V \Delta T$$

$$dQ = \mu \times \frac{3}{2} \times R (473 - 273)$$

$$\therefore 4 \times \frac{3}{2} \times R \times 200 (\because \mu = 4)$$

$$\therefore dQ = 4 \times 300 R$$

$$\therefore 1200 R$$

293 (b)

Universal gas constant

$$R = C_p - C_V$$

294 (a)

22 g of CO_2 is half mole of CO_2 i.e., $n_1 = 0.5$

16 g of O_2 is half mole of O_2 i.e., $n_2 = 0.5$

$$\therefore T = \frac{n_1 T_1 + n_2 T_2}{n_1 + n_2}$$

$$= \frac{0.5 \times (27 + 273) + 0.5 (37 + 273)}{0.5 + 0.5}$$

$$= 305 \text{ K}$$

$$= 305 - 273 = 32^\circ \text{C}$$

295 (a)

$$PV = mRT = m \left(\frac{R}{M}\right) T$$

$$\Rightarrow V = \left(\frac{m}{M}\right) \frac{RT}{P} = \left(\frac{2.2}{44}\right) \times \frac{8.31 \times (273 + 0)}{2 \times (1 \times 10^5)}$$

$$\therefore 5.67 \times 10^{-4} \text{ m}^3 = 0.56 \text{ litre}$$

296 (c)

If number of molecules in gas increases then number of collisions of molecules with walls of container would also increase and hence the pressure increases, *i.e.*, $P \propto N$.

$$\Rightarrow \frac{P_2}{P_1} = \frac{N_2}{N_1} = \frac{2}{1} \Rightarrow P_2 = 2P_1$$

297 (a)

Pressure of the gas will not be affected by motion of the system, hence by

$$v_{rms} = \sqrt{\frac{3P}{\rho}} \Rightarrow \dot{c}^2 = \frac{3P}{\rho} \Rightarrow P = \frac{1}{3} \rho \dot{c}^2$$

298 (b)

As the temperature increases, the average velocity increases. So the collisions are faster

299 (d)

$$(\Delta Q)_p = \mu C_p \Delta T \Rightarrow 207 = 1 \times C_p \times 10$$

$$\Rightarrow C_p = 20.7 \frac{\text{Joule}}{\text{mol-K}}. \text{ Also } C_p - C_v = R$$

$$\Rightarrow C_v = C_p - R = 20.7 - 8.3 = 12.4 \frac{\text{Joule}}{\text{mole-K}}$$

$$\text{So, } (\Delta Q)_v = \mu C_v \Delta T = 1 \times 12.4 \times 10 = 124 \text{ J}$$

300 (a)

At constant pressure

$$V \propto T \Rightarrow \frac{V_2}{V_1} = \frac{T_2}{T_1} \Rightarrow T_2 = \left(\frac{V_2}{V_1} \right) T_1$$

$$\Rightarrow T_2 = \left(\frac{3V}{V} \right) \times 273 = 819 \text{ K} = 546^\circ \text{C}$$

301 (c)

According to Boyle's law $(P_1 V_1)_{\text{bottom}} = (P_2 V_2)_{\text{top}}$

$$(10+h) \times \frac{4}{3} \pi r_1^3 = 10 \times \frac{4}{3} \pi r_2^3 \text{ but } r_2 = 2r_1$$

$$\therefore (10+h)r_1^3 = 10 \times 8r_1^3 \Rightarrow 10+h = 80 \therefore h = 70 \text{ m}$$

302 (c)

Here temperature remain constant

$$\text{So } P_1 V_1 = P_2 V_2 = 76 \times 5 = P_2 \times 35$$

$$\Rightarrow P_2 = \frac{76 \times 5}{35} = 10.85 \text{ cm of Hg}$$

303 (b)

$$\text{For diatomic gases } \frac{C_p}{C_v} = \gamma = 1.4$$

304 (a)

$$\text{Using } \frac{C}{5} = \frac{F-32}{9}$$

$$\frac{-183}{5} = \frac{F-32}{9}$$

$$F-32 = \frac{-183 \times 9}{5} = -329.4$$

$$F = -329.4 + 32 = -297.4^\circ$$

307 (d)

$$n_1 C_v \Delta T_1 = n_2 C_v \Delta T_2$$

$$10 \times (T-10) = 20(20-T)$$

$$T-10 = 40-2T$$

$$3T = 50 \Rightarrow T = 16.6^\circ \text{C}$$

308 (b)

Number of translational degrees of freedom (3) are same for all types of gases

309 (a)

$$\frac{T_A}{M_A} = 4 \frac{T_B}{M_B} \Rightarrow \sqrt{\frac{T_A}{M_A}} = 2 \sqrt{\frac{T_B}{M_B}}$$

$$\Rightarrow \sqrt{\frac{3RT_A}{M_A}} = 2 \sqrt{\frac{3RT_B}{M_B}} \Rightarrow C_A = 2C_B \Rightarrow \frac{C_A}{C_B} = 2$$

310 (b)

Neon gas is monoatomic and for monoatomic gases

$$C_v = \frac{3}{2} R$$

311 (b)

Thermal capacity = Mass \times Specific heat

Due to same material both spheres will have same specific heat.

Also mass = Volume (V) \times Density (ρ)

\therefore Ratio of thermal capacity

$$\therefore \frac{m_1}{m_2} = \frac{V_1 \rho}{V_2 \rho} = \frac{\frac{4}{3} \pi r_1^3}{\frac{4}{3} \pi r_2^3} = \left(\frac{r_1}{r_2} \right)^3$$

$$= \left(\frac{1}{2} \right)^3 = \frac{1}{8}$$

312 (c)

C_p is always greater than C_v

$$\text{ie, } C_p > C_v$$

313 (a)

$$\text{As } \theta_2 > \theta_1 \Rightarrow \tan \theta_2 > \tan \theta_1 \Rightarrow \left(\frac{T}{P} \right)_2 > \left(\frac{T}{P} \right)_1$$

$$\text{Also from } PV = \mu RT; \frac{T}{P} \propto V \Rightarrow V_2 > V_1$$

314 (a)

According to kinetic theory, molecules of a liquid are in a state of continuous random motion. They continuously collide against the walls of the container. During each collision, certain momentum is transferred to the walls of the container. So, kinetic energy of molecules increases, hence due to random motion, the temperature increase. So, random motion of molecules and not ordered motion cause rise of

temperature.

315 (d)

From Maxwell's velocity distribution law, we infer that

$$v_{rms} > v > v_{mp}$$

ie, most probable velocity is less than the root mean square velocity.

316 (a)

Mayer Formula

317 (b)

Temperature remain constant so

$$v_{rms} \propto \frac{1}{\sqrt{M}} \Rightarrow \frac{v_{O_2}}{v_{H_2}} = \sqrt{\frac{M_{H_2}}{M_{O_2}}} = \sqrt{\frac{1}{16}} = \frac{1}{4}$$

318 (c)

Mean kinetic energy of gas molecule

$$E = \frac{f}{2} kT = \frac{f}{2} k(t + 273) = \left(\frac{f}{2} k\right)t + \frac{f}{2} \times 273k;$$

Comparing it with standard equation of straight line

$$y = mx + c. \text{ We get } m = \frac{f}{2}k \text{ and } c = \frac{f}{2}273k$$

So the graph between E and t will be straight line with positive intercept on E -axis and positive slope with t -axis

319 (b)

In isothermal changes, temperature remains constant

320 (a)

$$E = \frac{3}{2} RT \Rightarrow \frac{E'}{E} = \frac{T'}{T} = \frac{400}{300} = \frac{4}{3} = 1.33$$

321 (c)

When saturated vapour is compressed some of the vapour condenses but pressure does not change

322 (d)

10 g of ice at -10°C to ice at 0°C

$$Q_1 = cm, \Delta\theta = 0.5 \times 10 \times 10 = 50 \text{ cal}$$

10 g of ice 0°C to water at 0°C

$$Q_2 = mL = 10 \times 80 = 800 \text{ cal}$$

10 g of water at 0°C to water at 100°C

$$Q_3 = cm, \Delta\theta = 1 \times 10 \times 100 = 1000 \text{ cal}$$

10 g water at 100°C to steam at 100°C

$$Q_4 = mL = 10 \times 540 = 5400 \text{ cal}$$

$$\text{Total heat required, } Q + Q_1 + Q_2 + Q_3 + Q_4 \\ = 50 + 800 + 1000 + 5400 = 7250 \text{ cal}$$

323 (a)

When the piston is in equilibrium, the pressure is same on both the sides of the piston. It is given that temperature and weight of gas on the two sides of piston not change. From ideal gas equation,

$pV = nRT$, we have $V \propto$ mass of the gas.

$$\text{So, } \frac{V_1}{V_2} = \frac{m_1}{m_2} \Rightarrow \frac{V_1}{V_2} + 1 = \frac{m_1}{m_2} + 1$$

$$\text{Or } \frac{V_1 + V_2}{V_2} = \frac{m_1 + m_2}{m_2}$$

$$\text{Or } \frac{V_2}{V_1 + V_2} = \frac{m_2}{m_1 + m_2} = \frac{2m}{m + 2m} = \frac{2}{3}$$

324 (d)

$$PV = \mu RT \Rightarrow P \left(\frac{m}{\rho}\right) = \mu RT \Rightarrow \rho \propto \frac{P}{T}$$

Since T becomes four times and P becomes twice so

ρ becomes $\frac{1}{2}$ times

325 (d)

Kinetic energy is function of temperature

327 (d)

For an ideal gas keeping the temperature same throughout,

$$pV = \text{constant}$$

Hence, for a given mass, the graph between $pV \wedge V$ will be a straight line parallel to V -axis whatever may be the volume.

328 (b)

$$P = \frac{\mu RT}{V} = \frac{mRT}{MV} \left(\mu = \frac{m}{M}\right)$$

So, at constant volume pressure-versus temperature graph is a straight line passing through origin with

slope $\frac{mR}{MV}$. As the mass is doubled and volume is

halved slope becomes four times. Therefore, pressure versus temperature graph will be shown by the line B

329 (a)

In free expansion of Vander waal's gas, its temperature decreases

330 (b)

The mean kinetic energy for gas molecules

$$E = \frac{3}{2} kT \Rightarrow E \propto T$$

$$\text{So, } \frac{E_1}{E_2} = \frac{T_1}{T_2} \dots (i)$$

According to question both gases are at the same temperature T .

$$\text{So, } \frac{E_1}{E_2} = \frac{T}{T} = \frac{1}{1}$$

$$\Rightarrow E_1 : E_2 = 1 : 1$$

331 (a)

$$v_{rms} = \sqrt{\frac{3RT}{M}} \Rightarrow T \propto M \Rightarrow \frac{T_{He}}{T_H} = \frac{M_{He}}{M_H}$$

$$\Rightarrow \frac{(273+0)}{T_{He}} = \frac{2}{4} \Rightarrow T_{He} = 546 K = 273^\circ C$$

332 (b)

$$P_1 = 720 \text{ kPa}, T_1 = 40^\circ C = 273 + 40 = 313 K$$

$$P \propto mT \Rightarrow \frac{P_2}{P_1} = \frac{m_2 T_2}{m_1 T_1} = \frac{3}{4} \times \frac{626}{313} = 1.5$$

$$\Rightarrow P_2 = 1.5 P_1 = 1.5 \times 720 = 1080 \text{ kPa}$$

333 (c)

Since the volume of cylinder is fixed, the heat required is determined by C_V

He is a monoatomic gas.

Therefore, its molar specific heat at constant volume is

$$C_V = \frac{3}{2} R$$

\therefore Heat required = no. of moles \times molar specific \times rise in temperature

$$i \times \frac{3}{2} R \times 20 = 60 R = 60 \times 8.31 = 498.6 J$$

334 (d)

$$l = l_0 \left(1 + \frac{1}{100} \right)$$

$$\therefore 2l^2 = 2l_0^2 \left(1 + \frac{1}{100} \right)^2$$

$$\text{Or } 2l^2 - 2l_0^2 = 2l_0^2 \times \frac{2}{100}$$

$$\text{Or } \Delta S = S \times \frac{2}{100} \vee \frac{\Delta S}{S} = \frac{2}{100} = 2\%$$

335 (d)

$$\frac{(v_{rms})_1}{(v_{rms})_2} = \sqrt{\frac{T_1}{T_2}} \Rightarrow \frac{500}{(v_{rms})_2} = \sqrt{\frac{0+273}{819+273}} = \sqrt{\frac{273}{1092}}$$

$$(v_{rms})_2 = 500 \sqrt{\frac{1092}{273}} = 500 \sqrt{4} = 1000 \frac{m}{s} = 1 \frac{km}{s}$$

336 (a)

The value of universal gas constant is approx.

$$2 \frac{\text{cal}}{\text{mole-Kelvin}}$$

337 (a)

Let V be the volume of solid ; d be its density and m be its mass ; if g coefficient of volume expansion of liquid, then

$$\text{Density at temperature } t_1 \text{ is, } d_1 = \frac{d_0}{1 + \gamma t_1}$$

$$\text{Density at temperature } t_2 \text{ is, } d_2 = \frac{d_0}{1 + \gamma t_2}$$

According to Archimede's principle,

$$f_1 V d_1 = m = f_2 V d_2$$

$$\text{Or } \frac{d_1}{d_2} = \frac{f_2}{f_1} = \frac{d_0}{(1 + \gamma t_1)} \frac{(1 + \gamma t_2)}{d_0}$$

$$\text{Or } f_1 + f_1 \gamma t_2 = f_2 + f_2 \gamma t_1$$

$$f_1 - f_2 = \gamma (f_2 t_1 - f_1 t_2)$$

$$\gamma = \frac{(f_1 - f_2)}{f_2 t_1 - f_1 t_2}$$

338 (c)

$$\gamma_{poly} = \frac{(4 + f_{vib})}{(3 + f_{vib})}$$

$f_{vib} = i$ degree of freedom due to vibration

$$\Rightarrow \gamma_{poly} < \frac{4}{3}$$

Or $\gamma_{poly} < 1.33$

Also you can remember that as the atomicity of gas increases the value of γ -decreases

339 (a)

For NH_3 , degree of freedom $f = 6$

$$\Rightarrow \frac{C_P}{C_V} = \gamma = 1 + \frac{2}{f} = 1 + \frac{2}{6} = \frac{4}{3} = 1.33$$

340 (d)

$$\text{From } C_V = \frac{1}{2} fR = \frac{1}{2} \times 6R = 3R$$

341 (c)

$$\text{Mean kinetic energy per molecule } E = \frac{f}{2} kT = \frac{n}{2} kT$$

342 (b)

Mean free path $\lambda \propto \frac{1}{P}$; If λ is doubled then P

becomes half

343 (a)

Average kinetic theory of one molecule is

$$E = \frac{3}{2} kT$$

where k is Boltzmann constant and T the absolute temperature.

$$\text{Given, } T_1 = -68^\circ C = 273 - 68 = 205 K,$$

$$E_1 = E, E_2 = 2E$$

$$\therefore \frac{E_1}{E_2} = \frac{T_1}{T_2}$$

$$\Rightarrow T_2 = \frac{T_1 E_2}{E_1}$$

$$\therefore T_2 = \frac{205 \times 2E}{E} = 410 \text{ K}$$

344 (c)

$$C_V = \frac{R}{0.67} = 1.5R = \frac{3}{2}R$$

This is the value for monoatomic gases

345 (c)

$$C_{\text{isothermal}} = \infty \wedge C_{\text{adiabatic}} = 0$$

346 (b)

$$C_p - C_V = \frac{R}{J} \Rightarrow C_p = \frac{R}{J} + C_V = \frac{R}{J} + \frac{R}{J(\gamma-1)}$$

$$\Rightarrow C_p = \frac{R}{J} \left(\frac{\gamma}{\gamma-1} \right) = \frac{R}{J} \left(\frac{1.5}{1.5-1} \right) = \frac{3R}{J}$$

347 (d)

$$\text{For any gas } C_p - C_V = 1.99 = 2 \frac{\text{cal}}{\text{mol-K}}$$

349 (a)

$$\frac{v_2}{v_1} = \sqrt{\frac{T_2}{T_1}} \Rightarrow \frac{v_2}{400} = \sqrt{\frac{273+227}{273+27}} = \sqrt{\frac{5}{3}}$$

$$\Rightarrow v_2 = 400 \sqrt{5/3} = 516 \text{ m/s}$$

350 (c)

$$\text{Using pressure or Gay-Lussac's law } \frac{P_1}{P_2} = \frac{T_1}{T_2}$$

$$\text{or } P_2 = \frac{P_1 T_2}{T_1} = \frac{P(273+927)}{(273+27)} = 4P$$

351 (a)

$$\frac{C_p}{C_V} = \gamma = 1 + \frac{2}{f}$$

352 (d)

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \Rightarrow T_2 = \frac{P_2 V_2}{P_1 V_1} \times T_1$$

$$\therefore T_2 = \frac{1}{30} \times \frac{10}{1} \times 300 = 100 \text{ K} = -173^\circ \text{C}$$

353 (a)

$$v_{\text{average}} = \sqrt{\frac{8RT}{\pi M}} \Rightarrow v_{\text{av}} \propto \sqrt{T}$$

354 (a)

$$\Delta p = mV - (-mV) = 2mV$$

355 (b)

$$\text{Kinetic energy for } 1g \Rightarrow E_{\text{trans}} = \frac{3}{2}rT = \frac{3}{2} \frac{RT}{M}$$

356 (b)

$$C_p - C_V = R$$

At constant pressure, Heat $\dot{Q} = n C_p \theta$

$$\Rightarrow 310 = 2 \times C_p \times (35 - 25) = 20 C_p$$

$$\Rightarrow C_p = \frac{310}{20} = 15.5$$

At constant volume, Heat required $\dot{Q} = n C_V \theta$

$$\Rightarrow Q = 2 \times (C_p - R) \times (32 - 25)$$

$$\therefore 2 \times (15.5 - 8.3) \times 10 = 2 \times 7.2 \times 10 = 144 \text{ J}$$

357 (b)

The collision of molecules of ideal gas is elastic collision

358 (c)

Mean kinetic energy of molecule depends upon temperature only. For O_2 it is same as that of H_2 at the same temperature of -73°C

359 (a)

When C_p and C_V are given with *calorie* and R with *Joule* then $C_p - C_V = R/J$

360 (c)

$$C_V = \frac{n_1 C_{V1} + n_2 C_{V2}}{n_1 + n_2}$$

$$\therefore \frac{1 \times \frac{3}{2}R + 1 \times \frac{5}{2}R}{1+1} = 2R$$

361 (b)

Molar specific heat of the mixture at constant volume is

$$C_V = \frac{n_1 C_{V1} + n_2 C_{V2}}{(n_1 + n_2)} = \frac{2 \left(\frac{3}{2}R \right) + 3 \left(\frac{5}{2}R \right)}{2+3} = 2.1R$$

363 (d)

$$v_{\text{rms}} = \sqrt{\frac{3P}{\rho}} \Rightarrow P = \frac{v_{\text{rms}}^2 \rho}{3} = \frac{(3180)^2 \times 8.99 \times 10^{-2}}{3}$$

$$\therefore 3.03 \times 10^5 \text{ N/m}^2 = 3 \text{ atm}$$

364 (c)

Kinetic energy of ideal gas depends only on its temperature. Hence, it remains constant whether its pressure is increased or decreased.

365 (a)

$$V \propto T \Rightarrow \frac{V_1}{V_2} = \frac{T_1}{T_2} \Rightarrow \frac{200}{V_2} = \frac{(273+20)}{(273-20)} = \frac{293}{253}$$

$$\Rightarrow V_2 = \frac{200 \times 253}{293} = 172.6 \text{ ml}$$

366 (b)

$$PV = \mu RT = \frac{m}{M} RT \Rightarrow \frac{m}{VP} = \frac{\text{density}}{P} = \frac{M}{RT}$$

$$\left(\frac{\text{density}}{P} \right)_{\text{At } 0^\circ \text{C}} = \frac{M}{R(273)} = x \quad \dots \text{(i)}$$

$$\left(\frac{\text{density}}{P} \right)_{\text{At } 100^\circ \text{C}} = \frac{M}{R(373)} \quad \dots \text{(ii)}$$

$$\Rightarrow \left(\frac{\text{density}}{P} \right)_{\text{At } 100^\circ\text{C}} = \frac{273 \times}{373}$$

367 (c)

Here, $\Delta l = 80.3 - 80.0 = 0.3 \text{ cm}$

$$l = 80 \text{ cm}, \alpha = 12 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$$

$$\text{Rise in temperature } \Delta T = \frac{\Delta l}{l\alpha}$$

$$\Delta T = \frac{0.3}{80 \times 12 \times 10^{-6}} = 312.5 \text{ }^\circ\text{C}$$

369 (a)

$$\left(P + \frac{aT^2}{V} \right) V^c = (RT + b) \Rightarrow P = (RT + b) V^{-c} - (aT^2) V^{-c}$$

Comparing this equation with $P = AV^m - BV^n$

We get $m = -c$ and $n = -1$

370 (d)

$$\Delta Q = KA \left(\frac{\Delta T}{\Delta x} \right) \Delta t, \text{ where } A = 4\pi r^2$$

$$= 0.008 \times 4 \times \frac{22}{7} (6 \times 10^8)^2 \times \left(\frac{32}{10^5} \right) \times 86400$$

$$= 10^{18} \text{ cal}$$

371 (c)

$$v_{rms} \propto \sqrt{T} \Rightarrow \frac{v_1}{v_2} = \sqrt{\frac{T_1}{T_2}} = \sqrt{\frac{200}{800}} = \frac{1}{2} \Rightarrow v_2 = 2v_1$$

372 (c)

$$p = \frac{n_1 RT + n_2 RT + n_3 RT}{V}$$

$$= (n_1 + n_2 + n_3) \frac{RT}{V}$$

$$= \left(\frac{8}{16} + \frac{14}{28} + \frac{22}{44} \right) \times \frac{0.082 \times 300}{10} = 3.69 \text{ atm}$$

373 (c)

As number of moles increases, pressure increases and at certain pressure vapour condenses hence pressure now decreases

374 (d)

Root mean square velocity

$$v_{rms} = \sqrt{\frac{3RT}{M}}$$

where R is gas constant, T the temperature and M molecular weight.

Given,

$$M_{N_2} = 28, M_{O_2} = 32, T_{O_2} = 127^\circ\text{C} = 127 + 273 = 400 \text{ K}$$

K

$$\therefore \frac{v_{O_2}}{v_{N_2}} = \sqrt{\frac{T_{O_2}}{M_{O_2}} \times \frac{M_{N_2}}{T_{N_2}}} = \sqrt{\frac{400}{32} \times \frac{28}{T_{N_2}}} = 1$$

$$\Rightarrow T_{N_2} = 350 \text{ K} = 77^\circ\text{C}.$$

375 (b)

Temperature becomes $\frac{1}{4}th$ of initial value

$$[1200 \text{ K} = 927^\circ\text{C} \rightarrow 300 \text{ K} = 27^\circ\text{C}]$$

So, using $v_{rms} \propto \sqrt{T}$, r.m.s. velocity will be half of the initial value

376 (d)

$$v_{rms} \propto \frac{1}{\sqrt{M}}; \text{ so } \frac{(v_{rms})_{O_2}}{(v_{rms})_{H_2}} = \sqrt{\frac{M_{H_2}}{M_{O_2}}} = \sqrt{\frac{2}{32}} = 1:4$$

377 (b)

Number of moles $n = 5 \text{ mol}, T_1 = 100^\circ\text{C}$,

$T_2 = 120^\circ\text{C}, \Delta U = 80 \text{ J}$

Rise in temperature $\Delta t = 120 - 100 = 20^\circ\text{C}$

$$\Delta U = ms \Delta t$$

$$\frac{80}{5} = 1 \times s \times 20$$

$$s = 0.8 \text{ J}$$

\therefore For 5 mol, $s = 0.8 \times 5 \text{ J K}^{-1} = 4 \text{ J K}^{-1}$

378 (a)

Ratio of specific heat for a monoatomic gas is $\frac{5}{3}$

and for diatomic gas is $\frac{7}{5}$.

Given, $n_1 = 1, n_2 = 3, n = 4$

$$\therefore \frac{n}{\gamma - 1} = \frac{n_1}{\gamma_1 - 1} + \frac{n_2}{\gamma_2 - 1}$$

$$\frac{4}{\gamma - 1} = \frac{1}{\frac{5}{3} - 1} + \frac{3}{\frac{7}{5} - 1}$$

$$\Rightarrow \frac{4}{\gamma - 1} = \frac{3}{2} + \frac{15}{2} = 9$$

$$\therefore 4 = 9\gamma - 9$$

$$\Rightarrow 9\gamma = 13 \Rightarrow \gamma = \frac{13}{9}$$

$$\text{Now, } C_v(\gamma - 1) = R$$

$$\text{or } C_v = \frac{R}{\gamma - 1} = \frac{8.3}{\frac{13}{9} - 1} = \frac{8.3 \times 9}{4}$$

$$\Rightarrow C_v = 18.7 \text{ J mol}^{-1} \text{ K}^{-1}$$

379 (b)

Using the relation $p = \frac{1}{3} \frac{mnv^2}{V}$

...(i)

and also
$$p' = \frac{1}{3} \frac{\frac{m}{2} n (2v)^2}{V}$$

...(ii)

Dividing Eq.(ii) by Eq. (i), we get

$$\frac{p'}{p} = 2$$

So, $p:p' = 1:2$

The ratio of initial and final pressures is 1:2.

380 (c)

Molar specific heat at constant pressure $C_p = \frac{7}{2} R$

Since, $C_p - C_v = R \Rightarrow C_v = C_p - R = \frac{7}{2} R - R = \frac{5}{2} R$

$$\therefore \frac{C_p}{C_v} = \frac{(7/2)R}{(5/2)R} = \frac{7}{5}$$

381 (d)

According to the equilibrium theorem, the molar heat capacities should be independent of temperature.

However, variations in C_v and C_p are observed as the temperature changes. At very high temperatures, vibrations are also important and that affects the values of C_v and C_p for diatomic and polyatomic gases. Here in this question according to given information (d) may be correct answer

382 (d)

We know $v_s = \sqrt{\frac{\gamma P}{\rho}} \wedge v_{rms} = \sqrt{\frac{3P}{\rho}}$

$$\therefore \frac{v_{rms}}{v_s} = \sqrt{\frac{\gamma}{3}}$$

383 (a)

For 1 g gas $PV = rT = \left(\frac{R}{M}\right)T$

Since P and V are constant $\Rightarrow T \propto M \Rightarrow \frac{T_{N_2}}{T_{O_2}} = \frac{M_{N_2}}{M_{O_2}}$

$$\Rightarrow \frac{T_{N_2}}{(273+15)} = \frac{28}{32} \Rightarrow T_{N_2} = 252 K = -21^\circ C$$

384 (a)

$(\Delta Q)_v = C_v \Delta T = \frac{f}{2} R \Delta T$

$$\Rightarrow \Delta T \propto \frac{1}{f}$$

Also $f_{Mono} < f_{Dia} \Rightarrow (\Delta T)_{Mono} > (\Delta T)_{Dia}$

385 (b)

$$v_{rms} = \sqrt{\frac{3RT}{M}} = \sqrt{3} \sqrt{\frac{RT}{M}} = 1.73 \sqrt{\frac{RT}{M}}$$

386 (d)

$PV = kT \Rightarrow P \left(\frac{m}{\rho}\right) = kT \Rightarrow \rho = \frac{Pm}{kT}$

387 (c)

Below 100 K only translational degree of freedom is considered. Hence

$$\gamma_{mixture} = \frac{\frac{\mu_1 \gamma_1}{\gamma_1 - 1} + \frac{\mu_2 \gamma_2}{\gamma_2 - 1}}{\frac{\mu_1}{\gamma_1 - 1} + \frac{\mu_2}{\gamma_2 - 1}} \text{ according}$$

to question, $\mu_1 = \mu_2$ and $\gamma_1 = \gamma_2 = 1 + \frac{2}{3} = \frac{5}{3}$

$$\Rightarrow \gamma_{mix} = \gamma_1 = \frac{5}{3}$$