

3.MATRICES

Single Correct Answer Type

1. If n is a natural number. Then $\begin{bmatrix} 2 & -1 \\ 3 & -2 \end{bmatrix}^n$, is
 - a) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ if n is even
 - b) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ if n is odd
 - c) $\begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$ if n is a natural number
 - d) None of these
2. $\begin{bmatrix} 0 & a \\ b & 0 \end{bmatrix}^4 = I$, then
 - a) $a = 1 = 2b$
 - b) $a = b$
 - c) $a = b^2$
 - d) $ab = 1$
3. If a square matrix A is such that $AA^T = I = A^T A$, then $|A|$ is equal to
 - a) 0
 - b) ± 1
 - c) ± 2
 - d) None of these
4. If k is a scalar and I is a unit matrix of order 3, then $\text{adj}(kI)$ is equal to
 - a) $k^3 I$
 - b) $k^2 I$
 - c) $-k^3 I$
 - d) $-k^2 I$
5. If A is a singular matrix, then $A \text{adj}(A)$ is a
 - a) Scalar matrix
 - b) Zero matrix
 - c) Identity matrix
 - d) Orthogonal matrix
6. If $A = \begin{bmatrix} \cos^2 \alpha & \cos \alpha \sin \alpha \\ \cos \alpha \sin \alpha & \sin^2 \alpha \end{bmatrix}$ and $A = \begin{bmatrix} \cos^2 \beta & \cos \beta \sin \beta \\ \cos \beta \sin \beta & \sin^2 \beta \end{bmatrix}$ are two matrices such that the product AB is null matrix, then $\alpha - \beta$ is
 - a) 0
 - b) Multiple of π
 - c) An odd multiple of $\pi/2$
 - d) None of the above
7. Consider the following statements:
 1. If A and B are two square matrices of same order and commute, then $(A + B)(A - B) = A^2 - B^2$
 2. If A and B are two square matrices of same order, then $(AB)^n = A^n B^n$
 3. If A and B are two matrices such that $AB = A$ and $BA = B$, then A and B are idempotent
 Which of these is/are not correct?
 - a) Only (1)
 - b) (2) and (3)
 - c) (3) and (1)
 - d) All of these
8. If $A = \begin{bmatrix} 2 & 4 & 5 \\ 4 & 8 & 10 \\ -6 & -12 & -15 \end{bmatrix}$, the rank of A is equal to
 - a) 0
 - b) 1
 - c) 2
 - d) 3
9. If A and B are 3×3 matrices such that $AB + B$ and $BA = A$, then
 - a) $A^2 = A$ and $B^2 \neq B$
 - b) $A^2 \neq A$ and $B^2 = B$
 - c) $A^2 = A$ and $B^2 = B$
 - d) $A^2 \neq A$ and $B^2 \neq B$
10. The inverse of the matrix $\begin{bmatrix} 7 & -3 & -3 \\ -1 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$ is
 - a) $\begin{bmatrix} 1 & 1 & 1 \\ 3 & 4 & 3 \\ 3 & 3 & 4 \end{bmatrix}$
 - b) $\begin{bmatrix} 1 & 3 & 1 \\ 4 & 3 & 8 \\ 3 & 4 & 1 \end{bmatrix}$
 - c) $\begin{bmatrix} 1 & 1 & 1 \\ 3 & 3 & 4 \\ 3 & 4 & 3 \end{bmatrix}$
 - d) $\begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$
11. If A is a skew-symmetric matrix and n is a positive integer, then A^n is
 - a) A symmetric matrix
 - b) Skew-symmetric matrix
 - c) Diagonal matrix
 - d) None of these
12. The number of 3×3 non-singular matrices, with four entries as 1 and all other entries as 0, is
 - a) Less than 4
 - b) 5
 - c) 6
 - d) At least 7

- a) $K = -2$ b) $K = -1$ c) $K = 0$ d) $K = 1$
27. The system of equation,
 $x + y + z = 6$
 $x + 2y + 3z = 10$
And $x + 2y + \lambda z = \mu$
Has no solution, if
- a) $\lambda = 3, \mu = 10$ b) $\lambda \neq 3, \mu = 10$ c) $\lambda \neq 3, \mu \neq 10$ d) $\lambda = 3, \mu \neq 10$
28. If a square matrix A is such that $AA^T = I = A^T A$, then $|A|$ is equal to
- a) 0 b) ± 1 c) ± 2 d) None of these
29. Let $A = \begin{bmatrix} \cos^2 \theta & \sin \theta \cos \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$ and $B = \begin{bmatrix} \cos^2 \phi & \sin \phi \cos \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix}$, then $AB = O$, if
- a) $\theta = n\phi, n = 0, 1, 2, \dots$ b) $\theta + \phi = n\pi, n = 0, 1, 2, \dots$ c) $\theta = \phi + (2n + 1)\frac{\pi}{2}, n = 0, 1, 2, \dots$ d) $\theta = \phi + n\frac{\pi}{2}, n = 0, 1, 2, \dots$
30. If A, B are two square matrices such that $AB = A$ and $BA = B$, then
- a) A, B are idempotent b) Only A is idempotent c) Only B is idempotent d) None of these
31. If $\begin{bmatrix} x - y - z \\ -y + z \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 5 \\ 3 \end{bmatrix}$, then the values of x, y and z are respectively
- a) 5, 2, 2 b) 1, -2, 3 c) 0, -3, 3 d) 11, 8, 3
32. Inverse of the matrix $\begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$ is
- a) $\begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$ b) $\begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix}$ c) $\begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$ d) $\begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$
33. If $A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{bmatrix}$, $10B = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{bmatrix}$ and B is the inverse of A , then the value of α is
- a) 2 b) 0 c) 5 d) 4
34. $A = \begin{bmatrix} 0 & 3 & 3 \\ -3 & 0 & -4 \\ -3 & 4 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, then $B'(AB)$ is
- a) Null matrix b) Singular matrix c) Unit matrix d) Symmetric matrix
35. If $A = \begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}$, then A^{-1} is
- a) $\begin{bmatrix} 1/a & 0 & 0 \\ 0 & 1/b & 0 \\ 0 & 0 & 1/c \end{bmatrix}$ b) $\begin{bmatrix} -1/a & 0 & 0 \\ 0 & -1/b & 0 \\ 0 & 0 & -1/c \end{bmatrix}$ c) $\begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & 1/c \end{bmatrix}$ d) None of these
36. If $A = \begin{bmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{bmatrix}$, where $a^2 + b^2 = 1$, then $\text{adj}(A)$ is equal to
(Here, A^T is the transpose of A)
- a) A^{-1} b) A^T c) A d) $-A$
37. The rank of a null matrix is
- a) 0 b) 1 c) Does not exist d) None of these
38. If A and B are matrices of the same order, then $(A + B)^2 = A^2 + 2AB + B^2$ is possible, iff
- a) $AB = I$ b) $BA = I$ c) $AB = BA$ d) None of these

50. The characteristic roots of the matrix $\begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}$ are
- a) 1, 3, 6 b) 1, 2, 4 c) 4, 5, 6 d) 2, 4, 6
51. If matrix $A = \begin{bmatrix} 3 & 2 & 4 \\ 1 & 2 & -1 \\ 0 & 1 & 1 \end{bmatrix}$ and $A^{-1} = \frac{1}{k} \text{adj } A$, then k is
- a) 7 b) -7 c) $\frac{1}{7}$ d) 11
52. The number of 3×3 matrices A whose entries are either 0 or 1 and for which the system $A \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ has exactly two distinct solutions, is
- a) 0 b) $2^9 - 1$ c) 168 d) 2
53. If A is an invertible matrix of order n , then the determinant of $\text{adj } (A)$ is equal to
- a) $|A|^n$ b) $|A|^{n+1}$ c) $|A|^{n-1}$ d) $|A|^{n+2}$
54. $\text{adj} \begin{bmatrix} 1 & 0 & 2 \\ -1 & 1 & -2 \\ 0 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 5 & a & -2 \\ 1 & 1 & 0 \\ -2 & -2 & b \end{bmatrix}$, then $[a \ b]$ is equal to
- a) $[-4 \ 1]$ b) $[-4 \ -1]$ c) $[4 \ 1]$ d) $[4 \ -1]$
55. If $A = [x \ y \ z]$, $B = \begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix}$ and $C = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ Then, $B'(AB) = O$, if
- a) $[ax^2 + by^2 + cz^2 + 2gxy + 2fyz + 2czx] = 0$ b) $[ax^2 + cy^2 + bz^2 + xy + yz + zx] = 0$
c) $[ax^2 + by^2 + cz^2 + 2hxy + 2by + 2cz] = 0$ d) $[ax^2 + by^2 + cz^2 + 2gzx + 2hxy + 2fyz] = 0$
56. The multiplicative inverse of matrix $\begin{bmatrix} 2 & 1 \\ 7 & 4 \end{bmatrix}$ is
- a) $\begin{bmatrix} 4 & -1 \\ -7 & -2 \end{bmatrix}$ b) $\begin{bmatrix} -4 & -1 \\ 7 & -2 \end{bmatrix}$ c) $\begin{bmatrix} 4 & -7 \\ 7 & 2 \end{bmatrix}$ d) $\begin{bmatrix} 4 & -1 \\ -7 & 2 \end{bmatrix}$
57. The rank of matrix $\begin{bmatrix} 4 & 1 & 0 & 0 \\ 3 & 0 & 1 & 0 \\ 6 & 0 & 2 & 0 \end{bmatrix}$ is
- a) 4 b) 3 c) 2 d) 1
58. The solution of (x, y, z) the equation $\begin{bmatrix} 1 & 0 & 1 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ is (x, y, z)
- a) $(1, 1, 1)$ b) $(0, -1, 2)$ c) $(-1, 2, 2)$ d) $(-1, 0, 2)$
59. $\begin{bmatrix} 7 & 1 & 2 \\ 9 & 2 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix} + 2 \begin{bmatrix} 4 \\ 2 \end{bmatrix}$ is equal to
- a) $\begin{bmatrix} 43 \\ 44 \end{bmatrix}$ b) $\begin{bmatrix} 43 \\ 45 \end{bmatrix}$ c) $\begin{bmatrix} 45 \\ 44 \end{bmatrix}$ d) $\begin{bmatrix} 44 \\ 45 \end{bmatrix}$
60. Let A, B and C be $n \times n$ matrices. Which one of the following is a correct statements?
- a) If $AB = AC$, then $B = C$ b) If $A^3 + 2A^2 + 3A + 5I = O$; then A is invertible
c) If $A^2 = O$, then $A = O$ d) None of the above
61. If A and B are square matrices of order 3 such that $|A| = -1, |B| = 3$, then $|3 AB|$ equals
- a) -9 b) -81 c) -27 d) 81
62. The rank of the matrix $\begin{bmatrix} 1 & 2 & 3 & 0 \\ 2 & 4 & 3 & 2 \\ 3 & 2 & 1 & 3 \\ 6 & 8 & 7 & 5 \end{bmatrix}$ is
- a) 1 b) 2 c) 3 d) 4
63. If $A + I = \begin{bmatrix} 3 & -2 \\ 4 & 1 \end{bmatrix}$, then $(A - I)$ is equal to
- a) $\begin{bmatrix} -5 & -4 \\ 8 & -9 \end{bmatrix}$ b) $\begin{bmatrix} -5 & 4 \\ -8 & 9 \end{bmatrix}$ c) $\begin{bmatrix} 5 & 4 \\ 8 & 9 \end{bmatrix}$ d) $\begin{bmatrix} -5 & -4 \\ -8 & -9 \end{bmatrix}$

64. If $A = [a_{ij}]$ is a scalar matrix of order $n \times n$ such that $a_{ii} = k$ for all i , then $|A| =$
a) nk b) $n + k$ c) n^k d) k^n
65. If $A \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is such that $|A|=0$ and $A^2 - (a - d)A + kI = 0$, then k is equal to
a) $b + c$ b) $a + d$ c) $ab + cd$ d) Zero
66. If $A = \begin{bmatrix} 1 & -2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 \\ 3 & 2 \\ 1 & 1 \end{bmatrix}$, then $(AB)^T$ is equal to
a) $\begin{bmatrix} -3 & -2 \\ 10 & 7 \end{bmatrix}$ b) $\begin{bmatrix} -3 & 10 \\ -2 & 7 \end{bmatrix}$ c) $\begin{bmatrix} -3 & 7 \\ 10 & 2 \end{bmatrix}$ d) None of these
67. If A is a matrix such that there exists a square submatrix of order r which is non-singular and every square submatrix of order $r + 1$ or more is singular, then
a) $\text{rank}(A) = r + 1$ b) $\text{rank}(A) = r$ c) $\text{rank}(A) > r$ d) $\text{rank}(A) \geq r + 1$
68. If $A = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$, then $A^2 - 5A =$
a) I b) $14I$ c) 0 d) None of these
69. The order of $\begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ is
a) 3×1 b) 1×1 c) 1×3 d) 3×3
70. If $A + B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ and $A - 2B = \begin{bmatrix} -1 & 1 \\ 0 & -1 \end{bmatrix}$, then $A =$
a) $\begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix}$ b) $\begin{bmatrix} 2/3 & 1/3 \\ 1/3 & 2/3 \end{bmatrix}$ c) $\begin{bmatrix} 1/3 & 1/3 \\ 2/3 & 1/3 \end{bmatrix}$ d) None of these
71. If A is singular matrix, then $A \text{adj}(A)$
a) Is a scalar matrix b) Is a zero matrix
c) Is an identity matrix d) Is an orthogonal matrix
72. If $\begin{bmatrix} 3 & 1 \\ 4 & 1 \end{bmatrix} X = \begin{bmatrix} 5 & -1 \\ 2 & 3 \end{bmatrix}$, then X is equal to
a) $\begin{bmatrix} -3 & 4 \\ 14 & -13 \end{bmatrix}$ b) $\begin{bmatrix} 3 & -4 \\ -14 & 13 \end{bmatrix}$ c) $\begin{bmatrix} 3 & 4 \\ 14 & 13 \end{bmatrix}$ d) $\begin{bmatrix} -3 & 4 \\ -14 & 13 \end{bmatrix}$
73. For the matrix $A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 2 & 1 & 0 \end{bmatrix}$, which of the following is correct?
a) $A^3 + 3A^2 - I = 0$ b) $A^3 - 3A^2 - I = 0$ c) $A^3 + 2A^2 - I = 0$ d) $A^3 - A^2 + I = 0$
74. The matrices $P = \begin{bmatrix} u_1 & v_1 & w_1 \\ u_2 & v_2 & w_2 \\ u_3 & v_3 & w_3 \end{bmatrix}; Q = \frac{1}{9} \begin{bmatrix} 2 & 2 & 1 \\ 12 & -5 & m \\ -8 & 1 & 5 \end{bmatrix}$ are such that $PQ = I$, an identity matrix. Solving the equation $\begin{bmatrix} u_1 & v_1 & w_1 \\ u_2 & v_2 & w_2 \\ u_3 & v_3 & w_3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 5 \end{bmatrix}$, the value of y comes out to be -3 . Then, the value of m is equal to
a) 27 b) 7 c) -27 d) -7
75. If A is an invertible matrix, then which of the following is correct
a) A^{-1} is multivalued b) A^{-1} is singular c) $(A^{-1})^T \neq (A^T)^{-1}$ d) $|A| \neq 0$
76. If $\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} A \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then the matrix A equals
a) $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ b) $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ c) $\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ d) $\begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$
77. If A is any square matrix, then $\det(A - A^T)^T$ is equal to
a) 0 b) 1
c) Can be 0 or a perfect square d) Cannot be determined
78. If $O(A) = 2 \times 3, O(B) = 3 \times 2$, and $O(C) = 3 \times 3$, Which one of the following is not defined?
a) $CB + A'$ b) BAC c) $C(A + B)'$ d) $C(A + B')$
79. Suppose A is a matrix of order 3 and $B = |A|A^{-1}$. If $|A| = -5$, then $|B|$ is equal to
a) 1 b) -5 c) -1 d) 25

80. $A = \begin{bmatrix} -2 & 4 \\ -1 & 2 \end{bmatrix}$, the A^2 is equal to
 a) Null matrix b) Unit matrix c) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ d) $\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$
81. The system of simultaneous equations $kx + 2y - z = 1$, $(k - 1)y - 2z = 2$ and $(k + 2)z = 3$ has a unique solution, if k equals
 a) -2 b) -1 c) 0 d) 1
82. If $A = \begin{bmatrix} -1 & 2 \\ 2 & -1 \end{bmatrix}$ and $B = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$, $Ax = B$, then X is equal to
 a) $[0 \ 7]$ b) $\frac{1}{3} \begin{bmatrix} 5 \\ 7 \end{bmatrix}$ c) $\frac{1}{3} [5 \ 7]$ d) $\begin{bmatrix} 5 \\ 7 \end{bmatrix}$
83. For the equations $x + 2y + 3z = 1$, $2x + y + 3z = 2$ and $5x + 5y + 9z = 4$
 a) There is only one solution b) There exists infinitely many solutions
 c) There is no solution d) None of the above
84. If $A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ and $n \in N$, then A^n is equal to
 a) $2^n A$ b) $2^{n-1} A$ c) $n A$ d) None of these
85. If $A = \begin{bmatrix} 2 & 2 \\ -3 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$, then $(B^{-1}A^{-1})^{-1} =$
 a) $\begin{bmatrix} 2 & -2 \\ 2 & 3 \end{bmatrix}$ b) $\begin{bmatrix} 3 & -2 \\ 2 & 2 \end{bmatrix}$ c) $\frac{1}{10} \begin{bmatrix} 2 & 2 \\ -2 & 3 \end{bmatrix}$ d) $\frac{1}{10} \begin{bmatrix} 3 & 2 \\ -2 & 2 \end{bmatrix}$
86. If A is a skew symmetric matrix of order n and C is a column matrix of order $n \times 1$, then $C^T A C$ is
 a) An identity matrix of order n b) An identity matrix of order 1
 c) A zero matrix of order 1 d) None of the above
87. If A is a square matrix, then $A + A^T$ is
 a) Non-singular matrix b) Symmetric matrix
 c) Skew-symmetric matrix d) Unit matrix
88. If $A = [a_{ij}]_{2 \times 2}$, where $a_{ij} = i + j$, then A is equal to
 a) $\begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ b) $\begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix}$ c) $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ d) $\begin{bmatrix} 2 & 3 \\ 3 & 4 \end{bmatrix}$
89. If $P = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{bmatrix} \begin{bmatrix} -1 & -2 \\ -2 & 0 \\ 0 & -4 \end{bmatrix} \begin{bmatrix} -4 & -5 & -6 \\ 0 & 0 & 1 \end{bmatrix}$, then P_{22} is equal to
 a) 40 b) -40 c) -20 d) 20
90. Which of the following is/are incorrect?
 (i) adjoint of a symmetric matrix is symmetric
 (ii) adjoint of a unit matrix is a unit matrix
 (iii) $A(\text{adj } A) = (\text{adj } A)A = |A| I$
 (iv) adjoint of a diagonal matrix is a diagonal matrix
 a) (i) b) (ii) c) (iii) and (iv) d) None of these
91. If A and B are 3×3 matrices such that $AB = A$ and $BA = B$, then
 a) $A^2 = A$ and $B^2 \neq B$ b) $A^2 \neq A$ and $B^2 = B$
 c) $A^2 = A$ and $B^2 = B$ d) $A^2 \neq A$ and $B^2 \neq B$
92. Let A be a skew-symmetric matrix of even order, then $|A|$
 a) Is a square b) Is not a square c) Is always zero d) None of these
93. If $A = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 2 & -3 \\ 2 & 1 & 0 \end{bmatrix}$, $B = (\text{adj } A)$ and $C = 5A$, then $\frac{|\text{adj } B|}{|C|}$ is equal to
 a) 5 b) 25 c) -1 d) 1
94. If $A = \begin{bmatrix} 6 & 8 & 5 \\ 4 & 2 & 3 \\ 9 & 7 & 1 \end{bmatrix}$ is the sum of a symmetric matrix B and skew-symmetric matrix, C , then B is
 a) $\begin{bmatrix} 6 & 6 & 7 \\ 6 & 2 & 5 \\ 7 & 5 & 1 \end{bmatrix}$ b) $\begin{bmatrix} 0 & 2 & -2 \\ -2 & 5 & -2 \\ 2 & 2 & 0 \end{bmatrix}$ c) $\begin{bmatrix} 6 & 6 & 7 \\ -6 & 2 & -5 \\ -7 & 5 & 1 \end{bmatrix}$ d) $\begin{bmatrix} 0 & 6 & -2 \\ 2 & 0 & -2 \\ -2 & -2 & 0 \end{bmatrix}$

109. If $2X - \begin{bmatrix} 1 & 2 \\ 7 & 4 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 0 & -2 \end{bmatrix}$, then X is equal to
 a) $\begin{bmatrix} 2 & 2 \\ 7 & 4 \end{bmatrix}$ b) $\begin{bmatrix} 1 & 2 \\ 7 & 2 \end{bmatrix}$ c) $\begin{bmatrix} 2 & 2 \\ 7 & 1 \end{bmatrix}$ d) None of these
110. Let $A = \begin{bmatrix} 1 & 2 \\ -5 & 1 \end{bmatrix}$ and $A^{-1} = xA + yI$, then the values of x and y are
 a) $x = -\frac{1}{11}, y = \frac{2}{11}$ b) $x = -\frac{1}{11}, y = -\frac{2}{11}$ c) $x = \frac{1}{11}, y = \frac{2}{11}$ d) $x = \frac{1}{11}, y = -\frac{2}{11}$
111. Let A and B be two symmetric matrices of same order. Then, the matrix $AB - BA$ is
 a) A symmetric matrix b) A skew-symmetric matrix
 c) A null matrix d) The identity matrix
112. If $A = \begin{bmatrix} 1 & x \\ x^2 & 4y \end{bmatrix}$, $B = \begin{bmatrix} -3 & 1 \\ 1 & 0 \end{bmatrix}$ and $\text{adj } A + B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then the values of x and y are respectively
 a) (1, 1) b) (-1, 1) c) (1, 0) d) None of these
113. Let p is a non-singular matrix such that $1 + p + p^2 + \dots + p^n = O$ (O denotes the null matrix), then p^{-1} is
 a) p^n b) $-p^n$ c) $-(1 + p + \dots + p^n)$ d) None of these
114. If $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{40} \begin{bmatrix} 5 & 10 & -5 \\ -5 & -2 & 13 \\ 10 & -4 & 6 \end{bmatrix} \begin{bmatrix} 5 \\ 0 \\ 5 \end{bmatrix}$, then the value of $x + y + z$ is
 a) 3 b) 0 c) 2 d) 1
115. The matrix $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ is the matrix reflection in the line
 a) $x = 1$ b) $x + y = 1$ c) $y = 1$ d) $x = y$
116. If $\begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} \begin{bmatrix} 1 & \tan \theta \\ -\tan \theta & 1 \end{bmatrix}^{-1} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$, then
 a) $a = 1, b = 1$ b) $a = \sin 2\theta, b = \cos 2\theta$
 c) $a = \cos 2\theta, b = \sin 2\theta$ d) None of the above
117. If $A = \begin{bmatrix} -1 & -2 & -2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix}$, then $\text{adj } A$ is equal to
 a) A b) A' c) $3A$ d) $3A'$
118. Let the homogeneous system of linear equations $px + y + z = 0, x + qy + z = 0$, and $x + y + rz = 0$, where $p, q, r \neq 1$, have a non-zero solution, then the value of $\frac{1}{1-p} + \frac{1}{1-q} + \frac{1}{1-r}$ is
 a) -1 b) 0 c) 2 d) 1
119. If $A = \begin{bmatrix} 1 & \tan \frac{\theta}{2} \\ -\tan \frac{\theta}{2} & 1 \end{bmatrix}$ and $AB = I$, then B is equal to
 a) $\cos^2 \frac{\theta}{2} \cdot A$ b) $\cos^2 \frac{\theta}{2} \cdot A^T$ c) $\cos^2 \theta \cdot I$ d) $\sin^2 \frac{\theta}{2} \cdot A$
120. The values of x, y, z in order, if the system of equations $3x + y + 2z = 3, 2x - 3y - z = -3, x + 2y + z = 4$ has unique solution, are
 a) 2, 1, 5 b) 1, 1, 1 c) 1, -2, -1 d) 1, 2, -1
121. Matrix A is such that $A^2 = 2A - I$, where I is the identity matrix, then for $n \geq 2, A^n$ is equal to
 a) $nA - (n-1)I$ b) $nA - I$ c) $2^{n-1}A - (n-1)I$ d) $2^{n-1}A - I$
122. Matrix M_r is defined as $M_r = \begin{bmatrix} r & r-1 \\ r-1 & r \end{bmatrix}, r \in N$ value of $\det(M_1) + \det(M_2) + \det(M_3) + \dots + \det(M_{2007})$ is
 a) 2007 b) 2008 c) 2008^2 d) 2007^2
123. The number of solutions of the system of equations $x_2 - x_3 = 1, -x_1 + 2x_3 = -2, x_1 - 2, x_1 - 2x_2 = 3$ is
 a) Zero b) One c) Two d) Infinite
124. If $A = [a_{ij}]$ is a scalar matrix of order $n \times n$ such that $a_{ii} = k$ for all i , then trace of A is equal to
 a) nk b) $n + k$ c) n/k d) None of these
125. If $D = \text{diag}[d_1, d_2, d_3, \dots, d_n]$, where $d_i \neq 0 \forall i = 1, 2, \dots, n$ then D^{-1} is equal to

- b) $|A| = 0$ and $|B| = 0$
 c) Either $|A| = 0$ or $|B| = 0$
 d) $A = 0$ or $B = 0$

140. Let $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $D = \begin{bmatrix} 3 \\ 5 \\ 11 \end{bmatrix}$ and $A = \begin{bmatrix} 1 & -1 & -2 \\ 2 & 1 & 1 \\ 4 & -1 & -2 \end{bmatrix}$, if $X = A^{-1}D$, then X is equal to

- a) $\begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$ b) $\begin{bmatrix} \frac{8}{3} \\ -1 \\ \frac{3}{0} \end{bmatrix}$ c) $\begin{bmatrix} -\frac{8}{3} \\ 1 \\ 0 \end{bmatrix}$ d) $\begin{bmatrix} \frac{8}{3} \\ 1 \\ \frac{3}{-1} \end{bmatrix}$

141. If A and B are matrices such that AB and $A + B$ both are defined, then

- a) A and B can be any two matrices
 b) A and B are square matrices not necessarily of the same order
 c) A, B are square matrices of the same order
 d) Number of columns of A is same as the number of rows of B

142. Let a, b, c be any real numbers. Suppose that there are real numbers x, y, z not all zero such that $x = cy + bz, y = az + cx$, and $z = bx + ay$ have non-zero solution. Then, $a^2 + b^2 + c^2 + 2abc$ is equal to

- a) 1 b) 2 c) -1 d) 0

143. If I_n is the identity matrix of order n , then rank of I_n is

- a) 1 b) n c) 0 d) None of these

144. If the matrix $A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & \lambda \end{bmatrix}$ is singular, then λ is equal to

- a) 3 b) 4 c) 2 d) 5

145. If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, then $I + A + A^2 + A^3 + \dots \infty$ equals to

- a) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ b) $\begin{bmatrix} -1 & -2 \\ -3 & -4 \end{bmatrix}$ c) $\begin{bmatrix} 1/2 & -1/3 \\ -1/2 & 0 \end{bmatrix}$ d) $\begin{bmatrix} -1/4 & 1/3 \\ 1/2 & 0 \end{bmatrix}$

146. If A is a non-singular square matrix of order n , then the rank of A is

- a) Equal to n b) Less than n c) Greater than n d) None of these

147. If $A = \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix}$ and $f(t) = t^2 - 3t + 7$, then $f(A) + \begin{bmatrix} 3 & 6 \\ -12 & -9 \end{bmatrix}$ is equal to

- a) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ b) $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ c) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ d) $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$

148. The system of linear equations

$$x + y + z = 2$$

$$2x + y - z = 3$$

$$3x + 2y + kz = 4$$

has a unique solution if

- a) $k \neq 0$ b) $-1 < k < 1$ c) $-2 < k < 2$ d) $k = 0$

149. The number of solutions of the system of equations

$$2x + y - z = 7, x - 3y + 2z = 1, x + 4y - 3z = 5$$

- a) 0 b) 1 c) 2 d) 3

150. If $X = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, the value of X^n is equal to

- a) $\begin{bmatrix} 3^n & -4^n \\ n & -n \end{bmatrix}$ b) $\begin{bmatrix} 2+n & 5-n \\ n & -n \end{bmatrix}$ c) $\begin{bmatrix} 3^n & (-4)^n \\ 1^n & (-1)^n \end{bmatrix}$ d) None of these

151. If I_3 is the identity matrix of order 3, then $(I_3)^{-1} =$

- a) 0 b) $3I_3$ c) I_3 d) Not necessarily exists

152. If $A = [a_{ij}]$ is a square matrix of order $n \times n$ and k is a scalar, then $|kA| =$

- a) $k^n|A|$ b) $k|A|$ c) $k^{n-1}|A|$ d) None of these

153. If $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & -1 \end{bmatrix}$, then A^2 is equal to

154. a) Null matrix b) Unit matrix c) $-A$ d) A
 If $A = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix}$, then value of α for which $A^2 = B$ is
155. a) 1 b) -1 c) 4 d) No real values
 If A is a square matrix such that $A(\text{adj } A) = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix}$, then $|\text{adj } A| =$
156. a) $\omega^2 A$ b) ωA c) A d) 0
 If ω is a complex cube root of unity and $A = \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix}$, then A^{50} is
157. a) 0 b) -1 c) 2 d) None of these
 If $A = \begin{bmatrix} 1 & 2 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -2 & y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $AB = I_3$, then $x + y$ equals
158. a) $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ b) $\begin{bmatrix} \sin \theta & \cos \theta \\ \cos \theta & \sin \theta \end{bmatrix}$ c) $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ d) $\begin{bmatrix} -\sin \theta & \cos \theta \\ \cos \theta & \sin \theta \end{bmatrix}$
 The adjoint of the matrix $\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ is
159. a) $\begin{bmatrix} \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \\ -4 & 3 & -1 \\ \frac{5}{2} & \frac{3}{2} & \frac{1}{2} \end{bmatrix}$ b) $\begin{bmatrix} \frac{1}{2} & -4 & \frac{5}{2} \\ 1 & -6 & 3 \\ 1 & 2 & -1 \end{bmatrix}$ c) $\frac{1}{2} \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \\ 4 & 2 & 3 \end{bmatrix}$ d) $\frac{1}{2} \begin{bmatrix} 1 & -1 & -1 \\ -8 & 6 & -2 \\ 5 & -3 & 1 \end{bmatrix}$
 The inverse matrix of $A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$ is
160. a) $f(-\theta)$ b) $f(\theta)^{-1}$ c) $f(2\theta)$ d) None of these
 If $f(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$, then $\{f(\theta)^{-1}\}$ is equal to
161. a) $2ac$ b) $-ac$ c) ac d) $-2ac$
 If the three linear equations
 $x + 4ay + az = 0$
 $x + 3by + bz = 0$
 $x + 2cy + cz = 0$
 Have a non-trivial solution, where $a \neq 0, b \neq 0, c \neq 0$, then $ab + bc$ is equal to
162. a) $\text{rank}(AB) = mn$
 b) $\text{rank}(AB) \geq \text{rank}(A)$
 c) $\text{rank}(AB) \geq \text{rank}(B)$
 d) $\text{rank}(AB) \leq \min(\text{rank } A, \text{rank } B)$
 If A and B are two matrices such that $\text{rank of } A = m$ and $\text{rank of } B = n$, then
163. a) 1 b) 2 c) 3 d) 4
 If A is a non-zero column matrix of order $m \times 1$ and B is a non-zero row matrix of order $1 \times n$, then $\text{rank of } AB$ equals
164. a) $-\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ b) $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ c) $\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ d) $\begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$
 If $\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} A \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then A is equal to
165. a) $I - A$ b) $A - I$ c) A d) $A + I$
 If $A^2 - A + I = 0$, then the inverse of A is
166. a) $\text{rank}(BA) = \text{rank}(A)$ b) $\text{rank}(BA) \geq \text{rank}(B)$ c) $\text{rank}(BA) > \text{rank}(A)$ d) $\text{rank}(BA) > \text{rank}(B)$
 If B is an invertible matrix and A is a matrix, then
167. a) 6 b) 16 c) 10 d) None of these
 If $A = \begin{bmatrix} 4 & 2 \\ 3 & 4 \end{bmatrix}$, $|\text{adj } A|$ is equal to

168. $\cos \theta \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} + \sin \theta \begin{bmatrix} \sin \theta & -\cos \theta \\ \cos \theta & \sin \theta \end{bmatrix}$ is equal to
a) $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ b) $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ c) $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ d) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
169. Let $A = [a_{ij}]_{m \times n}$ be a matrix such that $a_{ij} = 1$ for all i, j . Then,
a) $\text{rank}(A^T) > 1$ b) $\text{rank}(A) = 1$ c) $\text{rank}(A) = m$ d) $\text{rank}(A) = n$
170. Let A be a square matrix all of whose entries are integers. Then, which one of the following is true?
a) If $\det(A) = \pm 1$, then A^{-1} need not exist
b) If $\det(A) = \pm 1$, then A^{-1} exists but all its entries are not necessarily integers
c) If $\det(A) \neq \pm 1$, then A^{-1} exists and all its entries are non – integers
d) If $\det(A) = \pm 1$, then A^{-1} exists and all its entries are integers
171. Matrix $A = \begin{bmatrix} 1 & 0 & -k \\ 2 & 1 & 3 \\ k & 0 & 1 \end{bmatrix}$ is invertible for
a) $k = 1$ b) $k = -1$ c) $k = \pm 1$ d) None of these
172. If $\begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} \begin{bmatrix} 1 & \tan \theta \\ -\tan \theta & 1 \end{bmatrix}^{-1} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$, then
a) $a = 1, b = 1$
b) $a = \cos 2\theta, b = \sin 2\theta$
c) $a = \sin 2\theta, b = \cos 2\theta$
d) None of these
173. If $x^2 + y^2 + z^2 \neq 0, x = cy + bz, y = az + cx$ and $z = bx + ay$, then $a^2 + b^2 + c^2 + 2abc =$
a) 2 b) $a + b + c$ c) 1 d) $ab + bc + ca$
174. If $A = \begin{bmatrix} 1 & -3 \\ 2 & k \end{bmatrix}$ and $A^2 - 4A = 10I = A$ then k is equal to
a) 0 b) -4 c) 4 and not 1 d) 1 or 4
175. Matrix A such that $A^2 = 2A - I$, where I is the identity matrix. Then, for $n \geq 2, A^n$ is equal to
a) $nA - (n - 1)I$ b) $nA - I$ c) $2^{n-1}A - (n - 1)I$ d) $2^{n-1}A - I$
176. The matrix A satisfying the equation $\begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} A = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$ is
a) $\begin{bmatrix} 1 & 4 \\ -1 & 0 \end{bmatrix}$ b) $\begin{bmatrix} 1 & -4 \\ 1 & 0 \end{bmatrix}$ c) $\begin{bmatrix} 1 & 4 \\ 0 & -1 \end{bmatrix}$ d) None of these
177. If A is an orthogonal matrix, then A^{-1} equals
a) A b) A^T c) A^2 d) None of these
178. By elementary transformation method, the inverse of $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 6 \end{bmatrix}$ is
a) $\begin{bmatrix} -2 & 0 & 1 \\ 0 & 3 & -2 \\ 1 & -2 & 1 \end{bmatrix}$ b) $\begin{bmatrix} 2 & 0 & -1 \\ 0 & -3 & 2 \\ -1 & 2 & -1 \end{bmatrix}$ c) $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 6 \end{bmatrix}$ d) None of these
179. What must be the matrix X , if $2X + \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 3 & 8 \\ 7 & 2 \end{bmatrix}$?
a) $\begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix}$ b) $\begin{bmatrix} 1 & -3 \\ 2 & -1 \end{bmatrix}$ c) $\begin{bmatrix} 2 & 6 \\ 4 & -2 \end{bmatrix}$ d) $\begin{bmatrix} 2 & -6 \\ 4 & -2 \end{bmatrix}$
180. If $A = \begin{bmatrix} 4 & x + 2 \\ 2x - 3 & x + 1 \end{bmatrix}$ is symmetric, then $x =$
a) 3 b) 5 c) 2 d) 4
181. If $A = \begin{bmatrix} 3 & 3 & 3 \\ 3 & 3 & 3 \\ 3 & 3 & 3 \end{bmatrix}$, A^4 is equal to
a) $27A$ b) $81A$ c) $243A$ d) $729A$
182. If ω is a complex cube root of unity, then the matrix $A = \begin{bmatrix} 1 & \omega^2 & \omega \\ \omega^2 & \omega & 1 \\ \omega & 1 & \omega^2 \end{bmatrix}$ is a
a) Singular matrix b) Non-symmetric matrix

$$a) \begin{bmatrix} 2 & 2 & -4 \\ 2 & 3 & 4 \\ -4 & 4 & 2 \end{bmatrix}$$

$$b) \begin{bmatrix} 2 & 4 & -5 \\ 0 & 3 & 7 \\ -3 & 1 & 2 \end{bmatrix}$$

$$c) \begin{bmatrix} 4 & 4 & -8 \\ 4 & 6 & 8 \\ -8 & 8 & 4 \end{bmatrix}$$

$$d) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

211. If the system of linear equations $x + 2ay + az = 0$, $x + 3by + bz = 0$ and $x + 4cy + cz = 0$ has a non-zero solution, then a, b, c

a) Are in AP

b) Are in GP

c) Are in HP

d) Satisfy $a + 2b + 3c = 0$

212. For what value of k the following system of linear equations will have infinite solutions

$$x - y + z = 3, 2x + y - z = 2$$

$$\text{and } -3x + 2ky + 6z = 3$$

a) $k \neq 2$

b) $k = 0$

c) $k = 3$

d) $k \in [2, 3]$

213. The product of two orthogonal matrices is

a) Orthogonal

b) Involutory

c) Unitary

d) Idempotent

214. The system of equations $x + y + z = 8$, $x - y + 2z = 6$, $3x + 5y - 7z = 14$ has

a) No solution

b) Unique solution

c) Infinitely many solution

d) None of the above

215. If the system of equations $x + ay = 0$, $az + y = 0$ and $ax + z = 0$ has infinite solutions, then the value of a is

a) -1

b) 1

c) 0

d) No real values

$$216. \begin{bmatrix} -6 & 5 \\ -7 & 6 \end{bmatrix}^{-1} =$$

$$a) \begin{bmatrix} -6 & 5 \\ -7 & 6 \end{bmatrix}$$

$$b) \begin{bmatrix} 6 & -5 \\ -7 & 6 \end{bmatrix}$$

$$c) \begin{bmatrix} 6 & 5 \\ 7 & 6 \end{bmatrix}$$

$$d) \begin{bmatrix} 6 & -5 \\ 7 & -6 \end{bmatrix}$$

217. Let $F(\alpha) = \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$, then $[F(\alpha)]^{-1}$ is equal to

a) $F(-\alpha)$

b) $F(\alpha^{-1})$

c) $F(2\alpha)$

d) None of these

218. Let for any matrix M , M^{-1} exist which of the following is not true?

a) $|M^{-1}| = |M|^{-1}$

b) $(M^2)^{-1} = (M^{-1})^2$

c) $(M^T)^{-1} = (M^{-1})^T$

d) $(M^{-1})^{-1} = M$

219. If A and B are square matrices of size $n \times n$ such that

$$A^2 - B^2 = (A - B)(A + B), \text{ then which of the following will be always true?}$$

a) $AB = BA$

b) Either of A or B is a zero matrix

c) Either of A or B is an identity matrix

d) $A = B$

220. $x_1 + 2x_2 + 3x_3 = 2x_1 + 3x_2 + x_3 = 3x_1 + x_2 + 2x_3 = 0$.

This system of equations has

a) Infinite solution

b) No solution

c) No solution

d) Unique solution

221. If A is a 3×4 matrix and B is a matrix such that $A^T B$ and BA^T are both defined, then order of B is

a) 3×4

b) 3×3

c) 4×4

d) 4×3

222. If $X = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, then the value of X^n is

$$a) \begin{bmatrix} 3^n & -4^n \\ n & -n \end{bmatrix}$$

$$b) \begin{bmatrix} 2+n & 5-n \\ n & -n \end{bmatrix}$$

$$c) \begin{bmatrix} 3^n & (-4)^n \\ 1^n & (-1)^n \end{bmatrix}$$

d) None of these

223. Let $f(\alpha) = \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$, where $\alpha \in R$. Then, $(F(\alpha))^{-1}$ is equal to

a) $F(-\alpha)$

b) $F(\alpha^{-1})$

c) $F(2\alpha)$

d) None of these

224. For any square matrix A , AA^T is a

a) Unit matrix

b) Symmetric matrix

c) Skew-symmetric matrix

d) Diagonal matrix

225. If A is a square matrix of order $n \times n$, then $\text{adj}(\text{adj} A)$ is equal to

a) $|A|^n A$

b) $|A|^{n-1} A$

c) $|A|^{n-2} A$

d) $|A|^{n-3} A$

226. If a system of the equations $(\alpha + 1)^3 x + (\alpha + 2)^3 y - (\alpha + 3)^3 = 0$,

$(\alpha + 1)x + (\alpha + 2)y - (\alpha + 3) = 0$, and $x + y - 1 = 0$ is consistent. What is the value of α ?

- a) Possesses a trivial solution only
 c) Does not have a common non-zero solution
- b) Possesses a non-zero unique solution
 d) Has infinitely many solutions
241. Consider the following statements:
 1. A square matrix A is hermitian, if $A = A'$
 2. Let $A = [a_{ij}]$ be a skew-hermitian matrix, then a_{ij} is purely imaginary
 3. All integer powers of a symmetric matrix are symmetric. Which of these is/are correct?
 a) (1) and (2) b) (2) and (3) c) (3) and (1) d) (1), (2) and (3)
242. If $a_1, a_2, a_3, a_4, a_5, a_6$ are in AP with common difference $d \neq 0$, then the system of equations $a_1x + a_2y = a_3, a_4x + a_5y = a_6$ has
 a) Infinite number of solutions b) Unique solution
 c) No solution d) Cannot say anything
243. If I_n is the identity matrix of order n , then $(I_n)^{-1}$ is equal to
 a) Does not exist b) I_n c) 0 d) nI_n
244. If A is a square matrix, then $\text{adj } A^T - (\text{adj } A)^T$ is equal to
 a) $2|A|$ b) $2|A|I$ c) Null matrix d) Unit matrix
245. If $\begin{bmatrix} 2 & -1 & 3 \\ 1 & 3 & -1 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 9 \\ 4 \\ 10 \end{bmatrix}$, then $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ is equal to
 a) $\begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ b) $\begin{bmatrix} 2 \\ 3 \\ 1 \end{bmatrix}$ c) $\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ d) $\begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$
246. Consider the system of equations
 $a_1x + b_1y + c_1z = 0$
 $a_2x + b_2y + c_2z = 0$
 $a_3x + b_3y + c_3z = 0$
 if $\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = 0$, then the system has
 a) More than two solutions
 b) One trivial and one non-trivial solutions
 c) No solution
 d) Only trivial solution (0,0,0)
247. The number of solutions of the system of equations
 $x - y + z = 2$
 $2x + y - z = 5$
 $4x + y + z = 10$ is
 a) ∞ b) 1 c) 2 d) 0
248. If $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$, then which of the following statement is not correct?
 a) A is orthogonal matrix b) A' is orthogonal matrix
 c) Determinant $A = 1$ d) A is not invertible
249. If $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$, then A^{-1} is equal to
 a) $2A$ b) A c) $-A$ d) I
250. The rank of the matrix $\begin{bmatrix} -1 & 2 & 5 \\ 2 & -4 & a-4 \\ 1 & -2 & a+1 \end{bmatrix}$ is
 a) 1 if $a = 6$ b) 2 if $a = 1$ c) 3 if $a = 2$ d) 4 if $a = -6$
251. If $D = \text{diag}(d_1, d_2, d_3, \dots, d_n)$, where $d_i \neq 0$ for all $i = 1, 2, \dots, n$, then D^{-1} is equal to
 a) D b) $\text{diag}(d_1^{-1}, d_2^{-1}, \dots, d_n^{-1})$ c) I_n d) None of these
252. If $f(x) = x^2 + 4x - 5$ and $A = \begin{bmatrix} 1 & 2 \\ 4 & -3 \end{bmatrix}$, then $f(A)$ is equal to

253. If $A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$ and I is the unit matrix of order 2, then A^2 equals
- a) $4A - 3I$ b) $3A - 4I$ c) $A - I$ d) $A + I$
254. Which one of the following is true always for any two non-singular matrices A and B of same order?
- a) $AB = BA$ b) $(AB)^t = A^t B^t$
c) $(A + B)(A - B) = A^2 - B^2$ d) $(AB)^{-1} = B^{-1} A^{-1}$
255. The inverse of $\begin{bmatrix} 0 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 0 \end{bmatrix}$ is
- a) $\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$ b) $\begin{bmatrix} 1/2 & 0 & 2 \\ 0 & 1/2 & 0 \\ 0 & 0 & 1/2 \end{bmatrix}$ c) $\begin{bmatrix} 0 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 0 \end{bmatrix}$ d) $\begin{bmatrix} 0 & 0 & 1/2 \\ 0 & 1/2 & 0 \\ 1/2 & 0 & 0 \end{bmatrix}$
256. The values of a for which the system of equations $ax + y + z = 0$, $x - ay + z = 0$, $x + y + z = 0$ possesses non-zero solution, are given by
- a) 1, 2 b) 1, -1 c) 0 d) None of these
257. If A is square matrix, then
- a) $A + A^T$ is symmetric b) AA^T is skew-symmetric
c) $A^T + A$ is skew-symmetric d) $A^T A$ is skew-symmetric
258. If $A = \begin{bmatrix} 1 & 2 & -1 \\ -1 & 1 & 2 \\ 2 & -1 & 1 \end{bmatrix}$, then $\det [\text{adj}(\text{adj } A)]$ is equal to
- a) 12^4 b) 13^4 c) 14^4 d) None of these
259. If $\begin{bmatrix} 1 & 1 & 1 \\ 1 & -2 & -2 \\ 1 & 3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ 4 \end{bmatrix}$, then $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ is equal to
- a) $\begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ b) $\begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix}$ c) $\begin{bmatrix} 5 \\ -2 \\ 1 \end{bmatrix}$ d) $\begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}$
260. If $A = \begin{bmatrix} 1 & 2 \\ -4 & -1 \end{bmatrix}$, then A^{-1} is
- a) $\frac{1}{7} \begin{bmatrix} -1 & -2 \\ 4 & 1 \end{bmatrix}$ b) $\frac{1}{7} \begin{bmatrix} 1 & 2 \\ -4 & -1 \end{bmatrix}$ c) $\frac{1}{7} \begin{bmatrix} 1 & 2 \\ 4 & 1 \end{bmatrix}$ d) Does not exist
261. If the matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is commutative with the matrix $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, then
- a) $a = 0, b = c$ b) $b = 0, c = d$ c) $c = 0, d = a$ d) $d = 0, a = b$
262. If $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 2 \\ 4 & 5 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$, then AB is equal to
- a) $\begin{bmatrix} 5 & 1 & -3 \\ 3 & 2 & 6 \\ 14 & 5 & 0 \end{bmatrix}$ b) $\begin{bmatrix} 11 & 4 & 3 \\ 1 & 2 & 3 \\ 0 & 3 & 3 \end{bmatrix}$ c) $\begin{bmatrix} 1 & 8 & 4 \\ 2 & 9 & 6 \\ 0 & 2 & 0 \end{bmatrix}$ d) $\begin{bmatrix} 0 & 1 & 2 \\ 5 & 4 & 3 \\ 1 & 8 & 2 \end{bmatrix}$
263. Let A be a skew-symmetric matrix of odd order, then $|A|$ is equal to
- a) 0 b) 1 c) -1 d) None of these
264. If $P = \begin{bmatrix} \sqrt{3}/2 & 1/2 \\ -1/2 & \sqrt{3}/2 \end{bmatrix}$, $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $Q = PAP^T$, then $P^T Q^{2005} P$ is
- a) $\begin{bmatrix} 1 & 2005 \\ 0 & 1 \end{bmatrix}$ b) $\begin{bmatrix} 1 & 2005 \\ 2005 & 1 \end{bmatrix}$
c) $\begin{bmatrix} 1 & 0 \\ 2005 & 1 \end{bmatrix}$ d) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
265. If X and Y are 2×2 matrices such that $2X + 3Y = O$ and $X + 2Y = I$, where O and I denote the 2×2 zero matrix and the 2×2 identity matrix, then X is equal to

a) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

b) $\begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$

c) $\begin{bmatrix} -3 & 0 \\ 0 & -3 \end{bmatrix}$

d) $\begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$

266. Consider the system of linear equations

$$x_1 + 2x_2 + x_3 = 3$$

$$2x_1 + 3x_2 + x_3 = 3$$

$$3x_1 + 5x_2 + 2x_3 = 1$$

The system has

a) Infinite number of solutions

b) Exactly 3 solutions

c) A unique solution

d) No solution

267.

If $A = \begin{bmatrix} x & -2 \\ 3 & 7 \end{bmatrix}$ and $A^{-1} = \begin{bmatrix} \frac{7}{34} & \frac{1}{17} \\ -\frac{3}{34} & \frac{2}{17} \end{bmatrix}$, then the value of x is

a) 2

b) 3

c) -4

d) 4

268.

If $A = \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix}$. The only correct statement about the matrix A is

a) A is a zero matrix

b) $A = (-1)I$, where I is a unit matrix

c) A^{-1} does not exist

d) $A^2 = I$

269.

The inverse of the matrix $\begin{bmatrix} 1 & 3 \\ 3 & 10 \end{bmatrix}$ is equal to

a) $\begin{bmatrix} 10 & 3 \\ 3 & 1 \end{bmatrix}$

b) $\begin{bmatrix} 10 & -3 \\ -3 & 1 \end{bmatrix}$

c) $\begin{bmatrix} 1 & 3 \\ 3 & 10 \end{bmatrix}$

d) $\begin{bmatrix} -1 & -3 \\ -3 & -10 \end{bmatrix}$

270.

If $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 2 \\ 4 & 5 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$, then AB is equal to

a) $\begin{bmatrix} 5 & 1 & -3 \\ 3 & 2 & 6 \\ 14 & 5 & 0 \end{bmatrix}$

b) $\begin{bmatrix} 11 & 4 & 3 \\ 1 & 2 & 3 \\ 0 & 3 & 3 \end{bmatrix}$

c) $\begin{bmatrix} 1 & 8 & 4 \\ 2 & 9 & 6 \\ 0 & 2 & 0 \end{bmatrix}$

d) $\begin{bmatrix} 0 & 1 & 2 \\ 5 & 4 & 3 \\ 1 & 8 & 2 \end{bmatrix}$

271.

If $F(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $G(y) = \begin{bmatrix} \cos y & 0 & \sin y \\ 0 & 1 & 0 \\ -\sin y & 0 & \cos y \end{bmatrix}$, then $[F(x)G(y)]^{-1}$ is equal to

a) $F(-x)G(-y)$

b) $F(x^{-1})G(y^{-1})$

c) $G(-y)F(-x)$

d) $G(y^{-1})F(x^{-1})$

272.

Let $A = \begin{bmatrix} 1 & 2 \\ -5 & 1 \end{bmatrix}$ and $A^{-1} = xA + yI$, then the value of x and y are

a) $x = \frac{-1}{11}, y = \frac{2}{11}$

b) $x = \frac{-1}{11}, y = \frac{-2}{11}$

c) $x = \frac{1}{11}, y = \frac{2}{11}$

d) $x = \frac{1}{11}, y = \frac{-2}{11}$

273.

If A^T, B^T are transpose matrices of the square matrices A, B respectively, then $(AB)^T$ is equal to

a) $A^T B^T$

b) AB^T

c) BA^T

d) $B^T A^T$

274.

If $\begin{bmatrix} x+y+z \\ x+y \\ y+z \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \\ 7 \end{bmatrix}$, then the value of (x, y, z) is

a) (4, 3, 2)

b) (3, 2, 4)

c) (2, 3, 4)

d) None of the above

275.

If $A = \begin{bmatrix} ab & b^2 \\ -a^2 & -ab \end{bmatrix}$, then A is equal to

a) Idempotent

b) Involutary

c) Nilpotent

d) Scalar

276.

For non-singular square matrices A, B and C of the same order, $(AB^{-1}C)^{-1}$ is equal to

a) $A^{-1}BC^{-2}$

b) $C^{-1}B^{-1}A^{-1}$

c) CBA^{-1}

d) $C^{-1}BA^{-1}$

277.

The matrix $\begin{bmatrix} \lambda - 1 & 4 \\ -3 & 0 & 1 \\ -1 & 1 & 2 \end{bmatrix}$ is invertible, if

a) $\lambda \neq -17$

b) $\lambda \neq -18$

c) $\lambda \neq -19$

d) $\lambda \neq -20$

278.

If a, b, c are non-zero, then the number of solutions of following system of equation is

$$\frac{2x^2}{a^2} - \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0 \quad \dots(i)$$

$$-\frac{x^2}{a^2} + \frac{2y^2}{b^2} - \frac{z^2}{c^2} = 0 \quad \dots(ii)$$

- b) Skew-symmetric matrix
 c) Null matrix
 d) Unit matrix
293. If A is any $m \times n$ matrix such that AB and BA are both defined, then B is an
 a) $m \times n$ matrix b) $n \times m$ matrix c) $n \times n$ matrix d) $m \times m$ matrix
294. If A is a square matrix of order $n \times n$ and k is a scalar, then $\text{adj}(kA)$ is equal to
 a) $k \text{adj} A$ b) $k^n \text{adj} A$ c) $k^{n-1} \text{adj} A$ d) $k^{n+1} \text{adj} A$
295. $x + ky - z = 0, 3x - ky - z = 0$ and $x - 3y + z = 0$ has non-zero solution for k is equal to
 a) -1 b) 0 c) 1 d) 2
296. If $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, I is the unit matrix of order 2 and a, b are arbitrary constants, then $(aI + bA)^2$ is equal to
 a) $a^2I - abA$ b) $a^2I + 2abA$ c) $a^2I + b^2A$ d) None of the above
297. If A is an orthogonal matrix, then
 a) $|A| = 0$ b) $|A| = \pm 1$ c) $|A| = \pm 2$ d) None of these
298. Given $2x - y + 2z = 2, x - 2y + 2z = -4, x + y + \lambda z = 4$ then the value of λ such that the given system of equations has no solution, is
 a) 3 b) 1 c) 0 d) -3
299. If $A = \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & x \end{bmatrix}$ is an idempotent matrix, then x is equal to
 a) -5 b) -1 c) -3 d) -4
300. If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 3 & 0 \\ 0 & 1 & 2 \end{bmatrix}$ and $\text{adj} A = \begin{bmatrix} 6 & -2 & -6 \\ -4 & 2 & x \\ y & -1 & -1 \end{bmatrix}$, then $x + y =$
 a) 6 b) -1 c) 3 d) 1
301. If $\begin{bmatrix} x + y & 2x + z \\ x - y & 2z + w \end{bmatrix} = \begin{bmatrix} 4 & 7 \\ 0 & 10 \end{bmatrix}$, then the value of x, y, z, w are
 a) 2, 2, 3, 4 b) 2, 3, 1, 2 c) 3, 3, 0, 1 d) None of these
302. If for a matrix $A, A^2 + I = O$, where I is the identity matrix, then A equals
 a) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ b) $\begin{bmatrix} -i & 0 \\ 0 & -i \end{bmatrix}$ c) $\begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix}$ d) $\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$
303. If $m[-3 \ 4] + n[4 \ -3] = [10 \ -11]$, then $3m + 7n$ is equal to
 a) 3 b) 5 c) 10 d) 1
304. If $A = \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix}$ and $A(\text{adj} A) = k \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, then the value of k is
 a) $\sin x \cos x$ b) 1 c) 2 d) 3
305. If $A = \begin{bmatrix} 3 & 2 \\ 0 & 1 \end{bmatrix}$, then $(A^{-1})^3$ is equal to
 a) $\frac{1}{27} \begin{bmatrix} 1 & -26 \\ 0 & 27 \end{bmatrix}$ b) $\frac{1}{27} \begin{bmatrix} -1 & 26 \\ 0 & 27 \end{bmatrix}$ c) $\frac{1}{27} \begin{bmatrix} 1 & -26 \\ 0 & -27 \end{bmatrix}$ d) $\frac{1}{27} \begin{bmatrix} -1 & -26 \\ 0 & -27 \end{bmatrix}$
306. Let $M = [a_{uv}]_{n \times n}$ be a matrix, where $a_{uv} = \sin(\theta_u - \theta_v) + i \cos(\theta_u - \theta_v)$, the M is equal to
 a) \bar{M} b) $-M$ c) \bar{M}^T d) $-\bar{M}^T$
307. If the system of homogeneous equations $2x - y + z = 0, x - 2y + z = 0, \lambda x - y + 2z = 0$ has infinitely many solutions, then
 a) $\lambda = 5$ b) $\lambda = -5$ c) $\lambda \neq \pm 5$ d) None of these
308. Assuming that the sum and product given below are defined, which of the following is not true for matrices?
 a) $A + B = B + A$ b) $AB = AC$ does not imply $B = C$
 c) $AB = O$ implies $A = O$ or $B = O$ d) $(AB)' = B'A'$
309. If $E(\theta) = \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix}$ and θ and ϕ differ by an odd multiple of $\frac{\pi}{2}$, then $E(\theta) E(\phi)$ is a
 a) Unit matrix b) Null matrix c) Diagonal matrix d) None of these
310. Consider the system of equations in x, y, z as

$$x \sin 3\theta - y + z = 0$$

$$x \cos 2\theta + 4y + 3z = 0$$

$$\text{and } 2x + 7y + 7z = 0$$

If this system has a non-trivial solution, then for integer n , values of θ are given by

a) $\pi \left(n + \frac{(-1)^n}{3} \right)$ b) $\pi \left(n + \frac{(-1)^n}{4} \right)$ c) $\pi \left(n + \frac{(-1)^n}{6} \right)$ d) $\frac{n\pi}{2}$

311. If $A = \begin{bmatrix} 0 & 2 \\ 3 & -4 \end{bmatrix}$ and $kA = \begin{bmatrix} 0 & 3a \\ 2b & 24 \end{bmatrix}$, then the values of k, a, b are respectively

a) $-6, -12, -18$ b) $-6, 4, 9$ c) $-6, -4, -9$ d) $-6, 12, 18$

312. If A is a non-singular matrix of order 3, then $\text{adj}(\text{adj } A)$ is equal to

a) A b) A^{-1} c) $\frac{1}{|A|}A$ d) $|A|A$

313. If A is a matrix such that $A^2 = A + I$, where I is the unit matrix, then A^5 is equal to

a) $5A + I$ b) $5A + 2I$ c) $5A + 3I$ d) $5A + 4I$

314. If A and B are two matrices such that both $A+B$ and AB are defined, then

a) A and B are of same order b) A is of order $m \times m$ and B is of order $n \times n$
c) Both A and B are of same order $n \times n$ d) A is of order $m \times n$ and B is of order $n \times m$

315. If $U = [2 \ -3 \ 4], X = [0 \ 2 \ 3], V = \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}$ and $Y = \begin{bmatrix} 2 \\ 2 \\ 4 \end{bmatrix}$, then $UV + XY$

a) $[20]$ b) 20 c) $[-20]$ d) -20

316. The matrix $A = \begin{bmatrix} a & 2 \\ 2 & 4 \end{bmatrix}$ is singular, if

a) $a \neq 1$ b) $a = 1$ c) $a = 0$ d) $a = -1$

317. If $A = \begin{bmatrix} 1 & x \\ x^2 & 4y \end{bmatrix}$ and $B = \begin{bmatrix} -3 & 1 \\ 1 & 0 \end{bmatrix}$, $\text{adj } A + B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$,

Then values of x and y are

a) $1, 1$ b) $\pm 1, 1$ c) $1, 0$ d) None of these

318. $A = \begin{bmatrix} \cos \alpha - \sin \alpha & 0 \\ \sin \alpha & \cos \alpha \\ 0 & 0 & 1 \end{bmatrix}$, then A^{-1} is

a) A b) $-A$ c) $\text{adj}(A)$ d) $-\text{adj}(A)$

319. Which of the following is incorrect?

a) $A^2 - B^2 = (A + B)(A - B)$
b) $(A^T)^T = A$
c) $(AB)^n = A^n B^n$ where A, B commute
d) $(A - I)(I + A) = 0 \Leftrightarrow A^2 = I$

320. If $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $f(x) = \frac{1+x}{1-x}$, then $f(A)$ is

a) $\begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix}$ b) $\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ c) $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ d) None of these

321. If A is an invertible matrix, then $\det(A^{-1})$ is equal to

a) $\det(A)$ b) $\frac{1}{\det(A)}$ c) 1 d) None of these

322. If A and B are square matrices of the same order and $AB = 3I$, then A^{-1} is equal to

a) $3B$ b) $\frac{1}{3}B$ c) $3B^{-1}$ d) $\frac{1}{3}B^{-1}$

323. If the points $(x_1, y_1), (x_2, y_2)$ and (x_3, y_3) are collinear,

then the rank of the matrix $\begin{bmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{bmatrix}$ will always be less than

a) 2 b) 3 c) 1 d) None of these

324. If $A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$ and I is the unit matrix of order 2, then A^2 equals

325. a) $4A - 3I$ b) $3A - 4I$ c) $A - I$ d) $A + I$
 If $A = \begin{bmatrix} -2 & 6 \\ -5 & 7 \end{bmatrix}$, then $\text{adj } A$ is
326. a) $\begin{bmatrix} 7 & -6 \\ 5 & -2 \end{bmatrix}$ b) $\begin{bmatrix} 2 & -6 \\ 5 & -7 \end{bmatrix}$ c) $\begin{bmatrix} 7 & -5 \\ 6 & -2 \end{bmatrix}$ d) None of these
 If $A = \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix}$ then $A \cdot (\text{adj } A)$ is equal to
327. a) A b) $|A|$ c) $|A|I$ d) None of these
 Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$, $a, b \in N$. Then
- a) There exist more than one but finite number of B 's such that $AB = BA$
 b) There exist exactly one B such $AB = BA$
 c) There exists infinitely many B 's such that $AB = BA$
 d) There cannot exist any B such that $AB = BA$
328. If $2A + 3B = \begin{bmatrix} 2 & -1 & 4 \\ 3 & 2 & 5 \end{bmatrix}$ and $A + 2B = \begin{bmatrix} 5 & 0 & 3 \\ 1 & 6 & 2 \end{bmatrix}$, then B is
- a) $\begin{bmatrix} 8 & -1 & 2 \\ -1 & 10 & -1 \end{bmatrix}$ b) $\begin{bmatrix} 8 & 1 & 2 \\ -1 & 10 & -1 \end{bmatrix}$ c) $\begin{bmatrix} 8 & 1 & -2 \\ -1 & 10 & -1 \end{bmatrix}$ d) $\begin{bmatrix} 8 & 1 & 2 \\ 1 & 10 & 1 \end{bmatrix}$
329. If I is unit matrix of order 10, then the determinant of I is equal to
- a) 10 b) 1 c) $1/10$ d) 9
330. Let $A = \begin{bmatrix} 3 & 5 \\ 2 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 17 \\ 0 & -10 \end{bmatrix}$, then $|AB|$ is equal to
- a) 80 b) 100 c) -110 d) 92
331. If $A = \begin{bmatrix} 6 & 8 & 5 \\ 4 & 2 & 3 \\ 9 & 7 & 1 \end{bmatrix}$ is the sum of a symmetric matrix B and skew-symmetric matrix C , then B is
- a) $\begin{bmatrix} 6 & 6 & 7 \\ 6 & 2 & 5 \\ 7 & 5 & 1 \end{bmatrix}$ b) $\begin{bmatrix} 0 & 2 & -2 \\ -2 & 5 & -2 \\ 2 & 2 & 0 \end{bmatrix}$ c) $\begin{bmatrix} 6 & 6 & 7 \\ -6 & 2 & -5 \\ -7 & 5 & 1 \end{bmatrix}$ d) $\begin{bmatrix} 0 & 6 & -2 \\ 2 & 0 & -2 \\ -2 & -2 & 0 \end{bmatrix}$
332. Let A be a non-singular square matrix. Then, $|\text{adj } A|$ is equal to
- a) $|A|^n$ b) $|A|^{n-1}$ c) $|A|^{n-2}$ d) None of these
333. If $A = \begin{bmatrix} 1 & -1 \\ 2 & -1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & a \\ 4 & b \end{bmatrix}$ $(A + B)^2 = A^2 + B^2$. Then a and b are respectively
- a) 1, -1 b) 2, -3 c) -1, 1 d) 3, -2
334. For non-singular square matrices A, B and C of same order, $(AB^{-1}C)^{-1}$ is equal to
- a) $A^{-1}BC^{-1}$ b) $C^{-1}B^{-1}A^{-1}$ c) $CB^{-1}A^{-1}$ d) $C^{-1}BA^{-1}$
335. If $A = \begin{bmatrix} 5 & 2 \\ 3 & 1 \end{bmatrix}$, then $A^{-1} =$
- a) $\begin{bmatrix} 1 & -2 \\ -3 & 5 \end{bmatrix}$ b) $\begin{bmatrix} -1 & 2 \\ 3 & -5 \end{bmatrix}$ c) $\begin{bmatrix} -1 & -2 \\ -3 & -5 \end{bmatrix}$ d) $\begin{bmatrix} 1 & 2 \\ 3 & 5 \end{bmatrix}$

: ANSWER KEY :

1)	a	2)	d	3)	b	4)	b	189)	d	190)	a	191)	b	192)	b
5)	b	6)	c	7)	a	8)	b	193)	d	194)	b	195)	a	196)	d
9)	c	10)	d	11)	d	12)	d	197)	b	198)	a	199)	b	200)	d
13)	a	14)	d	15)	a	16)	b	201)	a	202)	a	203)	d	204)	b
17)	c	18)	a	19)	d	20)	b	205)	c	206)	a	207)	a	208)	a
21)	c	22)	b	23)	d	24)	d	209)	a	210)	a	211)	c	212)	c
25)	a	26)	b	27)	d	28)	b	213)	a	214)	b	215)	a	216)	a
29)	c	30)	a	31)	b	32)	d	217)	a	218)	b	219)	a	220)	c
33)	c	34)	a	35)	a	36)	a	221)	a	222)	d	223)	a	224)	b
37)	c	38)	c	39)	d	40)	c	225)	c	226)	d	227)	a	228)	c
41)	a	42)	a	43)	d	44)	a	229)	c	230)	c	231)	a	232)	a
45)	c	46)	d	47)	c	48)	c	233)	a	234)	d	235)	c	236)	c
49)	a	50)	a	51)	d	52)	a	237)	c	238)	d	239)	a	240)	d
53)	c	54)	c	55)	d	56)	d	241)	b	242)	b	243)	b	244)	c
57)	c	58)	d	59)	a	60)	b	245)	c	246)	a	247)	b	248)	d
61)	a	62)	c	63)	a	64)	d	249)	b	250)	b	251)	b	252)	d
65)	d	66)	b	67)	b	68)	b	253)	a	254)	d	255)	d	256)	b
69)	b	70)	c	71)	b	72)	a	257)	a	258)	c	259)	b	260)	a
73)	b	74)	d	75)	d	76)	a	261)	c	262)	a	263)	a	264)	a
77)	c	78)	d	79)	d	80)	a	265)	c	266)	d	267)	d	268)	d
81)	b	82)	b	83)	a	84)	b	269)	b	270)	a	271)	c	272)	a
85)	a	86)	c	87)	b	88)	d	273)	d	274)	c	275)	c	276)	d
89)	a	90)	d	91)	c	92)	a	277)	a	278)	d	279)	b	280)	a
93)	d	94)	a	95)	d	96)	b	281)	b	282)	d	283)	d	284)	b
97)	b	98)	b	99)	d	100)	b	285)	a	286)	d	287)	a	288)	c
101)	b	102)	b	103)	b	104)	a	289)	c	290)	b	291)	c	292)	b
105)	b	106)	d	107)	b	108)	b	293)	b	294)	c	295)	c	296)	b
109)	c	110)	b	111)	b	112)	a	297)	b	298)	b	299)	c	300)	a
113)	a	114)	a	115)	d	116)	c	301)	a	302)	b	303)	d	304)	b
117)	d	118)	d	119)	b	120)	d	305)	a	306)	d	307)	a	308)	c
121)	a	122)	d	123)	a	124)	a	309)	b	310)	c	311)	c	312)	d
125)	c	126)	a	127)	d	128)	d	313)	c	314)	c	315)	a	316)	b
129)	b	130)	b	131)	a	132)	d	317)	a	318)	c	319)	a	320)	a
133)	d	134)	b	135)	d	136)	c	321)	b	322)	b	323)	b	324)	a
137)	a	138)	a	139)	c	140)	b	325)	a	326)	c	327)	c	328)	b
141)	b	142)	a	143)	b	144)	a	329)	b	330)	b	331)	a	332)	b
145)	c	146)	a	147)	b	148)	a	333)	a	334)	d	335)	b		
149)	a	150)	d	151)	c	152)	a								
153)	b	154)	d	155)	b	156)	a								
157)	a	158)	a	159)	a	160)	a								
161)	a	162)	d	163)	c	164)	c								
165)	a	166)	a	167)	c	168)	d								
169)	b	170)	d	171)	c	172)	b								
173)	c	174)	c	175)	a	176)	c								
177)	b	178)	a	179)	a	180)	b								
181)	d	182)	a	183)	a	184)	a								
185)	c	186)	c	187)	b	188)	a								

: HINTS AND SOLUTIONS :

1 (a)

Let $A = \begin{bmatrix} 2 & -1 \\ 3 & -2 \end{bmatrix}$. Then,

$$A^2 = AA = \begin{bmatrix} 2 & -1 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ 3 & -2 \end{bmatrix}$$

$$\Rightarrow A^2 = \begin{bmatrix} 4-3 & -2+2 \\ 6-6 & -3+4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

 $\therefore A^3 = A, A^4 = A^2 = I, A^2 = A^4 A = IA = A$ etc.
Hence, $A^n = \begin{cases} A, & \text{if } n \text{ is odd} \\ I, & \text{if } n \text{ is even} \end{cases}$

2 (d)

$$\begin{bmatrix} 0 & a \\ b & 0 \end{bmatrix}^2 = \begin{bmatrix} 0 & a \\ b & 0 \end{bmatrix} \begin{bmatrix} 0 & a \\ b & 0 \end{bmatrix} = \begin{bmatrix} ab & 0 \\ 0 & ab \end{bmatrix}$$

$$\text{and } \begin{bmatrix} 0 & a \\ b & 0 \end{bmatrix}^4 = \begin{bmatrix} 0 & a \\ b & 0 \end{bmatrix}^2 \begin{bmatrix} 0 & a \\ b & 0 \end{bmatrix}^2$$

$$= \begin{bmatrix} a^2 b^2 & 0 \\ 0 & a^2 b^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad [\text{given}]$$

$$\Rightarrow a^2 b^2 = 1 \Rightarrow ab = 1$$

3 (b)

Given matrix A is a square matrixAnd $AA^T = I = A^T A$

$$\Rightarrow |AA^T| = |I| = |A^T A|$$

$$\Rightarrow |A||A^T| = 1 = |A^T||A|$$

$$\Rightarrow |A|^2 = 1 \quad [\because A \cdot A^T = |A|^2]$$

$$\Rightarrow |A| = \pm 1$$

4 (b)

$$\text{Let } I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{ then, } kI = \begin{bmatrix} k & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & k \end{bmatrix}$$

$$\Rightarrow \text{adj}(kI) = \begin{bmatrix} k^2 & 0 & 0 \\ 0 & k^2 & 0 \\ 0 & 0 & k^2 \end{bmatrix} = k^2 I$$

5 (b)

Given, A is singular $\Rightarrow |A| = 0$ Now, $A(\text{adj} A) = |A|I_n = 0$ $\therefore A(\text{adj} A) = 0$ i.e., $A(\text{adj} A)$ is a zero matrix.

6 (c)

Given, $AB = O$

$$\therefore \begin{bmatrix} \cos^2 \alpha & \cos \alpha \sin \alpha \\ \cos \alpha \sin \alpha & \sin^2 \alpha \end{bmatrix} \times \begin{bmatrix} \cos^2 \beta & \cos \beta \sin \beta \\ \cos \beta \sin \beta & \sin^2 \beta \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} \cos \alpha \cos \beta \cos(\alpha - \beta) & \cos \alpha \sin \beta \cos(\alpha - \beta) \\ \cos \beta \sin \alpha \cos(\alpha - \beta) & \sin \alpha \sin \beta \cos(\alpha - \beta) \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\Rightarrow \cos(\alpha - \beta) = 0$$

$$\Rightarrow \alpha - \beta \text{ is an odd multiple of } \frac{\pi}{2}$$

8 (b)

$$\text{Given, } A = \begin{bmatrix} 2 & 4 & 5 \\ 4 & 8 & 10 \\ -6 & -12 & -15 \end{bmatrix}$$

Applying $R_2 \rightarrow R_2 - 2R_1$ and $R_3 \rightarrow R_3 + 3R_1$

$$\Rightarrow A = \begin{bmatrix} 2 & 4 & 5 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Since the equivalent matrix in echelon form has only one non-zero row,

$$\therefore \text{Rank}(A) = 1$$

9 (c)

 $\therefore AB = B, BA = A$

$$\therefore A^2 + B^2 = AA + BB = A(BA) + B(AB)$$

$$= (AB)A + (BA)B = BA + AB$$

$$= A + B$$

$$\Rightarrow A^2 = A \text{ and } B^2 = B$$

10 (d)

$$|A| = 7(1 - 0) + 3(-1 - 0) - 3(0 + 1) = 1$$

Cofactors of matrix A are

$$C_{11} = 1, \quad C_{12} = 1, \quad C_{13} = 1$$

$$C_{21} = 3, \quad C_{22} = 4, \quad C_{23} = 3$$

$$C_{31} = 3, \quad C_{32} = 3, \quad C_{33} = 4$$

$$\therefore \text{adj}(A) = \begin{bmatrix} 1 & 1 & 1 \\ 3 & 4 & 3 \\ 3 & 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$$

11 (d)

It is given that A is a skew-symmetric matrix

$$\therefore A^T = -A$$

$$\Rightarrow (A^T)^n = (-A)^n$$

$$\Rightarrow (A^n)^T = (-1)^n A^n$$

$$\Rightarrow (A^n)^T = \begin{cases} A^n & \text{if } n \text{ is even} \\ -A^n & \text{if } n \text{ is odd} \end{cases}$$

Hence, A^n is skew-symmetric when n is odd

12 (d)

Consider $\begin{bmatrix} 1 & * & * \\ * & 1 & * \\ * & * & 1 \end{bmatrix}$. By placing a_1 in any one of the

6*

Position and 0 elsewhere. We get 6 non-singular matrices.

Similarly, $\begin{bmatrix} * & * & 1 \\ * & 1 & * \\ 1 & * & * \end{bmatrix}$ gives at least one non-singular.

13 (a)

We know that, $\text{Tr}(A) = \sum_{i=1}^3 a_{ii}$

$$\text{Tr}(A) = 1 + 7 + 9 = 17$$

14 (d)

$$\text{Now, } A + B = \begin{bmatrix} 3 & 4 \\ 2 & 4 \end{bmatrix} + \begin{bmatrix} -2 & -2 \\ 0 & -2 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 2 & 2 \end{bmatrix}$$

$$\Rightarrow |A + B| = \begin{vmatrix} 1 & 2 \\ 2 & 2 \end{vmatrix} = 2 - 4 = -2$$

$$\text{Also, } \text{adj}(A + B) = \begin{bmatrix} 2 & -2 \\ -2 & 1 \end{bmatrix}$$

$$\therefore (A + B)^{-1} = -\frac{1}{2} \begin{bmatrix} 2 & -2 \\ -2 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} -1 & 1 \\ 1 & -1/2 \end{bmatrix}$$

$$A^{-1} = \frac{1}{4} \begin{bmatrix} 4 & -4 \\ -2 & 3 \end{bmatrix} \text{ and } B^{-1} = \frac{1}{4} \begin{bmatrix} -2 & 2 \\ 0 & -2 \end{bmatrix}$$

$$\therefore A^{-1} + B^{-1} = \begin{bmatrix} 1 & -1 \\ -1/2 & 3/4 \end{bmatrix} + \begin{bmatrix} -1/2 & 1/2 \\ 0 & -1/2 \end{bmatrix}$$

$$= \begin{bmatrix} 1/2 & -1/2 \\ -1/2 & 1/4 \end{bmatrix}$$

$$\therefore (A + B)^{-1} \neq A^{-1} + B^{-1}$$

15 (a)

$$f(A) = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix} - 5 \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$$

$$= \begin{bmatrix} 8 & 5 \\ -5 & 3 \end{bmatrix} - \begin{bmatrix} 15 & 5 \\ -5 & 10 \end{bmatrix} = \begin{bmatrix} -7 & 0 \\ 0 & -7 \end{bmatrix}$$

16 (b)

$$C_{11} = 1, C_{12} = -2, C_{13} = -2$$

$$C_{21} = -1, C_{22} = 3, C_{23} = 3$$

$$C_{31} = 0, C_{32} = -4, C_{33} = -3$$

$$\therefore \text{adj}(A) = \begin{bmatrix} 1 & -2 & -2 \\ -1 & 3 & 3 \\ 0 & -4 & -3 \end{bmatrix}^T = \begin{bmatrix} 1 & -1 & 0 \\ -2 & 3 & -4 \\ -2 & 3 & -3 \end{bmatrix}$$

17 (c)

Since, the rank of given matrix is 1, then

$$\begin{bmatrix} 2 & 5 \\ -4 & a - 4 \end{bmatrix} = 0$$

$$\Rightarrow 2a - 8 + 20 = 0$$

$$\Rightarrow a = -6$$

18 (a)

Since, given equations have a non-trivial solution

$$\therefore \Delta = 0$$

$$\Rightarrow \begin{vmatrix} 1 & 3 & \lambda \\ 2 & 4 & -1 \\ 1 & 5 & -2 \end{vmatrix} = 0$$

$$\Rightarrow 1(-8 + 5) - 3(-4 + 1) + \lambda(10 - 4) = 0$$

$$\Rightarrow 6\lambda = -6 \Rightarrow \lambda = -1$$

20 (b)

Since, $|a| \neq 0$ So, A^{-1} exists

$$\therefore AB = AC$$

$$\Rightarrow A^{-1}(AB) = A^{-1}(AC)$$

$$\Rightarrow (A^{-1}A)B = (A^{-1}A)C \Rightarrow B = C$$

21 (c)

$$\text{Let } A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 4 & 10 \end{bmatrix}, B = \begin{bmatrix} 1 \\ \alpha \\ \alpha^2 \end{bmatrix} \text{ and } X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$\text{For consistent, } |A| = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 4 & 10 \end{vmatrix} = 0$$

$$\Rightarrow 1(20 - 16) - 1(10 - 4) + 1(4 - 2) = 0$$

$$\Rightarrow 4 - 6 + 2 = 0$$

$$\Rightarrow 0 = 0$$

and $(\text{adj } A)B = 0$

$$\Rightarrow \begin{bmatrix} 4 & -6 & 2 \\ -6 & 9 & -3 \\ 2 & -3 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ \alpha \\ \alpha^2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 4 - 6\alpha + 2\alpha^2 \\ -6 + 9\alpha - 3\alpha^2 \\ 2 - 3\alpha + \alpha^2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow 2\alpha^2 - 6\alpha + 4 = 0, -3\alpha^2 + 9\alpha - 6 = 0$$

and $\alpha^2 - 3\alpha + 2 = 0$

Now, $2\alpha^2 - 6\alpha + 4 = 0$

$$\Rightarrow (2\alpha - 2)(\alpha - 2) = 0$$

$$\Rightarrow \alpha = 1, 2$$

Similarly from other equations we also get the same value

22 (b)

Given, $B = A^6 - A^5$, where $A^3 = A + I$

$$\Rightarrow B = (A^3)^2 - A^3 A^2$$

$$= (A + I)^2 - (A + I)A^2$$

$$= A^2 + I^2 + 2AI - A^3 - A^2 I$$

$$= I + 2A - (A + I)$$

$$\Rightarrow B = A$$

$$\therefore \text{Inverse of } B = A^{-1}$$

23 (d)

Since, $A^T A = I$

$$\Rightarrow \begin{bmatrix} a & b & c \\ b & c & a \\ c & a & b \end{bmatrix} \begin{bmatrix} a & b & c \\ b & c & a \\ c & a & b \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} a^2 + b^2 + c^2 & ab + bc + ca & ab + bc + ca \\ ab + bc + ca & a^2 + b^2 + c^2 & ab + bc + ca \\ ab + bc + ca & ab + bc + ca & a^2 + b^2 + c^2 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow a^2 + b^2 + c^2 = 1 \text{ and } ab + bc + ca = 0$$

Now, $(a + b + c)^2 =$

$$a^2 + b^2 + c^2 + 2(ab + bc + ca)$$

$$= 1 + 2 \cdot 0 = 1$$

$$\Rightarrow a + b + c = 1 \dots (i)$$

Now, $(a^3 + b^3 + c^3) = (a + b + c)$

$$(a^2 + b^2 + c^2 - ab - bc - ca) + 3abc$$

$$= (a + b + c) + 3$$

$$\Rightarrow a^3 + b^3 + c^3 = 1 + 3 = 4 \text{ [using Eq. (i)]}$$

24 (d)

$$A^2 = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 8 \\ 8 & 8 & 9 \end{bmatrix}$$

Now,

$$A^2 - 4A = \begin{bmatrix} 9 & 8 & 8 \\ 8 & 9 & 8 \\ 8 & 8 & 9 \end{bmatrix} - \begin{bmatrix} 4 & 8 & 8 \\ 8 & 4 & 8 \\ 8 & 8 & 4 \end{bmatrix} = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix} = 5I_3$$

25 (a)

We have,

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

$$\Rightarrow A^2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2$$

$$\Rightarrow A^4 = A^2 A^2 = I_2 I_2 = I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

26 (b)

The system of given equation are

$$Kx + 2y - z = 1 \quad \dots(i)$$

$$(K - 1)y - 2z = 2 \quad \dots(ii)$$

$$\text{and } (K + 2)z = 3 \quad \dots(iii)$$

This system of equations has a unique solution, if

$$\begin{vmatrix} K & 2 & -1 \\ 0 & K-1 & -2 \\ 0 & 0 & K+2 \end{vmatrix} \neq 0$$

$$\Rightarrow (K - 2)(K)(K - 1) \neq 0$$

$$\Rightarrow K \neq 2, 0, 1$$

ie, $K = -1$, is a required answer.

27 (d)

Given, $x + y + z = 6$, $x + 2y + 3z = 10$ and $x + 2y + \lambda z = \mu$

$$\text{For no solution, } \Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 2 & \lambda \end{vmatrix} = 0$$

$$\Rightarrow 1(2\lambda - 6) - 1(\lambda - 3) + 1(2 - 2) = 0$$

$$\lambda - 3 = 0 \Rightarrow \lambda = 3$$

$$\text{and } \Delta_1 = \begin{vmatrix} 6 & 1 & 1 \\ 10 & 2 & 3 \\ \mu & 2 & 3 \end{vmatrix} \neq 0$$

$$\Rightarrow 6(6 - 6) - 1(30 - 3\mu) + 1(20 - 2\mu) \neq 0$$

$$\Rightarrow \mu - 10 \neq 0 \Rightarrow \mu \neq 10$$

28 (b)

$$|AA^T| = |I| = |A^T A|$$

$$\Rightarrow |A||A^T| = 1 = |A^2||A|$$

$$\Rightarrow |A|^2 = 1 \quad [\because |A^T| = |A|]$$

$$\Rightarrow |A| = \pm 1$$

29 (c)

AB

$$= \begin{bmatrix} \cos^2 \theta & \sin \theta \cos \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix} \begin{bmatrix} \cos^2 \phi & \sin \phi \cos \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 \theta \cos^2 \phi + \sin \theta \cos \theta \sin \phi \cos \phi & \cos^2 \theta \sin \phi \cos \phi + \sin \theta \cos \theta \sin^2 \phi \\ \cos^2 \theta \sin \phi \cos \phi + \sin^2 \theta \sin \theta \cos \theta & \cos^2 \theta \sin^2 \phi + \sin \theta \cos \theta \sin \phi \cos \phi \end{bmatrix}$$

$$= \begin{bmatrix} \cos \theta \cos \phi \cos(\theta - \phi) & \sin \phi \cos \theta \cos(\theta - \phi) \\ \sin \theta \cos \phi \cos(\theta - \phi) & \sin \theta \sin \phi \cos(\theta - \phi) \end{bmatrix}$$

$$\because AB = O$$

$$\Rightarrow \cos(\theta - \phi) = 0$$

$$\Rightarrow \cos(\theta - \phi) = \cos(2n + 1) \frac{\pi}{2}$$

$$\Rightarrow \theta = (2n + 1) \frac{\pi}{2} + \phi, \text{ where } n = 0, 1, 2, \dots$$

30 (a)

We have,

$$AB = A \text{ and } BA = B$$

Now, $AB = A$

$$\Rightarrow (AB)A = AA$$

$$\Rightarrow A(BA) = A^2$$

$$\Rightarrow AB = A^2 \quad [\because BA = B]$$

$$\Rightarrow A = A^2 \quad [\because AB = A]$$

and, $BA = B$

$$\Rightarrow (BA)B = B^2$$

$$\Rightarrow B(AB) = B^2$$

$$\Rightarrow BA = B^2 \quad [\because AB = A]$$

$$\Rightarrow B = B^2 \quad [\because BA = B]$$

$$\therefore A^2 = A \text{ and } B^2 = B$$

$\Rightarrow A$ and B are idempotent matrices

31 (b)

From given matrix equation, we have

$$x - y - z = 0$$

$$-y + z = 5$$

$$z = 3$$

$$\Rightarrow x = 1, y = -2, z = 3$$

32 (d)

Here, cofactors are

$$C_{11} = \cos 2\theta, \quad C_{12} = -\sin 2\theta$$

$$C_{21} = \sin 2\theta, \quad C_{22} = \cos 2\theta$$

$$\therefore |A| = |\cos^2 2\theta + \sin^2 2\theta| = 1$$

$$\therefore A^{-1} = \frac{1}{|A|} \begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix} = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$$

33 (c)

Since, B is the inverse of A .

$$\text{ie, } B = A^{-1}$$

$$\text{So, } 10A^{-1} = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{bmatrix}$$

$$\Rightarrow 10A^{-1}A = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{bmatrix} A$$

$$= \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{bmatrix} \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 10 & 0 & 0 \\ 0 & 10 & 0 \\ 0 & 0 & 10 \end{bmatrix} = \begin{bmatrix} 10 & 0 & 0 \\ -5 + \alpha & 5 + \alpha & -5 + \alpha \\ 0 & 0 & 10 \end{bmatrix}$$

$$\Rightarrow -5 + \alpha = 0 \Rightarrow \alpha = 5$$

34 (a)

$$AB = \begin{bmatrix} 0 & 3 & 3 \\ -3 & 0 & -4 \\ -3 & 4 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3y + 3z \\ -3x - 4z \\ -3x + 4y \end{bmatrix}$$

$$\text{Now, } B'(AB) = [x \ y \ z] \begin{bmatrix} 3y + 3z \\ -3x - 4z \\ -3x + 4y \end{bmatrix}$$

$$= [3xy + 3zx - 3xy - 4yz - 3xz + 4yz]$$

$$= [0]$$

$\therefore B'(AB)$ is a null matrix.

36 (a)

$$\text{Given, } A = \begin{bmatrix} a & b & 0 \\ -b & a & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow |A| = a^2 + b^2 = 1$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj}(A) = \text{adj}(A)$$

38 (c)

We have,

$$(A + B)^2 = A^2 + 2AB + B^2$$

$$\Leftrightarrow (A + B)(A + B) = A^2 + 2AB + B^2$$

$$\Leftrightarrow A^2 + AB + BA + B^2 = A^2 + 2AB + B^2$$

$$\Leftrightarrow AB = BA$$

$$\text{Hence, } (A + B)^2 = A^2 + 2AB + B^2 \Leftrightarrow AB = BA$$

39 (d)

$$A^{-1} = \frac{1}{2x^2} \begin{bmatrix} x & 0 \\ -x & 2x \end{bmatrix} = \begin{bmatrix} \frac{1}{2x} & 0 \\ -\frac{1}{2x} & \frac{1}{x} \end{bmatrix}$$

$$\text{But } A^{-1} = \begin{bmatrix} 1 & 0 \\ -1 & 2 \end{bmatrix}$$

$$\therefore \begin{bmatrix} \frac{1}{2x} & 0 \\ -\frac{1}{2x} & \frac{1}{x} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ -1 & 2 \end{bmatrix} \Rightarrow \frac{1}{2x} = 1 \Rightarrow x = \frac{1}{2}$$

40 (c)

Given,

$$\therefore \begin{vmatrix} a & a-1 \\ b-1 & b \\ -1 & c & c \end{vmatrix} = 0$$

Applying $C_2 \rightarrow C_2 - C_1, C_3 \rightarrow C_3 - C_1$

$$\Rightarrow \begin{vmatrix} a & 0 & -(1+a) \\ b & -(1+b) & 0 \\ -1 & c+1 & c+1 \end{vmatrix} = 0$$

$$\Rightarrow -a(1+b)(1+c) - b[0 + (1+a)(1+c)]$$

$$-1[0 - (1+a)(1+b)] = 0$$

$$\Rightarrow -a(1+b)(1+c) - b(1+a)(1+c)$$

$$+1(1+a)(1+b) = 0$$

On dividing by $(1+a)(1+b)(1+c)$, we get

$$-\frac{a}{1+a} - \frac{b}{1+b} + \frac{1}{1+c} = 0$$

$$\Rightarrow -\frac{a}{1+a} + 1 - \frac{b}{1+b} + 1 + \frac{1}{1+c} = 2$$

$$\Rightarrow \frac{1}{1+a} + \frac{1}{1+b} + \frac{1}{1+c} = 2$$

41 (a)

The given system of equation has at least one solution, if

$$\begin{vmatrix} 3 & -1 & 4 \\ 1 & 2 & -3 \\ 6 & 5 & \lambda \end{vmatrix} = 0$$

$$\Rightarrow 3(2\lambda + 15) + 1(\lambda + 18) + 4(5 - 12) = 0$$

$$\Rightarrow 7\lambda = -35 \Rightarrow \lambda = -5$$

43 (d)

$$\text{Given, } A = \begin{bmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 1 & 0 & 2 \end{bmatrix}$$

$$C_{11} = 4, C_{12} = 1, C_{13} = -2$$

$$C_{21} = -2, C_{22} = 4, C_{23} = 1$$

$$C_{31} = 1, C_{32} = -2, C_{33} = 4$$

$$\therefore \text{adj}(A) = \begin{bmatrix} 4 & 1 & -2 \\ -2 & 4 & 1 \\ 1 & -2 & 4 \end{bmatrix}^T = \begin{bmatrix} 4 & -2 & 1 \\ 1 & 4 & -2 \\ -2 & 1 & 4 \end{bmatrix}$$

$$\therefore |\text{adj} A| = 4(16 + 2) + 2(4 - 4) + (1 + 8) = 72 + 0 + 9 = 81$$

Alternate

$$|A| \begin{vmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 1 & 0 & 2 \end{vmatrix}$$

$$= 2(4 - 0) - 1(0 - 1) = 9$$

$$\therefore |\text{adj} A| = |A|^{3-1} = (9)^2 = 81$$

44 (a)

Since, $A' = -A$

$$\therefore A^3 = AAA$$

$$\text{And } (A^3)' = A'A'A' = -A^3$$

Hence, matrix A^3 is a skew-symmetric matrix

45 (c)

$$\text{Given, } [1 \ x \ 1] \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 1 \\ 0 & 3 & 2 \end{bmatrix} \begin{bmatrix} x \\ 1 \\ -2 \end{bmatrix} = 0$$

$$\Rightarrow [1 \ x \ 1] \begin{bmatrix} x+2-6 \\ 0+5-2 \\ 0+3-4 \end{bmatrix} = 0$$

$$\Rightarrow [1 \ x \ 1] \begin{bmatrix} x-4 \\ 3 \\ -1 \end{bmatrix} = 0$$

$$\Rightarrow x - 4 + 3x - 1 = 0 \Rightarrow x = \frac{5}{4}$$

46 (d)

$$\text{We have, } A + B = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$$

Clearly, $A + B$ is a symmetric matrix

Now,

$$(A + B)^{-1} = -1 \begin{bmatrix} 3 & -2 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} -3 & 2 \\ 2 & -1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{4} \begin{bmatrix} 4 & -4 \\ -2 & 3 \end{bmatrix}, B^{-1} = \frac{1}{2} \begin{bmatrix} -1 & 2 \\ 0 & -2 \end{bmatrix}$$

$$\therefore A^{-1} + B^{-1} = \frac{1}{4} \begin{bmatrix} 2 & 0 \\ -2 & -1 \end{bmatrix}$$

We observe that $(A + B)^{-1}$ is a symmetric matrix and $(A + B)^{-1} \neq A^{-1} + B^{-1}$

47 (c)

$$\therefore A \cdot A^T = I_n$$

$$\Rightarrow A - I_n = A - AA^T = A(I_n - A^T)$$

$$\Rightarrow |A - I_n| = |A(I_n - A^T)|$$

$$= |A||I_n - A^T|$$

$$= |A||I_n - A|$$

48 (c)

$$A^2 = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1}{2} - \frac{1}{2} & \frac{1}{2} - \frac{1}{2} \\ -\frac{1}{2} + \frac{1}{2} & -\frac{1}{2} + \frac{1}{2} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

∴ Matrix A is nilpotent of order 2.

49 (a)

Let A be a symmetric matrix. Then,

$$A A^{-1} = I$$

$$\Rightarrow (A A^{-1})^T = I$$

$$\Rightarrow (A^{-1})^T A^T = I$$

$$\Rightarrow (A^{-1})^T = (A^T)^{-1}$$

$$\Rightarrow (A^{-1})^T = (A)^{-1} \quad [\because A^T = A]$$

∴ A^{-1} is a symmetric matrix

50 (a)

Since, given matrix is a triangular matrix, so its characteristic roots are the diagonal elements.

Hence, required roots are 1, 3, 6.

51 (d)

$$\text{Given that, } A = \begin{bmatrix} 3 & 2 & 4 \\ 1 & 2 & -1 \\ 0 & 1 & 1 \end{bmatrix}$$

$$\text{and } A^{-1} = \frac{1}{k} \text{adj } A$$

$$\Rightarrow k = |A|$$

$$= \begin{vmatrix} 3 & 2 & 4 \\ 1 & 2 & -1 \\ 0 & 1 & 1 \end{vmatrix}$$

$$= 3(2+1) - 2(1-0) + 4(1-0)$$

$$= 9 - 2 + 4 = 11$$

52 (a)

Since, $A \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ is linear equation in three

variables and that could have only unique, no solution or infinitely many solution.

∴ It is not possible to have two solutions.

Hence, number of matrices A is zero.

53 (c)

Since, A is invertible matrix of order n , then the determinant of $\text{adj } A = |A|^{n-1}$

54 (c)

$$\text{Given, } \text{adj} \begin{bmatrix} 1 & 0 & 2 \\ -1 & 1 & -2 \\ 0 & 2 & 1 \end{bmatrix} = \begin{bmatrix} 5 & a & -2 \\ 1 & 1 & 0 \\ -2 & -2 & b \end{bmatrix} \quad \dots(i)$$

Cofactors of $\begin{bmatrix} 1 & 0 & 2 \\ -1 & 1 & -2 \\ 0 & 2 & 1 \end{bmatrix}$ are

$$C_{11} = 5, C_{12} = 1, C_{13} = -2$$

$$C_{21} = 4, C_{22} = 1, C_{23} = -2$$

$$C_{31} = -2, C_{32} = 0, C_{33} = 1$$

$$\Rightarrow \begin{bmatrix} 5 & 4 & -2 \\ 1 & 1 & 0 \\ -2 & -2 & 1 \end{bmatrix} = \begin{bmatrix} 5 & a & -2 \\ 1 & 1 & 0 \\ -2 & -2 & b \end{bmatrix}$$

On comparing the corresponding elements, we get

$$a = 4, \quad b = 1$$

$$\therefore [a \ b] = [4 \ 1]$$

55 (d)

$$AB = [x \ y \ z] \begin{bmatrix} a & h & g \\ h & b & f \\ g & f & c \end{bmatrix}$$

$$= [xa + yh + zg \quad xh + yb + zf \quad xg + yf + zc]$$

Now, $ABC = O$

$$\Rightarrow [xa + yh + zg \quad xh + yb + zf \quad xg + yf$$

$$+ zc] \begin{bmatrix} x \\ y \\ z \end{bmatrix} = O$$

$$\Rightarrow [ax^2 + hxy + gxz + hxy + y^2b + fzy + gxz + yfz + z^2c]$$

$$= O$$

$$\Rightarrow [ax^2 + by^2 + cz^2 + 2gzx + 2hxy + 2fyz] = O$$

56 (d)

$$\text{Let } A = \begin{bmatrix} 2 & 1 \\ 7 & 4 \end{bmatrix}$$

Let B be the multiplicative inverse of A , then

$$AB = I$$

$$\Rightarrow B = A^{-1}$$

$$= \frac{1}{8-7} \begin{bmatrix} 4 & -1 \\ -7 & 2 \end{bmatrix} = \begin{bmatrix} 4 & -1 \\ -7 & 2 \end{bmatrix}$$

57 (c)

$$\text{Let } A = \begin{bmatrix} 4 & 1 & 0 & 0 \\ 3 & 0 & 1 & 0 \\ 6 & 0 & 2 & 0 \end{bmatrix}$$

Now, we take a submatrix of order 3×3

$$B = \begin{bmatrix} 4 & 1 & 0 \\ 3 & 0 & 1 \\ 6 & 0 & 2 \end{bmatrix}$$

$$|B| = -1(6-6) = 0$$

Now, we take a submatrix of order 2×2 .

$$\text{ie, } C = \begin{bmatrix} 4 & 1 \\ 3 & 0 \end{bmatrix}$$

$$|C| = 0 - 3 \neq 0$$

∴ Rank of matrix A is 2.

58 (d)

$$\text{Let } A = \begin{bmatrix} 1 & 0 & 1 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{2} \begin{bmatrix} 1 & -1 & -1 \\ 1 & 1 & -1 \\ 1 & 1 & 1 \end{bmatrix}$$

Now, $X = A^{-1}B$

$$= \frac{1}{2} \begin{bmatrix} 1 & -1 & -1 \\ 1 & 1 & -1 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$$

$$= \frac{1}{2} \begin{bmatrix} -2 \\ 0 \\ 4 \end{bmatrix} = \begin{bmatrix} -1 \\ 0 \\ 2 \end{bmatrix}$$

59 (a)

$$\begin{bmatrix} 7 & 1 & 2 \\ 9 & 2 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix} + 2 \begin{bmatrix} 4 \\ 2 \end{bmatrix} = \begin{bmatrix} 21 + 4 + 10 \\ 27 + 8 + 5 \end{bmatrix} + \begin{bmatrix} 8 \\ 4 \end{bmatrix} \\ = \begin{bmatrix} 35 \\ 40 \end{bmatrix} + \begin{bmatrix} 8 \\ 4 \end{bmatrix} = \begin{bmatrix} 43 \\ 44 \end{bmatrix}$$

60 (b)

Since, A, B and C are $n \times n$ matrices and, if $A^3 + 2A^2 + 3A + 5I = 0$, then A is invertible.

61 (a)

We have,

$$|A| = -1, |B| = 3$$

$$\therefore |3AB| = 3^3 |AB| = 3^3 |A| |B| = 3 \times -1 \times 3 \\ = -9$$

62 (c)

Let $A = \begin{bmatrix} 1 & 2 & 3 & 0 \\ 2 & 4 & 3 & 2 \\ 3 & 2 & 1 & 3 \\ 6 & 8 & 7 & 5 \end{bmatrix}$. Then,

$$A \sim \begin{bmatrix} 1 & 2 & 3 & 0 \\ 2 & 4 & 3 & 2 \\ 3 & 2 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ By applying } R_4 \rightarrow R_4 - R_3 - R_2 - R_1$$

$$\Rightarrow A \sim \begin{bmatrix} 1 & 2 & 3 & 0 \\ 0 & 0 & -3 & 2 \\ 0 & -4 & -8 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ By applying } R_2 \rightarrow R_2 - 2R_1, \\ R_3 \rightarrow R_3 - 3R_1$$

We observe that the leading minor of the third order of this matrix is non-zero i.e.

$$\begin{vmatrix} 1 & 2 & 3 \\ 0 & 0 & -3 \\ 0 & -4 & -8 \end{vmatrix} = -12 \neq 0. \text{ Hence, rank}(A) = 3$$

63 (a)

Given, $A + I = \begin{bmatrix} 3 & -2 \\ 4 & 1 \end{bmatrix}$

$$\therefore A - I = A + I - 2I = \begin{bmatrix} 3 & -2 \\ 4 & 1 \end{bmatrix} - \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \\ = \begin{bmatrix} 1 & -2 \\ 4 & -1 \end{bmatrix}$$

$$\therefore (A + I)(A - I) = \begin{bmatrix} 3 & -2 \\ 4 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 4 & -1 \end{bmatrix} = \begin{bmatrix} -5 & -4 \\ 8 & -9 \end{bmatrix}$$

64 (d)

We have,

$$A = kI_n \Rightarrow |A| = |kI_n| = k^n |I_n| = k^n$$

65 (d)

Given, $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$

$$A^2 = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} \\ = \begin{bmatrix} a^2 + bc & ab + bd \\ ac + dc & bc + d^2 \end{bmatrix}$$

$$\therefore A^2 - (a + d)A + kI = 0$$

$$\Rightarrow \begin{bmatrix} a^2 + bc & ab + bd \\ ac + dc & bc + d^2 \end{bmatrix} - \begin{bmatrix} a^2 + ad & ab + bd \\ ac + dc & ad + d^2 \end{bmatrix} \\ + \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} bc - ad + k & 0 \\ 0 & bc - ad + k \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

On equating, we get

$$bc - ad + k = 0$$

$$\Rightarrow k = ad - bc \quad \dots(i)$$

Also, $\begin{vmatrix} a & b \\ c & d \end{vmatrix} = 0$

$$\Rightarrow ad - bc = 0$$

\therefore From Eq. (i), $k = 0$

66 (b)

Given that, $A = \begin{bmatrix} 1 & -2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 1 \\ 3 & 2 \\ 1 & 1 \end{bmatrix}$

$$\therefore AB = \begin{bmatrix} 1 & -2 & 1 \\ 2 & 1 & 3 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 3 & 2 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 2 - 6 + 1 & 1 - 4 + 1 \\ 4 + 3 + 3 & 2 + 2 + 3 \end{bmatrix}$$

$$= \begin{bmatrix} -3 & -2 \\ 10 & 7 \end{bmatrix}$$

$$\Rightarrow (AB)^T = \begin{bmatrix} -3 & 10 \\ -2 & 7 \end{bmatrix}$$

68 (b)

We have,

$$A^2 = AA = \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix} \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$$

$$\Rightarrow A^2 = \begin{bmatrix} 29 & -25 \\ -20 & 24 \end{bmatrix}$$

$$\therefore A^2 - 5A = \begin{bmatrix} 29 & -25 \\ -20 & 24 \end{bmatrix} - 5 \begin{bmatrix} 3 & -5 \\ -4 & 2 \end{bmatrix}$$

$$\Rightarrow A^2 - 5A = \begin{bmatrix} 14 & 0 \\ 0 & 14 \end{bmatrix} = 14I$$

71 (b)

Given that A is a singular matrix

$$\therefore |A| = 0$$

$$\therefore A \text{ adj } A = |A| = 0$$

$\therefore A \text{ adj } A$ is a zero matrix

72 (a)

$$\begin{bmatrix} 3 & 1 \\ 4 & 1 \end{bmatrix} X = \begin{bmatrix} 5 & -1 \\ 2 & 3 \end{bmatrix}$$

$$\Rightarrow X = \begin{bmatrix} 3 & 1 \\ 4 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 5 & -1 \\ 2 & 3 \end{bmatrix}$$

$$= \frac{1}{-1} \begin{bmatrix} 1 & -1 \\ -4 & 3 \end{bmatrix} \begin{bmatrix} 5 & -1 \\ 2 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} -1 & 1 \\ 4 & -3 \end{bmatrix} \begin{bmatrix} 5 & -1 \\ 2 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} -3 & 4 \\ 14 & -13 \end{bmatrix}$$

73 (b)

Given, $A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 2 & 1 \\ 2 & 1 & 0 \end{bmatrix}$

$$\therefore A^2 = \begin{bmatrix} 2 & 3 & 1 \\ 5 & 6 & 2 \\ 3 & 4 & 1 \end{bmatrix} \text{ and } A^3 = \begin{bmatrix} 7 & 9 & 3 \\ 15 & 19 & 6 \\ 9 & 12 & 4 \end{bmatrix}$$

$$\text{Hence, } A^3 - 3A^2 - I = 0$$

- 74 **(d)**
 $\because PQ = I \Rightarrow P^{-1} = Q$
 Now, the system in matrix notation is $PX = B$
 $\therefore X = P^{-1}B = QB$
 $\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{9} \begin{bmatrix} 2 & 2 & 1 \\ 13 & -5 & m \\ -8 & 1 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 5 \end{bmatrix}$
 $\Rightarrow y = \frac{1}{9}(13 - 5 + 5m)$
 $\Rightarrow -27 = 8 + 5m$ (given $y = -3$)
 $\therefore m = -7$
- 75 **(d)**
 Since A is invertible. Therefore, $|A| \neq 0$
 Thus, option (d) is correct.
- 76 **(a)**
 We have,
 $\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} A \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
 $\therefore A = \begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix}^{-1} \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix}^{-1} \right)$
 $\Rightarrow A = \begin{bmatrix} 2 & -1 \\ -3 & 2 \end{bmatrix} \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix} \right)$
 $\Rightarrow A = \begin{bmatrix} 2 & -1 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$
- 77 **(c)**
 If A is a square matrix, then $A - A^T$ is a skew-symmetric matrix, then $|A - A^T|$ is '0' or a perfect square as A is of odd order or even order
- 78 **(d)**
 $O(A') = 3 \times 2, O(B') = 2 \times 3$
 (a) $CB + A'$
 Now, order of CB
 $= (\text{order of } C \text{ is } 3 \times 3)(\text{order of } B \text{ is } 3 \times 2)$
 $= \text{order of } CB \text{ is } 3 \times 2$
 Since, $O(A') = 3 \times 2$
 \therefore Matrix $CB + A'$ can be determined.
 (b) $O(BA) = 3 \times 3$
 and $O(C) = 3 \times 3$
 \therefore Matrix BAC can be determined.
 (c) $O(A + B') = 2 \times 3$
 $\Rightarrow O(A + B')' = 3 \times 2$
 and $O(C) = 3 \times 3$
 \therefore Matrix $C(A + B')$ can be determined.
 (d) $O(A + B') = 2 \times 3$
 And $O(C) = 3 \times 3$
 \therefore Matrix $C(A + B')$ cannot be determined.
- 79 **(d)**
 Given A is a matrix of order 3 and $B = |A|A^{-1}, |A| = -5$
 $\therefore B = |A| \frac{(\text{adj } A)}{|A|} \Rightarrow B = (\text{adj } A)$
 $\Rightarrow |B| = |A|^{3-1} = 25$
- 80 **(a)**

$$\because A^2 = \begin{bmatrix} -2 & 4 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} -2 & 4 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

$\therefore A^2$ is null matrix.

81 **(b)**

For unique solution,

$$\begin{vmatrix} k & 2 & -1 \\ 0 & k-1 & -2 \\ 0 & 0 & k+2 \end{vmatrix} \neq 0$$

$$\Rightarrow k(k-1)(k+2) \neq 0$$

$$\Rightarrow k \neq 0, 1 \text{ or } -2$$

82 **(b)**

$$\text{Given, } A = \begin{bmatrix} -1 & 2 \\ 2 & -1 \end{bmatrix} \text{ and } B = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

$$\therefore A^{-1} = -\frac{1}{3} \begin{bmatrix} -1 & -2 \\ -2 & -1 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$$

Also, $AX = B$

$$\Rightarrow X = A^{-1}B = \frac{1}{3} \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

$$= \frac{1}{3} \begin{bmatrix} 3+2 \\ 6+1 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 5 \\ 7 \end{bmatrix}$$

83 **(a)**

$$\Delta = \begin{vmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \\ 5 & 5 & 9 \end{vmatrix}$$

$$= 1(9 - 15) - 2(18 - 15) + 3(10 - 5)$$

$$= -6 - 6 + 15$$

$$= 3 \neq 0$$

Hence, the system of equations has a unique solution.

84 **(b)**

It is given that

$$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

$$\therefore A^2 = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix} = 2A$$

$$\Rightarrow A^3 = 2(AA) = 2A^2 = 2(2A) = 2^2A$$

Continuing in this manner, we have $A^n = 2^{n-1}A$

85 **(a)**

We have,

$$(B^{-1}A^{-1})^{-1} = (A^{-1})^{-1}(B^{-1})^{-1} [\because (PQ)^{-1} = Q^{-1}P^{-1}]$$

$$\Rightarrow (B^{-1}A^{-1})^{-1} = AB$$

$$\Rightarrow (B^{-1}A^{-1}) = \begin{bmatrix} 2 & 2 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

$$\Rightarrow (B^{-1}A^{-1})^{-1} = \begin{bmatrix} 2 & -2 \\ 2 & 3 \end{bmatrix}$$

Hence, option (a) is correct

86 **(c)**

Here, C, A and C^T are matrix of order $n \times 1, n \times n$ and $1 \times n$ respectively.

Let $C^TAC = k$

$$\text{Then, } (C^TAC)^T = C^T A^T (C^T)^T$$

$$= C^T A^T C = C^T (-A)C$$

$$= -C^TAC = -k$$

$$\Rightarrow k = -k \Rightarrow k = 0$$

$\Rightarrow C^T AC$ is null matrix.

Which shows that $C^T AC$ is a zero matrix of order 1.

87 (b)

Since, $A + A^T$ is a square matrix

$$\therefore (A + A^T)^T = A^T + (A^T)^T = A^T + A$$

Hence, $A + A^T$ is symmetric matrix

88 (d)

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} 1+1 & 1+2 \\ 2+1 & 2+2 \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 3 & 4 \end{bmatrix}$$

89 (a)

$$\text{Given, } P = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{bmatrix} \begin{bmatrix} -1 & -2 \\ -2 & 0 \\ 0 & -4 \end{bmatrix} \begin{bmatrix} -4 & -5 & -6 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 5 \end{bmatrix} \begin{bmatrix} 4 & 5 & 4 \\ 8 & 10 & 12 \\ 0 & 0 & -4 \end{bmatrix}$$

$$\therefore P_{22} = \begin{bmatrix} 2 & 3 & 4 \\ 10 \end{bmatrix}$$

$$= 10 + 30 = 40$$

91 (c)

Given, $AB = A \Rightarrow B = I$

And $BA = B \Rightarrow A = I$

$\therefore B = I$ and $A = I$

$\Rightarrow B^2 = B$ and $A^2 = A$

92 (a)

Let $A = \begin{bmatrix} 0 & x & y & z \\ -x & 0 & a & b \\ -y & -a & 0 & c \\ -z & -b & -c & 0 \end{bmatrix}$ be a skew-symmetric

matrix. Then,

$$|A| = \begin{vmatrix} 0 & x & y & z \\ -x & 0 & a & b \\ -y & -a & 0 & c \\ -z & -b & -c & 0 \end{vmatrix} = (cx - by + az)^2$$

93 (d)

$$\text{Since, } A = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 2 & -3 \\ 2 & 1 & 0 \end{bmatrix}$$

$$\therefore B = \text{adj } A = \begin{bmatrix} 3 & 1 & 1 \\ -6 & -2 & 3 \\ -4 & -3 & 2 \end{bmatrix}$$

$$\Rightarrow \text{adj } B = \begin{bmatrix} 5 & -5 & 5 \\ 0 & 10 & -15 \\ 10 & 5 & 0 \end{bmatrix}$$

$$\Rightarrow |\text{adj } B| = \begin{vmatrix} 5 & -5 & 5 \\ 0 & 10 & -15 \\ 10 & 5 & 0 \end{vmatrix} = 625$$

Given that, $C = 5A$

$$\Rightarrow |C| = 5^3 |A| = 125 \begin{vmatrix} 1 & -1 & 1 \\ 0 & 2 & -3 \\ 2 & 1 & 2 \end{vmatrix} = 625$$

$$\text{Hence, } \frac{|\text{adj}(B)|}{|C|} = \frac{625}{625} = 1$$

94 (a)

$$\text{Given, } A = \begin{bmatrix} 6 & 8 & 5 \\ 4 & 2 & 3 \\ 9 & 7 & 1 \end{bmatrix}$$

and symmetric matrix $B = \frac{A+A'}{2}$

$$\therefore B = \frac{1}{2} \left\{ \begin{bmatrix} 6 & 8 & 5 \\ 4 & 2 & 3 \\ 9 & 7 & 1 \end{bmatrix} + \begin{bmatrix} 6 & 4 & 9 \\ 8 & 2 & 7 \\ 5 & 3 & 1 \end{bmatrix} \right\} = \begin{bmatrix} 6 & 6 & 7 \\ 6 & 2 & 5 \\ 7 & 5 & 1 \end{bmatrix}$$

95 (d)

The matrix $\begin{bmatrix} 5 & 10 & 3 \\ -2 & -4 & 6 \\ -1 & -2 & b \end{bmatrix}$ is singular, if

$$\begin{vmatrix} 5 & 10 & 3 \\ -2 & -4 & 6 \\ -2 & -2 & b \end{vmatrix} = 0$$

$$\Rightarrow -1(60 + 12) + 2(30 + 6) + b(-20 + 20) = 0$$

$$\Rightarrow -72 + 72 + 0b = 0$$

Hence, the given matrix is singular for any value of b

96 (b)

$$\det(2AB) = 2^3 \det(A) \det(B) \\ = 8 \det(A) \det(B)$$

97 (b)

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj} \text{ (In general)}$$

And in a diagonal matrix non-diagonal elements are zero i.e.,

$$a_{ij} = \begin{cases} 0, & \text{if } i \neq j \\ a_{ij}, & \text{if } i = j \end{cases}$$

So, $c_{ij} = a_{ii} b_{ij}$

98 (b)

$$\text{Here, } |A| = (1)(9) - 2(-11) - 3(6) \\ = 9 + 22 - 18 = 33$$

Since, $A^{-1} \text{adj}(A^{-1}) = |A^{-1}| I_3$

$$\Rightarrow A^{-1} \text{adj}(A^{-1}) = (|A|^{-1}) I_3$$

$$\Rightarrow A \cdot A^{-1} \text{adj}(A^{-1}) = (|A|)^{-1} A I_3$$

$$\Rightarrow \text{adj}(A^{-1}) = (|A|)^{-1} A$$

$$\Rightarrow |A| \text{adj}(A^{-1}) = A \quad (\text{But } |A| \neq 0)$$

99 (d)

$$\text{Let } A = \begin{bmatrix} 1 & -1 & 1 \\ 1 & 1 & -1 \\ -1 & 1 & 1 \end{bmatrix}$$

$$\therefore |A| = 1(1+1) + 1(1-1) + 1(1+1) = 4 \neq 0$$

\therefore Rank of matrix A is 3

100 (b)

Let $\frac{x^2}{a^2} = X$, $\frac{y^2}{b^2} = Y$ and $\frac{z^2}{c^2} = Z$, then given equation will be

$$X + Y - Z = 1, X - Y + Z = 1, -X + Y + Z = 1$$

$$\text{Here, } A = \begin{bmatrix} 1 & 1 & -1 \\ 1 & -1 & 1 \\ -1 & 1 & 1 \end{bmatrix}$$

$$\text{Now, } |A| = -4 \neq 0$$

Therefore, the given system of equation has

unique solution.

102 (b)

$$\begin{aligned} \left[\frac{1}{2}(A - A') \right]' &= \frac{1}{2}(A - A')' = \frac{1}{2}(A' - A) \\ &= -\frac{1}{2}(A - A') \end{aligned}$$

Hence, it is a skew-symmetric matrix.

103 (b)

$$\begin{aligned} \therefore \text{adj } A &= \begin{bmatrix} 4 & 2 \\ -3 & 1 \end{bmatrix} \\ \text{and } |A| &= \begin{vmatrix} 1 & -2 \\ 3 & 4 \end{vmatrix} = 10 \\ \therefore A^{-1} &= \frac{1}{10} \begin{bmatrix} 4 & 2 \\ -3 & 1 \end{bmatrix} \end{aligned}$$

105 (b)

$$\begin{aligned} \text{Let } A &= \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix} \\ \therefore \text{adj } A &= \begin{bmatrix} 1 & -2 & -2 \\ -1 & 3 & 3 \\ 0 & -4 & -3 \end{bmatrix}^T \\ &= \begin{bmatrix} 1 & -1 & 0 \\ -2 & 3 & -4 \\ -2 & 3 & -3 \end{bmatrix} \end{aligned}$$

106 (d)

We have,

$$\begin{aligned} 3A^3 + 2A^2 + 5A + I &= 0 \\ \Rightarrow I &= -3A^3 - 2A^2 - 5A \\ \Rightarrow IA^{-1} &= (-3A^3 - 2A^2 - 5A)A^{-1} \\ \Rightarrow A^{-1} &= -3A^2 - 2A - 5I \end{aligned}$$

108 (b)

The given system of equations are

$$\begin{aligned} x + y + z &= 0, \\ 2x + 3y + z &= 0 \text{ and } x + 2y = 0 \end{aligned}$$

$$\begin{aligned} \text{Here, } \begin{vmatrix} 1 & 1 & 1 \\ 2 & 3 & 1 \\ 1 & 2 & 0 \end{vmatrix} &= 1(0 - 2) - 1(0 - 1) + (4 - 3) \\ &= -2 + 1 + 1 = 0 \end{aligned}$$

\therefore This system has infinite solutions

109 (c)

$$\begin{aligned} 2X - \begin{bmatrix} 1 & 2 \\ 7 & 4 \end{bmatrix} &= \begin{bmatrix} 3 & 2 \\ 0 & -2 \end{bmatrix} \\ \Rightarrow 2X &= \begin{bmatrix} 3 & 2 \\ 0 & -2 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 7 & 4 \end{bmatrix} \\ \Rightarrow 2X &= \begin{bmatrix} 4 & 4 \\ 7 & 2 \end{bmatrix} \\ \Rightarrow X &= \begin{bmatrix} 2 & 2 \\ 7/2 & 1 \end{bmatrix} \end{aligned}$$

111 (b)

Given, $A = A', B = B'$

$$\begin{aligned} \text{Now, } (AB - BA)' &= (AB)' - (BA)' \\ &= B'A' - A'B' \\ &= BA - AB \\ &= -(AB - BA) \\ \therefore AB - BA &\text{ is a skew-symmetric matrix.} \end{aligned}$$

113 (a)

Given that, p is a non-singular matrix such that

$$\begin{aligned} 1 + p + p^2 + \dots + p^n &= 0 \\ \Rightarrow (1 + p)(1 + p + p^2 + \dots + p^n) &= 0 \\ \Rightarrow 1 - p^{n+1} &= 0 \\ \Rightarrow p^{n+1} &= 1 \\ \Rightarrow p^n \times p^1 &= 1 \\ \Rightarrow p^n &= 1/p \\ \therefore p^{-1} &= p^n \end{aligned}$$

114 (a)

$$\begin{aligned} \text{Given, } \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \frac{1}{40} \begin{bmatrix} 5 & 10 & -5 \\ -5 & -2 & 13 \\ 10 & -4 & 6 \end{bmatrix} \begin{bmatrix} 5 \\ 0 \\ 5 \end{bmatrix} \\ &= \frac{1}{40} \begin{bmatrix} 25 + 0 - 25 \\ -25 + 0 + 65 \\ 50 + 0 + 30 \end{bmatrix} \\ &= \frac{1}{40} \begin{bmatrix} 0 \\ 40 \\ 80 \end{bmatrix} \\ \Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} \\ \Rightarrow x = 0, y = 1, z = 2 \\ \therefore x + y + z &= 0 + 1 + 2 = 3 \end{aligned}$$

115 (d)

$$\begin{aligned} \therefore \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} &= \begin{bmatrix} y \\ x \end{bmatrix} = \begin{bmatrix} X \\ Y \end{bmatrix} \\ \text{Then, } X &= y \text{ and } Y = x \\ \text{ie, } y &= x \end{aligned}$$

116 (c)

$$\begin{aligned} \text{Given } \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} \begin{bmatrix} 1 & \tan \theta \\ -\tan \theta & 1 \end{bmatrix}^{-1} &= \\ \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \\ \Rightarrow \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} \cdot \frac{1}{1 + \tan^2 \theta} \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} &= \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \\ \Rightarrow \frac{1}{1 + \tan^2 \theta} \begin{bmatrix} 1 - \tan^2 \theta & -2 \tan \theta \\ 2 \tan \theta & 1 - \tan^2 \theta \end{bmatrix} &= \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \\ \Rightarrow \begin{bmatrix} 1 - \tan^2 \theta & 2 \tan \theta \\ 1 + \tan^2 \theta & 1 + \tan^2 \theta \end{bmatrix} &= \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \\ \Rightarrow \begin{bmatrix} \cos 2\theta - \sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix} &= \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \\ \Rightarrow a = \cos 2\theta, b = \sin 2\theta \end{aligned}$$

117 (d)

$$\begin{aligned} \text{Given, } A &= \begin{bmatrix} -1 & -2 & -2 \\ 2 & 1 & -2 \\ 2 & -2 & 1 \end{bmatrix} \\ \therefore B &= \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix} = \begin{bmatrix} -3 & -6 & -6 \\ 6 & 3 & -6 \\ 6 & -6 & 3 \end{bmatrix} \end{aligned}$$

$$\Rightarrow \text{adj } A = (B)' = \begin{bmatrix} -3 & 6 & 6 \\ -6 & 3 & -6 \\ -6 & -6 & 3 \end{bmatrix}$$

$$= 3 \begin{bmatrix} -1 & 2 & 2 \\ -2 & 1 & -2 \\ -2 & -2 & 1 \end{bmatrix} = 3A'$$

118 (d)

Given equations are

$$px + y + z = 0, x + qy + z = 0, x + y + rz = 0$$

Since, the system have a non-zero solution, then

$$\begin{bmatrix} p & 1 & 1 \\ 1 & q & 1 \\ 1 & 1 & r \end{bmatrix} = 0$$

Applying $C_2 \rightarrow C_2 - C_1$ and $C_3 \rightarrow C_3 - C_2$

$$\Rightarrow \begin{bmatrix} p & 1-p & 0 \\ 1 & q-1 & 1-q \\ 1 & 0 & r-1 \end{bmatrix} = 0$$

$$\Rightarrow (1-p)(1-q)(1-r) \begin{vmatrix} \frac{p}{1-p} & 1 & 0 \\ 1 & -1 & 1 \\ \frac{1}{1-r} & 0 & -1 \end{vmatrix} = 0$$

$$\Rightarrow (1-p)(1-q)(1-r)$$

$$\left[\frac{p}{1-p}(1) - 1 \left(-\frac{1}{1-q} - \frac{1}{1-r} \right) \right] = 0$$

Since, $p, q, r \neq 1$

$$\therefore \frac{p}{1-p} + \frac{1}{1-q} + \frac{1}{1-r} = 0$$

$$\Rightarrow \frac{1}{1-p} - 1 + \frac{1}{1-q} + \frac{1}{1-r} = 0$$

$$\Rightarrow \frac{1}{1-p} + \frac{1}{1-q} + \frac{1}{1-r} = 0$$

119 (b)

Given, $AB = I \Rightarrow B = A^{-1}$

Now, $A^{-1} = \frac{\text{adj } A}{|A|}$

$$= \frac{\begin{bmatrix} 1 & -\tan \frac{\theta}{2} \\ \tan \frac{\theta}{2} & 1 \end{bmatrix}}{1 + \tan^2 \frac{\theta}{2}}$$

$$= \frac{A^T}{\sec^2 \frac{\theta}{2}} = \cos^2 \frac{\theta}{2} A^T$$

120 (d)

Given equations are

$$3x + y + 2z = 3 \quad \dots(i)$$

$$2x - 3y - z = -3 \quad \dots(ii)$$

$$\text{and } x + 2y + z = 4 \quad \dots(iii)$$

$$\text{Let } A = \begin{bmatrix} 3 & 1 & 2 \\ 2 & -3 & -1 \\ 1 & 2 & 1 \end{bmatrix}, B = \begin{bmatrix} 3 \\ -3 \\ 4 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$\therefore |A| = \begin{vmatrix} 3 & 1 & 2 \\ 2 & -3 & -1 \\ 1 & 2 & 1 \end{vmatrix}$$

$$= 3 \begin{vmatrix} -3 & -1 \\ 2 & 1 \end{vmatrix} - 1 \begin{vmatrix} 2 & -1 \\ 1 & 1 \end{vmatrix} + 2 \begin{vmatrix} 2 & -3 \\ 1 & 2 \end{vmatrix}$$

$$= 3(-3+2) - 1(2+1) + 2(4+3)$$

$$= -3 - 3 + 14 = 8$$

$$\text{adj. } A = \begin{bmatrix} -1 & -3 & 7 \\ 3 & 1 & -5 \\ 5 & 7 & -11 \end{bmatrix}^T$$

$$= \begin{bmatrix} -1 & 3 & 5 \\ -3 & 1 & 7 \\ 7 & -5 & -11 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{8} \begin{bmatrix} -1 & 3 & 5 \\ -3 & 1 & 7 \\ 7 & -5 & -11 \end{bmatrix}$$

Now, $X = A^{-1}B$

$$= \frac{1}{8} \begin{bmatrix} -1 & 3 & 5 \\ -3 & 1 & 7 \\ 7 & -5 & -11 \end{bmatrix} \begin{bmatrix} 3 \\ -3 \\ 4 \end{bmatrix}$$

$$= \frac{1}{8} \begin{bmatrix} -3 - 9 + 20 \\ -9 - 3 + 28 \\ 21 + 15 - 44 \end{bmatrix} = \frac{1}{8} \begin{bmatrix} 8 \\ 16 \\ -8 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

$$\Rightarrow x = 1, y = 2, z = -1$$

121 (a)

$$A^2 = 2A - I$$

$$\therefore A^2 A = 2AA - IA$$

$$= 2A^2 - A = 2(2A - I) - A$$

$$\Rightarrow A^3 = 3A - 2I$$

$$\Rightarrow A^3 \cdot A = 3AA - 2IA = 3(2A - I) - 2A$$

$$\Rightarrow A^4 = 4A - 3I$$

Similarly, $A^n = nA - (n-1)I$

122 (d)

$$\det(M_r) = \begin{vmatrix} r & r-1 \\ r-1 & r \end{vmatrix} = 2r - 1$$

$$\sum_{r=1}^{2007} \det(M_r) = 2 \sum_{r=1}^{2007} r - 2007$$

$$= 2 \times \frac{2007 \times 2008}{2} - 2007 = (2007)^2$$

123 (a)

$$\text{Let } A = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 2 \\ 1 & -2 & 0 \end{bmatrix}$$

$$\therefore |A| = \begin{vmatrix} 0 & 1 & -1 \\ -1 & 0 & 2 \\ 1 & -2 & 0 \end{vmatrix}$$

$$= 0 \begin{vmatrix} 0 & 2 \\ -2 & 0 \end{vmatrix} - 1 \begin{vmatrix} -1 & 2 \\ 1 & 0 \end{vmatrix} - 1 \begin{vmatrix} -1 & 0 \\ 1 & -2 \end{vmatrix}$$

$$= 0 + 2 - 2 = 0$$

$$\Rightarrow |A| = 0$$

$$\text{Now, } (\text{adj } A)B = \begin{bmatrix} 4 & 2 & 2 \\ 2 & 1 & 1 \\ 2 & 1 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}$$

$$= \begin{bmatrix} 4 - 4 + 6 \\ 2 - 2 + 3 \\ 2 - 2 - 3 \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \\ -3 \end{bmatrix} \neq 0$$

∴ This system of equation is inconsistent, so it has no solution

125 (c)

Given, $D = \text{diag}(d_1, d_2, d_3, \dots, d_n)$

$$\Rightarrow D^{-1} = \text{diag}(d_1^{-1}, d_2^{-1}, \dots, d_n^{-1})$$

126 (a)

We have,

$$a = \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow A^n = \begin{bmatrix} 1 & na \\ 0 & 1 \end{bmatrix} \quad [\text{Using PMI}]$$

$$\Rightarrow \frac{1}{n}A^n = \begin{bmatrix} \frac{1}{n} & a \\ 0 & \frac{1}{n} \end{bmatrix} \Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n}A^n = \begin{bmatrix} 0 & a \\ 0 & 0 \end{bmatrix}$$

127 (d)

The given system of equations are

$$2x + y - 5 = 0 \quad \dots(\text{i})$$

$$x - 2y + 1 = 0 \quad \dots(\text{ii})$$

$$\text{and } 2x - 14y - a = 0 \quad \dots(\text{iii})$$

This system is consistent.

$$\therefore \begin{bmatrix} 2 & 1 & -5 \\ 1 & -2 & 1 \\ 2 & -14 & -a \end{bmatrix} = 0$$

$$\Rightarrow 2(2a + 14) - 1(-a - 2) - 5(-14 + 4) = 0$$

$$\Rightarrow 4a + 28 + a + 2 + 50 = 0$$

$$\Rightarrow 5a = -80 \Rightarrow a = -16$$

128 (d)

The system of given equations has no solution,

$$\text{if } \begin{bmatrix} \alpha & 1 & 1 \\ 1 & \alpha & 1 \\ 1 & 1 & \alpha \end{bmatrix} = 0$$

Applying $C_1 \rightarrow C_1 + C_2 + C_3$ and taking common $(\alpha + 2)$ from C_1 , we get

$$(\alpha + 2) \begin{bmatrix} 1 & 1 & 1 \\ 1 & \alpha & 1 \\ 1 & 1 & \alpha \end{bmatrix} = 0$$

Applying $R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1$

$$\Rightarrow (\alpha + 2) \begin{bmatrix} 1 & 1 & 1 \\ 0 & \alpha - 1 & 0 \\ 0 & 0 & \alpha - 1 \end{bmatrix} = 0$$

$$\Rightarrow (\alpha + 2)(\alpha - 1)^2 = 0$$

$$\Rightarrow \alpha = 1, -2$$

But $\alpha = 1$ makes given three equations same. So, the system of equation have infinite solution. So, answer is $\alpha = -2$ for which the system of equations has no solution

130 (b)

$$\text{Given, } A = \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix}$$

$$\therefore A^2 = \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x^2 + 1 & x \\ x & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow x^2 + 1 = 1, x = 0$$

$$\Rightarrow x = 0$$

131 (a)

Given that, $A^{-1} = \lambda (\text{adj } A)$

On comparing with $A^{-1} = \frac{1}{|A|} A$ we get

$$\lambda = \frac{1}{|A|}$$

$$\text{Now, } |A| = \begin{vmatrix} 0 & 3 \\ 2 & 0 \end{vmatrix} = 0 - 6 = -6$$

$$\Rightarrow \lambda = -\frac{1}{6}$$

132 (d)

$$a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13} + a_{14}C_{14} = |A|$$

133 (d)

Given equation are $x + y + z = 6, x + 2y + 3z = 10$ and $x + 2y + \lambda z = 10$

Since, it is consistent.

$$\therefore \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \\ 1 & 2 & \lambda \end{vmatrix} = 0$$

$$\Rightarrow 1(2\lambda - 6) - 1(\lambda - 3) + 1(2 - 2) = 0$$

$$\Rightarrow \lambda - 3 = 0 \Rightarrow \lambda = 3$$

134 (b)

$$A^2 = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = 2 \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} = 2A$$

$$\therefore A^4 = 2A \cdot 2A = 4A^2 = 4 \times 2A = 2^3 A$$

Similarly, $A^8 = 2^7 A$

$$\Rightarrow A^{100} = 2^{99} A$$

135 (d)

$$\text{Let } A = \begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$$

$$\therefore |A| = \cos^2 2\theta + \sin^2 2\theta = 1$$

$$\text{and } \text{adj } A = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{1} \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$$

$$= \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$$

138 (a)

Give equation can be written as,

$$2X = \begin{bmatrix} 3 & 8 \\ 7 & 2 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

$$\Rightarrow 2X = \begin{bmatrix} 2 & 6 \\ 4 & -2 \end{bmatrix} = 2 \begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix}$$

$$\Rightarrow X = \begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix}$$

139 (c)

We have,

$$A B = 0$$

$$\Rightarrow |A B| = 0$$

$$\Rightarrow |A| |B| = 0$$

$$\Rightarrow |A| = 0 \text{ or } |B| = 0$$

Let $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, B = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$. Then, $A B = O$. But

$$A \neq O, B \neq O$$

140 (b)

$$\text{Given, } A = \begin{bmatrix} 1 & -1 & -2 \\ 2 & 1 & 1 \\ 4 & -1 & -2 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{3} \begin{bmatrix} -1 & 0 & 1 \\ 8 & 6 & -5 \\ -6 & -3 & 3 \end{bmatrix}$$

$$\text{Now, } A^{-1}D = \frac{1}{3} \begin{bmatrix} -1 & 0 & 1 \\ 8 & 6 & -5 \\ -6 & -3 & 3 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \\ 11 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 8 \\ -1 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 8/3 \\ -1/3 \\ 0 \end{bmatrix}$$

142 (a)

Given equations are $x - cy - bz = 0$
 $cx - y + az = 0$ and $bx + ay - z = 0$

For non-zero solution

$$\begin{vmatrix} 1 & -c & -b \\ c & -1 & a \\ b & a & -1 \end{vmatrix} = 0$$

$$\Rightarrow 1(1 - a^2) + c(-c - ab) - b(ac + b) = 0$$

$$\Rightarrow a^2 + b^2 + c^2 + 2abc = 1$$

143 (b)

We have,

$$\text{Det}(I_n) = 1 (\neq 0) \Rightarrow \text{rank}(I_n) = n$$

144 (a)

The given matrix A is singular, if

$$|A| = \begin{vmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & \lambda \end{vmatrix} = 0$$

$$\Rightarrow 8(7\lambda - 16) + 6(-6\lambda + 8) + 2(24 - 14) = 0$$

$$\Rightarrow 56\lambda - 128 - 36\lambda + 48 + 20 = 0$$

$$\Rightarrow 20\lambda = 60$$

$$\Rightarrow \lambda = 3$$

145 (c)

$$\text{Let } B = I + A + A^2 + A^3 \dots \infty$$

$$\Rightarrow AB = A + A^2 + A^3 + \dots \infty$$

$$\Rightarrow B - AB = I$$

$$\Rightarrow B(I - A) = I$$

$$\Rightarrow B = (I - A)^{-1}$$

$$\Rightarrow B = \begin{bmatrix} 0 & -2 \\ -3 & -3 \end{bmatrix}^{-1} = -\frac{1}{6} \begin{bmatrix} -3 & 2 \\ 3 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1/2 & -1/3 \\ -1/2 & 0 \end{bmatrix}$$

146 (a)

Since A is non-singular matrix

$$\therefore |A| \neq 0 \Rightarrow \text{rank}(A) = n$$

147 (b)

$$A^2 = \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} = \begin{bmatrix} -7 & -12 \\ 24 & 17 \end{bmatrix}$$

$$\text{Now, } f(A) = A^2 - 3A + 7$$

$$= \begin{bmatrix} -7 & -12 \\ 24 & 17 \end{bmatrix} - 3 \begin{bmatrix} 1 & -2 \\ 4 & 5 \end{bmatrix} + 7 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} -3 & -6 \\ 12 & 9 \end{bmatrix}$$

$$\therefore f(A) + \begin{bmatrix} 3 & 6 \\ -12 & -9 \end{bmatrix} = \begin{bmatrix} -3 & -6 \\ 12 & 9 \end{bmatrix} + \begin{bmatrix} 3 & 6 \\ -12 & -9 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

148 (a)

The given system of equations will have a unique solution, if

$$\begin{vmatrix} 1 & 1 & 1 \\ 2 & 1 & -1 \\ 3 & 2 & k \end{vmatrix} \neq 0 \Rightarrow k \neq 0$$

149 (a)

$$\text{Given, } 2x + y - z = 7 \quad \dots(i)$$

$$x - 3y + 2z = 1 \quad \dots(ii)$$

$$\text{and } x + 4y - 3z = 5 \quad \dots(iii)$$

From Eqs.(i) and (ii), we get

$$5x - y = 15 \quad \dots(iv)$$

From Eqs. (i) and (iii)

$$5x - y = 16 \quad \dots(v)$$

Eqs. (iv) and (v) shows that they are parallel and solution does not exist.

150 (d)

We have,

$$X = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix} \Rightarrow X^2 = \begin{bmatrix} 5 & -8 \\ 2 & -3 \end{bmatrix}$$

Clearly for $n = 2$, the matrices in options (a), (b),

$$(c) \text{ do not tally with } \begin{bmatrix} 5 & -8 \\ 2 & -3 \end{bmatrix}$$

152 (a)

We have,

$$A = [a_{ij}] \therefore |kA| = k^n |A|$$

153 (b)

$$A^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & -1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

154 (d)

$$\text{Given that, } A = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix}$$

$$\Rightarrow A^2 = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} \alpha^2 + 0 & 0 + 0 \\ \alpha + 1 & 0 + 1 \end{bmatrix}$$

$$= \begin{bmatrix} \alpha^2 & 0 \\ \alpha + 1 & 1 \end{bmatrix}$$

Also, $B = A^2$ (given)

$$\Rightarrow \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix} = \begin{bmatrix} \alpha^2 & 0 \\ \alpha + 1 & 1 \end{bmatrix}$$

Clearly this is not satisfied by any real value of α

155 (b)

We have,

$$A(\text{adj } A) = \begin{bmatrix} 4 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 4 \end{bmatrix}$$

$$\Rightarrow |A| I = 4I \quad [\because A(\text{adj } A) = |A| I]$$

$$\Rightarrow |A| = 4$$

$$\Rightarrow |\text{adj } A| = |A|^2 \quad [|\text{adj } A| = |A|^{n-1}]$$

156 (a)

Given, $A = \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix}$

$$A^2 = \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix} \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix} = \begin{bmatrix} \omega^2 & 0 \\ 0 & \omega^2 \end{bmatrix}$$

$$\Rightarrow A^3 = \begin{bmatrix} \omega^2 & 0 \\ 0 & \omega^2 \end{bmatrix} \begin{bmatrix} \omega & 0 \\ 0 & \omega \end{bmatrix} = \begin{bmatrix} \omega^3 & 0 \\ 0 & \omega^3 \end{bmatrix}$$

Similarly, $A^{50} = \begin{bmatrix} \omega^{50} & 0 \\ 0 & \omega^{50} \end{bmatrix}$

$$= \begin{bmatrix} (\omega^2)^{16} \omega^2 & 0 \\ 0 & (\omega^3)^{16} \omega^2 \end{bmatrix}$$

$$= \begin{bmatrix} \omega^2 & 0 \\ 0 & \omega^2 \end{bmatrix}$$

$$= \omega^2 A$$

157 (a)

We have,

$$AB = I_3$$

$$\Rightarrow \begin{bmatrix} 1 & 2 & x \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & -2 & y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & 0 & x+y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow x + y = 0$$

158 (a)

Let $A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$

$$\therefore \text{adj}(A) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

159 (a)

$$\because |A| = 0 - 1(1 - 9) + 2(1 - 6)$$

$$= 8 - 10 = -2 \neq 0$$

$$\text{adj}(A) = \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{-2} \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix} = \begin{bmatrix} 1/2 & -1/2 & 1/2 \\ -4 & 3 & -1 \\ 5/2 & -3/2 & 1/2 \end{bmatrix}$$

160 (a)

$$|f(\theta)| = 1(\cos^2 \theta + \sin^2 \theta) = 1$$

Now, $\text{adj}\{f(\theta)\} = \begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$

$$\therefore \{f(\theta)\}^{-1} = \begin{bmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix} = f(-\theta)$$

161 (a)

Given, $x + 4ay + az = 0$... (i)

$x + 3by + bz = 0$... (ii)

And $x + 2cy + cz = 0$... (iii)

For non-trivial solution

$$\begin{vmatrix} 1 & 4a & a \\ 1 & 3b & b \\ 1 & 2c & c \end{vmatrix} = 0$$

Applying $R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1$

$$\Rightarrow \begin{vmatrix} 1 & 4a & a \\ 0 & 3b - 4a & b - a \\ 0 & 2c - 4a & c - a \end{vmatrix} = 0$$

$$\Rightarrow 1[(3b - 4a)(c - a) - 2(b - a)(c - 2a)] = 0$$

$$\Rightarrow 3bc - 3ab - 4ac + 4a^2$$

$$- 2(bc - 2ab - ac + 2a^2) = 0$$

$$\Rightarrow bc + ab - 2ac = 0$$

$$\Rightarrow ab + bc = 2ac$$

162 (d)

We know that

$$\text{rank}(AB) \leq \text{rank}(A)$$

$$\text{and, rank}(AB) \leq \text{rank}(B)$$

$$\therefore \text{rank}(AB) \leq \min(\text{rank } A, \text{rank } B)$$

163 (c)

Let $A = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}$ and $B = [b_{11} \ b_{12} \ b_{13} \ \dots \ b_{1n}]$ be two

non-zero column and row matrices respectively

We have,

$$AB = \begin{bmatrix} a_{11} b_{11} & a_{11} b_{12} & a_{11} b_{13} & \dots & a_{11} b_{1n} \\ a_{21} b_{11} & a_{21} b_{12} & a_{21} b_{13} & \dots & a_{21} b_{1n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{m1} b_{11} & a_{m1} b_{12} & a_{m1} b_{13} & \dots & a_{m1} b_{1n} \end{bmatrix}$$

Since A and B are non-zero matrices. Therefore, the matrix AB will also be a non-zero matrix. The matrix AB will have at least one non-zero element obtained by multiplying corresponding non-zero elements of A and B . All the two-rowed minors of A obviously vanish. But, A is a non-zero matrix.

Hence, $\text{rank}(A) = 1$

164 (c)

$$\begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix} A \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$A = \begin{bmatrix} 2 & 1 \\ 3 & 2 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix}^{-1}$$

$$= - \begin{bmatrix} 2 & -1 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -3 & -2 \\ -5 & -3 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & -1 \\ -3 & 2 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

165 (a)

If A is any square matrix, then

$$AA^{-1} = I \text{ and } A^{-1}A = A^{-1}$$

$$\text{Since, } A^2 - A + I = 0$$

$$\Rightarrow A^{-1}A^2 - A^{-1}A + A^{-1}I = 0$$

$$\Rightarrow (A^{-1}A)A - (A^{-1}A) + A^{-1} = 0$$

$$\Rightarrow A - 1 + A^{-1} = 0 \Rightarrow A^{-1} = I - A$$

166 (a)

Since, B is invertible, therefore B^{-1} exists

$$\text{Now, rank}(A) = \text{rank}[(AB)B^{-1}] \leq \text{rank}(AB)$$

But $\text{rank}(AB) \leq \text{rank}(A)$

$$\therefore \text{rank}(AB) = \text{rank}(A)$$

167 (c)

Given, $A = \begin{bmatrix} 4 & 2 \\ 3 & 4 \end{bmatrix}$ of order $n = 2$

$$\therefore |\text{adj}(A)| = |A|^{2-1} = \begin{vmatrix} 4 & 2 \\ 3 & 4 \end{vmatrix} = 10$$

168 (d)

$$\begin{aligned} & \cos \theta \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} + \sin \theta \begin{bmatrix} \sin \theta & -\cos \theta \\ \cos \theta & \sin \theta \end{bmatrix} \\ &= \begin{bmatrix} \cos^2 \theta + \sin^2 \theta & 0 \\ 0 & \cos^2 \theta + \sin^2 \theta \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

169 (b)

Let A denote the matrix every element of which is unity. Then, all the 2-rowed minors of A obviously vanish. But A is a non-null matrix. Hence, rank of A is 1

170 (d)

As $\det(A) = \pm 1, A^{-1}$ exists

$$\text{and } A^{-1} = \frac{1}{\det(A)} (\text{adj } A) = \pm (\text{adj } A)$$

All entries in $\text{adj}(A)$ are integers.

$\therefore A^{-1}$ has integer entries.

171 (c)

Since, A is invertible

$$\therefore |A| \neq 0 \Rightarrow \begin{vmatrix} 1 & 0 & -k \\ 2 & 1 & 3 \\ k & 0 & 1 \end{vmatrix} \neq 0$$

$$\Rightarrow 1(1-0) + k(0-k) \neq 0$$

$$\Rightarrow 1 - k^2 \neq 0 \Rightarrow k \neq \pm 1$$

172 (b)

We have,

$$\begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} \begin{bmatrix} 1 & \tan \theta \\ -\tan \theta & 1 \end{bmatrix}^{-1} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} \frac{1}{1 + \tan^2 \theta} \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

$$\Rightarrow \frac{1}{1 + \tan^2 \theta} \begin{bmatrix} 1 - \tan^2 \theta & -2 \tan \theta \\ 2 \tan \theta & 1 - \tan^2 \theta \end{bmatrix} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} & \frac{-2 \tan \theta}{1 + \tan^2 \theta} \\ \frac{2 \tan \theta}{1 + \tan^2 \theta} & \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \end{bmatrix} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

$$\Rightarrow a = \cos 2\theta, b = \sin 2\theta$$

173 (c)

We have,

$$x^2 + y^2 + z^2 \neq 0$$

\Rightarrow At least one of x, y, z is non-zero

Now,

$$x = cy + bz, y = az + cx, z = bx + ay$$

$$\Rightarrow x - cy - bz = 0$$

$$cx - y + az = 0$$

$$bx + zy - z = 0$$

As at least one of x, y, z is non-zero. Therefore, the above system of equations has non-trivial solutions

$$\therefore \begin{vmatrix} 1 & -c & -b \\ c & -1 & a \\ b & a & -1 \end{vmatrix} = 0 \Rightarrow a^2 + b^2 + c^2 + 2abc = 1$$

174 (c)

$$A^2 - 4A + 10I = A$$

$$\Rightarrow \begin{bmatrix} 1 & -3 \\ 2 & k \end{bmatrix} \begin{bmatrix} 1 & -3 \\ 2 & k \end{bmatrix} - 4 \begin{bmatrix} 1 & -3 \\ 2 & k \end{bmatrix} + 10 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -3 \\ 2 & k \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -5 & -3 - 3k \\ 2 + 2k & -6 + k^2 \end{bmatrix} - \begin{bmatrix} 4 & -12 \\ 8 & 4k \end{bmatrix} + \begin{bmatrix} 10 & 0 \\ 0 & 10 \end{bmatrix} = \begin{bmatrix} 1 & -3 \\ 2 & k \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & 9 - 3k \\ -6 + 2k & 4 + k^2 - 4k \end{bmatrix} = \begin{bmatrix} 1 & -3 \\ 2 & k \end{bmatrix}$$

$$\Rightarrow 9 - 3k = -3, -6 + 2k = 2 \quad \dots(i)$$

$$\text{and } 4 + k^2 - 4k = k$$

$$\Rightarrow k^2 - 5k + 4 = 0 \Rightarrow k = 4, 1$$

But $k = 1$ is not satisfied the Eq (i).

175 (a)

$$\text{Given, } A^2 = 2A - I$$

$$\text{Now, } A^3 = A^2 \cdot A = 2A^2 = -IA$$

$$= 2A^2 - A = 2(2A - I) - A$$

$$= 3A - 2I = 3A - (3 - 1)I$$

$$\dots \dots \dots \dots \dots$$

$$\dots \dots \dots \dots \dots$$

$$A^n = nA - (n - 1)I$$

176 (c)

$$\text{We have, } \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} A = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$$

$$\Rightarrow A = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$$

$$\Rightarrow A = \begin{bmatrix} 1 & -3 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 4 \\ 0 & -1 \end{bmatrix}$$

177 (b)

It is given that A is an orthogonal matrix

$$\therefore A A^T = I = A^T A \Rightarrow A^{-1} = A^T$$

178 (a)

$$\text{Let } A = IA$$

$$\Rightarrow \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 4 & 6 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A$$

Applying $R_2 \rightarrow R_2 - 2R_1$ and $R_3 \rightarrow R_3 - 3R_1$

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & -1 & -2 \\ 0 & -2 & -3 \end{bmatrix} \approx \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ -3 & 0 & 1 \end{bmatrix} A$$

Applying $R_3 \rightarrow R_3 - 2R_2$

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & -1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \approx \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & -2 & 1 \end{bmatrix} A$$

Applying $R_2 \rightarrow -R_2$ and $R_2 \rightarrow R_2 - 2R_3$

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \approx \begin{bmatrix} 1 & 0 & 0 \\ 0 & 3 & -2 \\ 1 & -2 & 1 \end{bmatrix} A$$

Applying $R_1 \rightarrow R_1 - 2R_2 - 3R_3$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \approx \begin{bmatrix} -2 & 0 & 1 \\ 0 & 3 & -2 \\ 1 & -2 & 1 \end{bmatrix} A$$

$$\therefore A^{-1} = \begin{bmatrix} -2 & 0 & 1 \\ 0 & 3 & -2 \\ 1 & -2 & 1 \end{bmatrix}$$

179 (a)

$$\text{Given that, } 2X + \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 3 & 8 \\ 7 & 2 \end{bmatrix}$$

$$\Rightarrow 2X = \begin{bmatrix} 3 & 8 \\ 7 & 2 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

$$\Rightarrow 2X = \begin{bmatrix} 2 & 6 \\ 4 & -2 \end{bmatrix} = 2 \begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix}$$

$$\Rightarrow X = \begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix}$$

180 (b)

Since the given matrix is symmetric

$$\therefore (A)_{12} = (A)_{21} \Rightarrow x + 2 = 2x - 3 \Rightarrow x = 5$$

181 (d)

$$\text{Given, } A = 3 \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

$$\therefore A^2 = 3 \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \cdot 3 \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} 9A$$

$$\therefore A^4 = A^2 \cdot A^2 = 9A \cdot 9A = 81 \cdot 9A = 729A$$

182 (a)

$$\text{Now, } \begin{vmatrix} 1 & \omega^2 & \omega \\ \omega^2 & \omega & 1 \\ \omega & 1 & \omega^2 \end{vmatrix}$$

$$= 1(\omega^3 - 1) - \omega^2(\omega^4 - \omega) + \omega(\omega^2 - \omega^2)$$

$$= 1(1 - 1) - \omega^2(\omega - \omega) + 0$$

$$= 0$$

Hence, matrix A is singular

183 (a)

Given system of equations are

$$x + y + z = 6, x + 2y + 3z = 10$$

$$\text{and } x + 2y + \lambda z = \mu$$

The given system of equations has infinite number of solutions, if any two equations will be same i.e., the last two equations will be same, if $\lambda = 3, \mu = 10$.

184 (a)

$$\text{Given, } (A + B)(A - B) = A^2 - B^2$$

$$\Rightarrow A^2 - AB + BA - B^2 = A^2 - B^2$$

$$\Rightarrow AB = BA$$

$$\text{Now, } (ABA^{-1})^2 = (BAA^{-1})^2 = B^2$$

185 (c)

Since diagonal elements of a skew-symmetric matrix are all zeros i.e. $a_{ii} = 0$ for all i

$$\therefore \text{tr}(A) = \sum_{i=1}^n a_{ii} = 0$$

186 (c)

$$\therefore P^6 = P(I - P) \quad \therefore P^2 = I - P$$

$$= PI - P^2 = PI - (I - P)$$

$$\text{Now, } P^4 = P \cdot P^3$$

$$\Rightarrow P^4 = P(2P - I)$$

$$\Rightarrow P^4 = 2P^2 - P$$

$$\Rightarrow P^4 = 2I - 2P - P$$

$$\Rightarrow P^4 = 2I - 3P$$

$$\text{And } P^5 = P(2I - 3P)$$

$$\Rightarrow P^5 = 2P - 3(I - P)$$

$$\Rightarrow P^5 = 5P - 3I$$

$$\text{Also, } P^6 = P(5P - 3I)$$

$$\Rightarrow P^6 = 5P^2 - 3P$$

$$\Rightarrow P^6 = 5(I - P) - 3P$$

$$\Rightarrow P^6 = 5I - 8P$$

$$\text{So, } n = 6$$

Alternate Solution

$$\therefore P^n = 5I - 8P$$

$$= 5(I - P) - 3P$$

$$= P(5P - 3I) \quad (\therefore P^2 = I - P)$$

$$= P(2P - 3P^2)$$

$$= P^2(2I - 3P)$$

$$= P^2[2(I - P) - P]$$

$$= P^2[2P^2 - P]$$

$$= P^3[2P - I]$$

$$= P^4[I - P]$$

$$= P^4 \cdot P^2 = P^6$$

$$\Rightarrow n = 6$$

187 (b)

$$A^2 = A \cdot A = AB \cdot A$$

$$= A \cdot BA = AB = A$$

189 (d)

$$\text{Let } A = \begin{bmatrix} 1 & 2 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\therefore |A| = \begin{vmatrix} 1 & 2 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 2 \\ 0 & 1 \end{vmatrix} = 1$$

$$\text{and } \text{adj } A = \begin{bmatrix} 1 & -2 & 7 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{hence, } A^{-1} = \frac{1}{|A|} \text{adj } A = \begin{bmatrix} 1 & -2 & 7 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{so, required element} = A_{13}^{-1} = 7$$

190 (a)

$$\therefore |A| = 1$$

$$\text{and } A^c = \begin{bmatrix} \cos x & \sin x & 0 \\ -\sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{and } \text{adj } A = (A^c)' = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{\text{adj } A}{|A|} = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix} = f(-x)$$

191 (b)

$$\therefore |A| = 1(0 - 1) = -1$$

\therefore Cofactors of A are

$$C_{11} = 0, C_{12} = 0, C_{13} = -1$$

$$C_{21} = 0, C_{22} = -1, C_{23} = 0$$

$$C_{31} = -1, C_{32} = 0, C_{33} = 0$$

$$\therefore A^{-1} = \frac{1}{-1} \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix} = A$$

192 (b)

We have,

$$A^2 - 5I_2 - \begin{bmatrix} 10 & 15 \\ 15 & 25 \end{bmatrix} - \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix} = \begin{bmatrix} 5 & 15 \\ 15 & 20 \end{bmatrix} = 5A$$

$$\therefore k = 5$$

194 (b)

$$\text{Let } A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -2 & -2 \\ 1 & 3 & 1 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

$$\text{and } B = \begin{bmatrix} 0 \\ 3 \\ 4 \end{bmatrix}$$

$$\therefore AX = B$$

$$\Rightarrow X = A^{-1}B$$

$$\text{Here, } A^{-1} = \frac{1}{6} \begin{bmatrix} 4 & 2 & 0 \\ -3 & 0 & 3 \\ 5 & -2 & -3 \end{bmatrix}$$

$$\therefore X = \frac{1}{6} \begin{bmatrix} 4 & 2 & 0 \\ -3 & 0 & 3 \\ 5 & -2 & -3 \end{bmatrix} \begin{bmatrix} 0 \\ 3 \\ 4 \end{bmatrix}$$

$$= \frac{1}{6} \begin{bmatrix} 0+6+0 \\ 0+0+12 \\ 0-6-12 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix}$$

$$\text{Thus, } \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ -3 \end{bmatrix}$$

195 (a)

The given system of equations can be rewritten as matrix form $AX = B$ as

$$\begin{bmatrix} 1 & -1 & 1 \\ 1 & 2 & -1 \\ 2 & 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\text{Now, } |A| = 1(6 + 1) + 1(3 + 2) + 1(1 - 4)$$

$$= 7 + 5 - 3 = 9 \neq 0$$

Since, $|A| \neq 0$. So, the given system of equations has only trivial solution. So, there is no non-trivial solution.

196 (d)

If matrix has no inverse it means the value of

determinant should be zero.

$$\therefore \begin{vmatrix} 1 & -1 & x \\ 1 & x & 1 \\ x & -1 & 1 \end{vmatrix} = 0$$

If we put $x = 1$, then column 1st and 3rd are identical.

197 (b)

Since, $\begin{bmatrix} 2+x & 3 & 4 \\ 1 & -1 & 2 \\ x & 1 & -5 \end{bmatrix}$ is a singular matrix

$$\therefore \begin{vmatrix} 2+x & 3 & 4 \\ 1 & -1 & 2 \\ x & 1 & -5 \end{vmatrix} = 0$$

$$\Rightarrow (2+x)(5-2) - 3(-5-2x) + 4(1+x) = 0$$

$$\Rightarrow 6 + 3x + 15 + 6x + 4 + 4x = 0$$

$$\Rightarrow 13x + 25 = 0 \Rightarrow x = -\frac{25}{13}$$

198 (a)

We have,

$$A = \begin{bmatrix} 2 & 3 & 1 & 4 \\ 0 & 1 & 2 & -1 \\ 0 & -2 & -4 & 2 \end{bmatrix}$$

$$\Rightarrow A \sim \begin{bmatrix} 2 & 3 & 1 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & -2 & -4 & 2 \end{bmatrix} \quad \text{Applying } R_2 \rightarrow 2R_2 +$$

$$R_3 \Rightarrow A \sim \begin{bmatrix} 2 & 3 & -5 & 7 \\ 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \end{bmatrix} \quad \begin{array}{l} \text{Applying } C_3 \rightarrow C_3 - 2C_2 \\ C_4 \rightarrow C_4 + C_2 \end{array}$$

$$\Rightarrow A \sim \begin{bmatrix} 2 & 3 & -5 & 7 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad \text{Applying } R_2 \leftrightarrow R_3$$

Clearly, $\begin{vmatrix} 2 & 3 \\ 0 & -2 \end{vmatrix} \neq 0$ and every minor of order 3 is zero

Hence, rank of A is 2

199 (b)

We have,

$$A^2 = AA = \begin{bmatrix} a & b \\ b & a \end{bmatrix} \begin{bmatrix} a & b \\ b & a \end{bmatrix} = \begin{bmatrix} a^2 + b^2 & 2ab \\ 2ab & a^2 + b^2 \end{bmatrix}$$

$$\therefore A^2 = \begin{bmatrix} \alpha & \beta \\ \beta & \alpha \end{bmatrix}$$

$$\Rightarrow \alpha = a^2 + b^2, \beta = 2ab$$

200 (d)

In a square matrix, the trace of A is defined as the sum of the diagonal elements

$$\text{Hence, trace of } A = \sum_{i=1}^n a_{ii}$$

201 (a)

Given system of equations is $x + 2y + 3z = 1$,

$$2x + y + 3z = 2 \quad \text{and} \quad 5x + 5y + 9z = 5$$

$$\text{Now, } \Delta = \begin{vmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \\ 5 & 5 & 9 \end{vmatrix}$$

$$= 1(9 - 15) - 2(18 - 15) + 3(10 - 5)$$

$$= -6 - 6 + 15$$

$$= 3 \neq 0$$

Hence, it has unique solution

202 (a)

$$\text{Let } \Delta = \begin{vmatrix} 4 & 2 & (1-x) \\ 5 & k & 1 \\ 6 & 3 & (1+x) \end{vmatrix}$$

Applying $R_1 \rightarrow R_1 + R_3$

$$\Rightarrow \Delta = \begin{vmatrix} 10 & 5 & 2 \\ 5 & k & 1 \\ 6 & 3 & 1+x \end{vmatrix}$$

Applying $C_1 \rightarrow C_1 - 2C_2$

$$\Rightarrow \Delta = \begin{vmatrix} 0 & 5 & 2 \\ 5-2k & k & 1 \\ 0 & 3 & 1+x \end{vmatrix}$$

$$\Rightarrow (5-2k)(5+5x-6) = 0$$

$$\Rightarrow k = \frac{5}{2}, \quad x = \frac{1}{5}$$

204 (b)

$$\text{Since, } \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{bmatrix} \cdot \frac{1}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{bmatrix}$$

205 (c)

It is a direct consequence of the definition of rank

206 (a)

$$\text{Now, } A(x)A(y) = (1-x)^{-1} \begin{bmatrix} 1 & -x \\ -x & 1 \end{bmatrix} (1-y-11-y-y1)$$

$$= [(1+xy) - (x+y)]^{-1} \begin{bmatrix} 1+xy & -(x+y) \\ -(x+y) & 1+xy \end{bmatrix}$$

$$= \left(1 - \frac{x+y}{1+xy}\right)^{-1} \begin{bmatrix} 1 & -\frac{x+y}{1+xy} \\ -\frac{x+y}{1+xy} & 1 \end{bmatrix}$$

$$= A(z)$$

207 (a)

$$A^2(\alpha) = \begin{vmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{vmatrix} \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 \alpha - \sin^2 \alpha & 2\cos \alpha \sin \alpha \\ -2\sin \alpha \cos \alpha & \cos^2 \alpha - \sin^2 \alpha \end{bmatrix}$$

$$= \begin{bmatrix} \cos 2\alpha & \sin 2\alpha \\ -\sin 2\alpha & \cos 2\alpha \end{bmatrix} = A(2\alpha)$$

208 (a)

Since, A is symmetric matrix, therefore $A^T = A$

$$\text{Now, } (A^n)^T = (A^T)^n = A^n$$

Hence, A^n is a symmetric matrix.

209 (a)

$$\text{Let } A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$$

$$\therefore |A| = \begin{vmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{vmatrix}$$

$$= 0 \begin{vmatrix} 2 & 3 \\ 1 & 1 \end{vmatrix} - 1 \begin{vmatrix} 1 & 3 \\ 3 & 1 \end{vmatrix} + 2 \begin{vmatrix} 1 & 2 \\ 3 & 1 \end{vmatrix}$$

$$= -(1-9) + 2(1-6) = 8 - 10 = -2$$

$$\text{and } \text{Adj } A = \begin{bmatrix} -1 & 8 & -5 \\ 1 & -6 & 3 \\ -1 & 2 & -1 \end{bmatrix}^T = \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix}$$

$$\text{Hence, } A^{-1} = \frac{1}{|A|} (\text{adj } A)$$

$$= -\frac{1}{2} \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} 1/2 & -1/2 & 1/2 \\ -4 & 3 & -1 \\ 5/2 & -3/2 & 1/2 \end{bmatrix}$$

210 (a)

We know that

$$A = \frac{1}{2}(A + A^T) + \frac{1}{2}(A - A^T)$$

Clearly, $\frac{1}{2}(A + A^T)$ is a symmetric matrix and

$\frac{1}{2}(A - A^T)$ is a skew-symmetric matrix

Now,

$$\frac{1}{2}(A + A^T) = \frac{1}{2} \left\{ \begin{bmatrix} 2 & 0 & -3 \\ 4 & 3 & 1 \\ -5 & 7 & 2 \end{bmatrix} + \begin{bmatrix} 2 & 4 & -5 \\ 0 & 3 & 7 \\ -3 & 1 & 2 \end{bmatrix} \right\}$$

$$\Rightarrow \frac{1}{2}(A + A^T) = \frac{1}{2} \begin{bmatrix} 4 & 4 & -8 \\ 4 & 6 & 8 \\ -8 & 8 & 4 \end{bmatrix} = \begin{bmatrix} 2 & 2 & -4 \\ 2 & 3 & 4 \\ -4 & 4 & 2 \end{bmatrix}$$

211 (c)

Since, the system of linear equations has a non-zero solution, then

$$\begin{bmatrix} 1 & 2a & a \\ 1 & 3b & b \\ 1 & 4c & c \end{bmatrix} = 0$$

Applying $R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1$

$$\Rightarrow \begin{vmatrix} 1 & 2a & a \\ 0 & 3b-2a & b-a \\ 0 & 4c-2a & c-a \end{vmatrix} = 0$$

$$\Rightarrow (3b-2a)(c-a) - (4c-2a)(b-a) = 0$$

$$\Rightarrow 3bc - 3ba - 2ac + 2a^2$$

$$= 4bc - 2ab - 4ac + 2a^2$$

$$\Rightarrow 2ac = bc + ab$$

On dividing by abc both sides, we get

$$\frac{2}{b} = \frac{1}{a} + \frac{1}{c}$$

$$\Rightarrow a, b, c \text{ are in HP.}$$

212 (c)

Given system of equations is

$$x - y + z = 3$$

$$2x + y - z = 2$$

$$\text{and } -3x - 2ky + 6z = 3$$

\therefore The given system will have infinite solutions.

$$\therefore \begin{vmatrix} 1 & -1 & 1 \\ 2 & 1 & -1 \\ -3 & -2k & 6 \end{vmatrix} = 0$$

$$\Rightarrow 6k - 18 = 0 \Rightarrow k = 3$$

213 (a)

The product of two orthogonal matrix is an orthogonal matrix

214 (b)

Given system of equations can be rewritten as

$$AX = B$$

$$\Rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 2 \\ 3 & 5 & -7 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 8 \\ 6 \\ 14 \end{bmatrix}$$

$$\therefore |A| = 1(7 - 10) - 1(-7 - 6) + 1(5 + 3)$$

$$= -3 + 13 + 8 = 18 \neq 0$$

\therefore Given system has unique solution.

215 (a)

Given, equations $(x + ay = 0, az + y = 0, ax + z = 0)$ has infinite solutions.

\therefore Using Cramer's rule, its determinant = 0

$$\Rightarrow \begin{vmatrix} 1 & a & 0 \\ 0 & 1 & a \\ a & 0 & 1 \end{vmatrix} = 0$$

$$\Rightarrow 1 + a^3 = 0 \Rightarrow a = -1$$

217 (a)

$$\text{Given that, } F(\alpha) = \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow F(-\alpha) = \begin{bmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$\therefore F(\alpha)F(-\alpha)$

$$= \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2 \alpha + \sin^2 \alpha & \cos \alpha \sin \alpha - \sin \alpha \cos \alpha \\ \sin \alpha \cos \alpha - \sin \alpha \cos \alpha & \sin^2 \alpha + \cos^2 \alpha \\ 0 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

$$\Rightarrow [F(\alpha)]^{-1} = F(-\alpha)$$

218 (b)

By using inverse of matrix, we know

$$|M^{-1}| = |M|^{-1} \text{ holds true}$$

$$(M^T)^{-1} = (M^{-1})^T \text{ holds true}$$

$$\text{and } (M^{-1})^{-1} = M \text{ holds true}$$

$$\text{but } (M^2)^{-1} = (M^{-1})^{-2} \text{ not true}$$

219 (a)

$$\text{Since, } A^2 - B^2 = (A - B)(A + B)$$

$$= A^2 - B^2 + AB - BA$$

$$\Rightarrow AB = BA$$

220 (c)

$$\text{Given, } x_1 + 2x_2 + 3x_3 = 0 \quad \dots(i)$$

$$2x_1 + 3x_2 + x_3 = 0 \quad \dots(ii)$$

$$3x_1 + x_2 + 2x_3 = 0 \quad \dots(iii)$$

$$\Rightarrow \Delta = \begin{vmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \\ 3 & 1 & 2 \end{vmatrix}$$

$$= 1(6 - 1) - 2(4 - 3) + 3(2 - 9)$$

$$= -18$$

Then, this system has the unique solution.

222 (d)

$$X^2 = X \cdot X = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 5 & -8 \\ 2 & -3 \end{bmatrix}$$

For $n = 2$, no option is satisfied

Hence, option (d) is correct

223 (a)

We have,

$$F(\alpha)F(-\alpha)$$

$$= \begin{bmatrix} \cos \alpha & -\sin \alpha & 1 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos \alpha & \sin \alpha & 0 \\ -\sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow F(\alpha)F(-\alpha) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = I$$

$$\Rightarrow F(-\alpha) = [F(\alpha)]^{-1}$$

224 (b)

$$\text{We have, } (AA^T)^T = (A^T)^T A^T = AA^T$$

$\therefore AA^T$ is symmetric matrix

225 (c)

For any square matrix X , we have

$$X(\text{adj } X) = |X| I_n$$

Taking $X = \text{adj } A$, we have

$$(\text{adj } A)(\text{adj } (\text{adj } A)) = |\text{adj } A| I_n$$

$$\Rightarrow \text{adj } A(\text{adj } (\text{adj } A)) = |A|^{n-1} I_n \quad [\because |\text{adj } A| = |A|^{n-1}]$$

$$\Rightarrow (A \text{ adj } A)(\text{adj } (\text{adj } A)) = |A|^{n-1} A \quad [\because A I_n = A]$$

$$\Rightarrow (|A| I_n)(\text{adj } (\text{adj } A)) = |A|^{n-1} A$$

$$\Rightarrow \text{adj } (\text{adj } A) = |A|^{n-2} A$$

226 (d)

Given equations are

$$(\alpha + 1)^3 x + (\alpha + 2)^3 y - (\alpha + 3)^3 z = 0$$

$$(\alpha + 1)x + (\alpha + 2)y - (\alpha + 3)z = 0$$

$$\text{and } x + y - z = 0$$

Since, this system of equations is consistent.

$$\therefore \begin{vmatrix} (\alpha + 1)^3 & (\alpha + 2)^3 & -(\alpha + 3)^3 \\ (\alpha + 1) & (\alpha + 2) & -(\alpha + 3) \\ 1 & 1 & -1 \end{vmatrix} = 0$$

Applying $C_2 \rightarrow C_2 - C_1$ and $C_3 \rightarrow C_3 + C_1$

$$\Rightarrow \begin{bmatrix} (\alpha + 1)^3 & (\alpha + 2)^3 - (\alpha + 1)^3 & -(\alpha + 1)^3 \\ (\alpha + 1)^3 - (\alpha + 1)^3 & (\alpha + 1)^3 - (\alpha + 3)^3 & -(\alpha + 1)^3 \\ (\alpha + 1) & (\alpha + 2) - (\alpha + 1) & -(\alpha + 3) + (\alpha + 1) \end{bmatrix} = 0$$

$$\Rightarrow \begin{bmatrix} (\alpha + 1)^3 & 3\alpha^2 + 9\alpha + 7 & -6\alpha^2 - 24\alpha - 26 \\ (\alpha + 1) & 1 & -2 \\ 1 & 0 & 0 \end{bmatrix} = 0$$

$$\Rightarrow -2(3\alpha^2 + 9\alpha + 7) + 6\alpha^2 + 24\alpha + 26 = 0$$

$$\Rightarrow 6\alpha + 12 = 0 \Rightarrow \alpha = -2$$

227 (a)

We have,

$$A^n = \begin{bmatrix} \cos n\theta & \sin n\theta \\ -\sin n\theta & \cos n\theta \end{bmatrix}$$

$$\Rightarrow \frac{1}{n}A^n = \begin{bmatrix} \frac{\cos n\theta}{n} & \frac{\sin n\theta}{n} \\ -\frac{\sin n\theta}{n} & \frac{\cos n\theta}{n} \end{bmatrix}$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n}A^n = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

228 (c)

We have,

$$AB = O$$

$$\Rightarrow \begin{bmatrix} \cos^2 \alpha & \cos \alpha \sin \alpha \\ \cos \alpha \sin \alpha & \sin^2 \alpha \end{bmatrix} \begin{bmatrix} \cos^2 \beta & \cos \beta \sin \beta \\ \cos \beta \sin \beta & \sin^2 \beta \end{bmatrix} = O$$

$$\Rightarrow \begin{bmatrix} \cos \alpha \cos \beta \cos(\alpha - \beta) & \cos \alpha \sin \beta \cos(\alpha - \beta) \\ \cos \beta \sin \alpha \cos(\alpha - \beta) & \sin \alpha \sin \beta \cos(\alpha - \beta) \end{bmatrix} = O$$

$$\Rightarrow \cos(\alpha - \beta) = 0$$

$$\Rightarrow \alpha - \beta \text{ is an odd multiple of } \frac{\pi}{2}$$

229 (c)

$$|A| |\text{adj } A| = |A|^n \text{ for order } n$$

$$\Rightarrow DD' = D^n$$

230 (c)

$$\text{Given, } \begin{bmatrix} 1 + \omega & 2\omega \\ -2\omega & -b \end{bmatrix} + \begin{bmatrix} a & -\omega \\ 3\omega & 2 \end{bmatrix} = \begin{bmatrix} 0 & \omega \\ \omega & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 + \omega + a & \omega \\ \omega & 2 - b \end{bmatrix} = \begin{bmatrix} 0 & \omega \\ \omega & 1 \end{bmatrix}$$

$$\Rightarrow 1 + \omega + a = 0, 2 - b = 1$$

$$\Rightarrow a = -1 - \omega, b = 1$$

$$\therefore a^2 + b^2 = (-1 - \omega)^2 + 1^2$$

$$= 1 + \omega^2 + 2\omega + 1^2$$

$$= 0 + \omega + 1 \quad (\because 1 + \omega + \omega^2 = 0)$$

$$= 1 + \omega$$

232 (a)

$$\text{Given, } 2kx - 2y + 3z = 0, x + ky + 2z = 0, 2x + kz = 0$$

For non-trivial solution

$$\begin{vmatrix} 2k & -2 & 3 \\ 1 & k & 2 \\ 2 & 0 & k \end{vmatrix} = 0$$

$$\Rightarrow 2k(k^2 - 0) + 2(k - 4) + 3(0 - 2k) = 0$$

$$\Rightarrow 2k^3 - 4k - 8 = 0$$

$$\Rightarrow (k - 2)(2k^2 + 4k + 4) = 0$$

$$\Rightarrow k = 2$$

233 (a)

$$AB = \begin{bmatrix} 2 & 1 & 3 \\ 4 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 5 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 + 0 + 15 & -2 + 2 + 0 \\ 4 + 0 + 0 & -4 + 2 + 0 \end{bmatrix}$$

$$= \begin{bmatrix} 17 & 0 \\ 4 & -2 \end{bmatrix}$$

234 (d)

$$A^2 = \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \alpha & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} \alpha^2 & 0 \\ \alpha + 1 & 1 \end{bmatrix}$$

$$\therefore A^2 = B \Rightarrow \begin{bmatrix} \alpha^2 & 0 \\ \alpha + 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix}$$

$$\Rightarrow \alpha^2 = 1 \text{ and } \alpha + 1 = 5$$

Which is not possible at the same time.

\(\therefore\) No real values of \(\alpha\) exists.

235 (c)

We have,

$$E(\alpha)E(\beta)$$

$$= \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{bmatrix}$$

$$= \begin{bmatrix} \cos(\alpha + \beta) & \sin(\alpha + \beta) \\ -\sin(\alpha + \beta) & \cos(\alpha + \beta) \end{bmatrix} = E(\alpha + \beta)$$

Hence, option (c) is correct.

237 (c)

$$\text{We have, } A = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix}$$

$$\therefore A^2 = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \end{bmatrix}$$

$$= \begin{bmatrix} \frac{1}{2} - \frac{1}{2} & \frac{1}{2} - \frac{1}{2} \\ -\frac{1}{2} + \frac{1}{2} & -\frac{1}{2} + \frac{1}{2} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = O$$

\(\therefore\) Matrix \(A\) is nilpotent

238 (d)

$$\text{Since, } A = \begin{bmatrix} 1 & -1 & 2 \\ 2 & 0 & 1 \\ 3 & 2 & 1 \end{bmatrix}$$

$$\text{Now, } |A| = 1(0 - 2) + 1(2 - 3) + 2(4 - 0) = 5$$

$$\therefore A^{-1} = \frac{1}{5} \begin{bmatrix} -2 & 5 & -1 \\ 1 & -5 & 3 \\ 4 & -5 & 2 \end{bmatrix}$$

$$\text{Now, } A^{-1}B = \frac{1}{5} \begin{bmatrix} -2 & 5 & -1 \\ 1 & -5 & 3 \\ 4 & -5 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \\ 4 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 3 \end{bmatrix}$$

239 (a)

Since, A is a skew-symmetric matrix. Therefore,

$$A^T = -A \Rightarrow |A^T| = |-A|$$

$$\Rightarrow |A| = (-1)^n |A|$$

Also, n is odd

$$\therefore 2|A| = 0 \Rightarrow |A| = 0$$

$$\text{Thus, } |\text{adj } A| = |A|^2 = 0$$

240 (d)

Given System of equations are

$$x + 3y + 2z = 0$$

$$3x + y + z = 0$$

$$\text{and } 2x - 2y - z = 0$$

$$\text{Now, } \Delta = \begin{vmatrix} 1 & 3 & 2 \\ 3 & 1 & 1 \\ 2 & -2 & -1 \end{vmatrix}$$

$$= 1(-1 + 2) - 3(-3 - 2) + 2(-6 - 2)$$

$$= 1 + 15 - 16$$

$$= 0$$

Since, determinant is zero, then it has infinitely many solutions.

242 (b)

$$\text{Let } \Delta = \begin{vmatrix} a_1 & a_2 \\ a_4 & a_5 \end{vmatrix}$$

$$= a_1 a_5 - a_2 a_4$$

$$= a_1(a_1 + 4d) - (a_1 + d)(a_1 + 3d)$$

$$= a_1^2 + 4a_1 d - a_1^2 - 4a_1 d - 3d^2 = -3d^2 \neq 0$$

Hence, given system of equations has unique solution.

243 (b)

$$\therefore I_n = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{bmatrix}, |I_n| = 1$$

$$\text{adj } (I_n) = I_n$$

$$\therefore (I_n)^{-1} = I_n$$

244 (c)

We know that

$$(\text{adj } A)^T = \text{adj } A^T$$

$$\Rightarrow \text{adj } A^T - (\text{adj } A)^T = O \text{ (Null matrix)}$$

245 (c)

$$\text{Given, } \begin{bmatrix} 2 & -1 & 3 \\ 1 & 3 & -1 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 9 \\ 4 \\ 10 \end{bmatrix}$$

It is of the form $AX = B$... (i)

$$|A| = 2(3 + 2) + 1(1 + 3) + 3(2 - 9) = -7$$

$$\therefore \text{adj } (A) = \begin{bmatrix} 5 & 7 & -8 \\ -4 & -7 & 5 \\ -7 & -7 & 7 \end{bmatrix}$$

$$\Rightarrow A^{-1} = \frac{1}{-7} \begin{bmatrix} 5 & 7 & -8 \\ -4 & -7 & 5 \\ -7 & -7 & 7 \end{bmatrix}$$

$$\text{From Eq.(i), } X = -\frac{1}{7} \begin{bmatrix} 5 & 7 & -8 \\ -4 & -7 & 5 \\ -7 & -7 & 7 \end{bmatrix} \begin{bmatrix} 9 \\ 4 \\ 10 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} x \\ y \\ z \end{bmatrix} = -\frac{1}{7} \begin{bmatrix} -7 \\ -14 \\ -21 \end{bmatrix}$$

$$\Rightarrow x = 1, y = 2, z = 3$$

247 (b)

$$A = \begin{bmatrix} 1 & -1 & 1 \\ 2 & 1 & -1 \\ 4 & 1 & 1 \end{bmatrix}$$

$$= 1(1 + 1) + 1(2 + 4) + 1(2 - 4) = 6 \neq 0$$

Hence, it has unique solution.

248 (d)

$$\text{Given, } A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$\text{Now, } |A| = \cos^2 \theta + \sin^2 \theta = 1 \neq 0.$$

$\therefore A$ is invertible.

249 (b)

$$|A| = -1 \text{ and } \text{adj } A = \begin{bmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{-1} \begin{bmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix} = A$$

250 (b)

$$A = \begin{bmatrix} -1 & 2 & 5 \\ 2 & -4 & a-4 \\ 1 & -2 & a+1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & a+6 \\ 0 & 0 & -a-6 \\ 1 & -2 & a+1 \end{bmatrix}$$

[using $R_1 \rightarrow R_1 + R_3$ and $R_2 \rightarrow R_2 - 2R_3$]

$$= \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -a-6 \\ 1 & -2 & a+1 \end{bmatrix} \quad [\text{using } R_1 \rightarrow R_1 + R_2]$$

$$\text{When } a = -6, A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & -2 & -5 \end{bmatrix} \quad \therefore \rho(A) = 1$$

$$\text{When } a = 6, A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -12 \\ 1 & -2 & 7 \end{bmatrix}, \quad \therefore \rho(A) = 2$$

$$\text{When } a = 1, A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -7 \\ 1 & -2 & 2 \end{bmatrix}, \quad \therefore \rho(A) = 2$$

$$\text{When } a = 2, A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -8 \\ 1 & -2 & 3 \end{bmatrix} \quad \therefore \rho(A) = 2$$

251 (b)

$$\text{Let } D = \begin{bmatrix} d_1 & 0 & 0 & \dots & 0 \\ 0 & d_2 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & d_n \end{bmatrix}$$

Then, $|D| = d_1 d_2 \dots d_n$

Now, Cofactor of $D_{11} = d_2 d_3 \dots d_n$

Cofactor of $D_{22} = d_1 d_3 \dots d_n$ etc

And, Cofactor of $D_{ij} = 0$ when $i \neq j$

$$\therefore D^{-1} = \frac{1}{|D|} \text{adj } D$$

$$= \frac{1}{d_1 d_2 \cdots d_n} \begin{bmatrix} d_2 d_3 \cdots d_n & 0 & 0 & \cdots & 0 \\ 0 & d_2 d_3 \cdots d_n & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & d_1 d_2 \cdots \end{bmatrix}$$

$$\therefore D^{-1} = \begin{bmatrix} \frac{1}{d_1} & 0 & 0 & \cdots & 0 \\ 0 & \frac{1}{d_2} & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \frac{1}{d_n} \end{bmatrix} = \text{diag}(s_1^{-1} d_2^{-1} \cdots d_n^{-1})$$

252 (d)

$$\therefore A^2 = \begin{bmatrix} 1 & 2 \\ 4 & -3 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 4 & -3 \end{bmatrix} = \begin{bmatrix} 9 & -4 \\ -8 & 17 \end{bmatrix}$$

$$\therefore f(A) = A^2 + 4A - 5$$

$$= \begin{bmatrix} 9 & -4 \\ -8 & 17 \end{bmatrix} + \begin{bmatrix} 4 & 8 \\ 16 & -12 \end{bmatrix} - \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$$

$$= \begin{bmatrix} 8 & 4 \\ 8 & 0 \end{bmatrix}$$

253 (a)

$$A^2 = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & -4 \\ -4 & 5 \end{bmatrix}$$

Again now, $4A - 3I = 4 \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} - \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} =$

$$\begin{bmatrix} 5 & -4 \\ -4 & 5 \end{bmatrix}$$

$$\therefore A^2 = 4A - 3I$$

254 (d)

$$\therefore (AB)^{-1} = B^{-1}A^{-1}$$

255 (d)

$$|A| = -8$$

$$\text{adj}(A) = \begin{bmatrix} 0 & 0 & -4 \\ 0 & -4 & 0 \\ -4 & 0 & 0 \end{bmatrix}$$

$$A^{-1} = \frac{1}{-8} \begin{bmatrix} 0 & 0 & -4 \\ 0 & -4 & 0 \\ -4 & 0 & 0 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 & 1/2 \\ 0 & 1/2 & 0 \\ 1/2 & 0 & 0 \end{bmatrix}$$

256 (b)

Since given system of equations possesses a non-zero solution.

$$\therefore \Delta = \begin{vmatrix} a & 1 & 1 \\ 1 & -a & 1 \\ 1 & 1 & 1 \end{vmatrix} = 0$$

$$\Rightarrow a(-a-1) - 1(1-1) + 1(1+a) = 0$$

$$\Rightarrow a^2 = 1 \Rightarrow a = \pm 1$$

257 (a)

$$\text{Now, } (A + A^T)^T = A^T + (A^T)^T = A^T + A$$

$\therefore A + A^T$ is symmetric matrix.

258 (c)

$$A = \begin{bmatrix} 1 & 2 & -1 \\ -1 & 1 & 2 \\ 2 & -1 & 1 \end{bmatrix} = 14$$

$$\therefore (\text{adj}(\text{adj } A)) = |A|^{n-2}A = 14^{3-2}A = 14A$$

$$\therefore |\text{adj}(\text{adj } A)| = |14A| = 14^3|A| = 14^4$$

259 (b)

Given,

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & -2 & -2 \\ 1 & 3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ 4 \end{bmatrix} \Rightarrow \begin{bmatrix} x+y+z \\ x-2y-2z \\ x+3y+z \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ 4 \end{bmatrix}$$

On Comparing both sides, we get

$$x + y + z = 0 \quad \dots(i)$$

$$x - 2y - 2z = 3 \quad \dots(ii)$$

$$\text{and } x + 3y + z = 4 \quad \dots(iii)$$

On solving Eqs. (i), (ii) and (iii), we get

$$x = 1, y = 2 \text{ and } z = -3$$

260 (a)

$$|A| = \begin{vmatrix} 1 & 2 \\ -4 & -1 \end{vmatrix}$$

$$= -1 + 8 = 7$$

$$\text{adj } A = \begin{bmatrix} -1 & -2 \\ 4 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{7} \begin{bmatrix} -1 & -2 \\ 4 & 1 \end{bmatrix}$$

261 (c)

It is given that

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} a & a+b \\ c & c+d \end{bmatrix} = \begin{bmatrix} a+c & b+d \\ c & d \end{bmatrix}$$

$$\Rightarrow a+c = a, a+b = b+d, c+d = d$$

$$\Rightarrow c = 0 \text{ and } a = d$$

262 (a)

$$AB = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 2 \\ 4 & 5 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix} = \begin{bmatrix} 5 & 1 & -3 \\ 3 & 2 & 6 \\ 14 & 5 & 0 \end{bmatrix}$$

263 (a)

Let A be a skew-symmetric matrix of odd order $(2n+1)$ say. Since A is skew-symmetric

$$\therefore A^T = -A$$

$$\Rightarrow |A^T| = |-A|$$

$$\Rightarrow |A^T| = (-1)^{2n+1}|A|$$

$$\Rightarrow |A^T| = -|A|$$

$$\Rightarrow |A| = -|A| \Rightarrow 2|A| = 0 \Rightarrow |A| = 0$$

264 (a)

$$\text{As, } PP^T = \begin{bmatrix} \sqrt{3}/2 & 1/2 \\ -1/2 & \sqrt{3}/2 \end{bmatrix} \begin{bmatrix} \sqrt{3}/2 & -1/2 \\ 1/2 & \sqrt{3}/2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow PP^T = I \text{ or } P^T = P^{-1} \quad \dots(i)$$

$$\text{As, } Q = PAP^T$$

$$\therefore P^T Q^{2005} P = P^T [PAP^T]^{2005} P \dots \text{2005 times} P$$

$$= \underbrace{(P^T P)A(P^T P)A(P^T P) \dots (P^T P)A(P^T P)}_{2005 \text{ times}}$$

$$= IA^{2005} = A^{2005}$$

$$\therefore A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, A^2 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \dots \text{and so on}$$

$$A^{2005} = \begin{bmatrix} 1 & 2005 \\ 0 & 1 \end{bmatrix}$$

$$\Rightarrow P^T Q^{2005} P = \begin{bmatrix} 1 & 2005 \\ 0 & 1 \end{bmatrix}$$

265 (c)

Given, $2X + 3Y = 0$... (i)

and $X + 2Y = I$... (ii)

where $O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ and $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

On solving Eqs. (i) and (ii), we get

$$X = -3I = \begin{bmatrix} -3 & 0 \\ 0 & -3 \end{bmatrix}$$

266 (d)

Subtracting the addition of first two equations from third equation, we get

$$0 = -5 \text{ which is an absurd result.}$$

267 (d)

Given $A = \begin{bmatrix} x & -2 \\ 3 & 7 \end{bmatrix}$

$$|A| = \begin{vmatrix} x & -2 \\ 3 & 7 \end{vmatrix} = 7x + 6$$

$$\therefore A^{-1} = \frac{1}{7x + 6} \begin{bmatrix} 7 & 2 \\ -3 & x \end{bmatrix}$$

But given $A^{-1} = \begin{bmatrix} \frac{7}{34} & \frac{1}{17} \\ \frac{-3}{34} & \frac{2}{17} \end{bmatrix}$

$$\therefore \frac{7}{7x + 6} = \frac{7}{34}$$

$$\Rightarrow 7x + 6 = 34 \Rightarrow 7x = 28 \Rightarrow x = 4$$

268 (d)

(a) It is clear that A is not a zero matrix.

(b) $(-1)I = -1 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \neq A$

ie, $(-1)I \neq A$

(c) $|A| = 0 \begin{vmatrix} -1 & 0 \\ 0 & 0 \end{vmatrix} - 0 \begin{vmatrix} 0 & 0 \\ -1 & 0 \end{vmatrix} - 1 \begin{vmatrix} 0 & -1 \\ -1 & 0 \end{vmatrix} = 0 - 0 - 1(-1) = 1$

Since, $|A| \neq 0$ so A^{-1} exists.

(d) $A^2 = \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{bmatrix}$

$$\Rightarrow A^2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \Rightarrow A^2 = I$$

270 (a)

Since, $A = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 2 \\ 4 & 5 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$

$$\therefore AB = \begin{bmatrix} 1 & 2 & -1 \\ 3 & 0 & 2 \\ 4 & 5 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$$

$$= \begin{bmatrix} 1+4+0 & 0+2-1 & 0+0-3 \\ 3+0+0 & 0+0+2 & 0+0+6 \\ 4+10+0 & 0+5+0 & 0+0+0 \end{bmatrix}$$

$$= \begin{bmatrix} 5 & 1 & -3 \\ 3 & 2 & 6 \\ 14 & 5 & 0 \end{bmatrix}$$

271 (c)

We have,

$$[F(x) G(y)]^{-1} = [G(y)]^{-1} [f(x)]^{-1}$$

$$\Rightarrow [F(x) G(y)]^{-1} = G(-y) F(-x)$$

272 (a)

$$A^{-1} = \frac{1}{1+10} \begin{bmatrix} 1 & -2 \\ 5 & 1 \end{bmatrix} = \frac{1}{11} \begin{bmatrix} 1 & -2 \\ 5 & 1 \end{bmatrix}$$

Also, $A^{-1} = xA + yI$

$$\Rightarrow \frac{1}{11} \begin{bmatrix} 1 & -2 \\ 5 & 1 \end{bmatrix} = \begin{bmatrix} x & 2x \\ -5x & x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 0 & y \end{bmatrix}$$

$$\Rightarrow x + y = \frac{1}{11}, 2x = \frac{-2}{11}$$

$$\Rightarrow x = \frac{-1}{11}, y = \frac{2}{11}$$

273 (d)

Now, $(AB)^T = B^T A^T$

274 (c)

On comparing corresponding elements, we get

$$x + y + z = 9$$

$$x + y = 5$$

$$\text{and } y + z = 7$$

On solving these, we get $x = 2, y = 3, z = 4$

$$\Rightarrow (x, y, z) = (2, 3, 4)$$

275 (c)

$$A \cdot A = \begin{bmatrix} ab & b^2 \\ -a^2 & -ab \end{bmatrix} \begin{bmatrix} ab & b^2 \\ -a^2 & -ab \end{bmatrix} = \begin{bmatrix} a^2b^2 - a^2b^2 & ab^3 - ab^3 \\ -a^3b + a^3b & -a^2b^2 + a^2b^2 \end{bmatrix} \Rightarrow A^2 = 0$$

$\therefore A$ is nilpotent matrix of order 2.

276 (d)

Since A, B and C are non-singular matrices, then

$$(AB^{-1}C)^{-1} = C^{-1}(AB^{-1})^{-1}$$

$$= C^{-1}((B^{-1})^{-1}A^{-1}) = C^{-1}BA^{-1}$$

277 (a)

Given matrix is invertible

$$\Rightarrow \begin{vmatrix} \lambda & -1 & 4 \\ -3 & 0 & 1 \\ -1 & 1 & 2 \end{vmatrix} \neq 0$$

$$\Rightarrow \lambda(0-1) + 1(-6+1) + 4(-3-0) \neq 0$$

$$\Rightarrow -\lambda - 5 - 12 \neq 0$$

$$\Rightarrow \lambda \neq -17$$

278 (d)

From Eqs. (ii) and (iii), we get

$$\frac{3y^2}{b^2} - \frac{3z^2}{c^2} = 0$$

$$\Rightarrow \frac{z^2}{c^2} = \frac{y^2}{b^2}$$

On putting this value in Eq. (i), we get

$$\frac{2x^2}{a^2} - \frac{2y^2}{b^2} = 0$$

$$\Rightarrow \frac{x^2}{a^2} = \frac{y^2}{b^2}$$

$$\therefore \frac{x^2}{a^2} = \frac{y^2}{b^2} = \frac{z^2}{c^2} = k^2 \text{ (say)}$$

$$\Rightarrow x = \pm ka, y = \pm kb, z = \pm kc, \forall k \in R$$

279 (b)

$$\text{We have, } A = \begin{bmatrix} 1 & \log_b a \\ \log_a b & 1 \end{bmatrix}$$

$$\therefore |A| = 1 - \log_a b \log_b a = 1 - 1 = 0$$

280 (a)

$$|A| = 4 - 6 = -2$$

$$\text{adj}(A) = \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix}$$

$$\therefore A^{-1} = -\frac{1}{2} \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix}$$

281 (b)

$$\text{Since, } P = \begin{bmatrix} i & 0 & -i \\ 0 & -i & i \\ -i & i & 0 \end{bmatrix} \text{ and } Q = \begin{bmatrix} -i & i \\ 0 & 0 \\ i & -i \end{bmatrix}$$

$$\therefore PQ = \begin{bmatrix} i & 0 & -i \\ 0 & -i & i \\ -i & i & 0 \end{bmatrix} \begin{bmatrix} -i & i \\ 0 & 0 \\ i & -i \end{bmatrix}$$

$$= \begin{bmatrix} -i^2 - i^2 & i^2 + i^2 \\ i^2 & -i^2 \\ i^2 & -i^2 \end{bmatrix}$$

$$= \begin{bmatrix} 1+1 & -1-1 \\ -1 & 1 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} 2 & -2 \\ -1 & 1 \\ -1 & 1 \end{bmatrix}$$

282 (d)

$$B = \text{adj}(A) = \begin{bmatrix} 3 & 1 & 1 \\ -6 & -2 & 3 \\ -4 & -3 & 2 \end{bmatrix}$$

$$\text{Therefore, } \text{adj}(B) = \begin{bmatrix} 5 & -5 & 5 \\ 0 & 10 & -15 \\ 10 & 5 & 0 \end{bmatrix}$$

$$\text{Now, } |\text{adj } B| = \begin{vmatrix} 5 & -5 & 5 \\ 0 & 10 & -15 \\ 10 & 5 & 0 \end{vmatrix} = 625$$

$$\text{and } |C| = 125|A| = 125 \begin{vmatrix} 1 & -1 & 1 \\ 0 & 2 & -3 \\ 2 & 5 & 0 \end{vmatrix} = 625$$

$$\therefore \frac{|\text{adj}(B)|}{|C|} = \frac{625}{625} = 1$$

Alternate

$$|A| = 1(0+3) + 1(0+6) + (0-4)$$

$$\text{Now, } \text{adj } B = \text{adj}(\text{adj } A)$$

$$= |A|A = 5A$$

$$\therefore \frac{|\text{adj } B|}{|C|} = \frac{|5A|}{|5A|} = 1$$

283 (d)

$$\text{Given, } A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$\therefore A^{-1} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} = A^T$$

284 (b)

The given system of equations posses non-zero solutions,

$$\therefore \begin{bmatrix} 1 & 1 & 1 \\ 1 & a & a \\ 1 & -a & 1 \end{bmatrix} = 0$$

Applying $R_2 \rightarrow R_2 - R_1$ and $R_3 \rightarrow R_3 - R_1$

$$\Rightarrow \begin{vmatrix} 1 & 1 & 1 \\ 0 & a-1 & a-1 \\ 0 & -a-1 & 0 \end{vmatrix} = 0$$

$$\Rightarrow 1(0 - (a^2 - 1)) = 0$$

$$\Rightarrow a^2 = 1 \Rightarrow a = \pm 1$$

285 (a)

$$\text{Given, } x \begin{bmatrix} -3 \\ 4 \end{bmatrix} + y \begin{bmatrix} 4 \\ 3 \end{bmatrix} = \begin{bmatrix} 10 \\ -5 \end{bmatrix}$$

$$\therefore -3x + 4y = 10 \quad \dots(i)$$

$$\text{and } 4x + 3y = -5 \quad \dots(ii)$$

On solving Eqs. (i) and (ii), we get

$$x = -2, y = 1$$

287 (a)

$$|A| = 5 + 6 = 11$$

$$\text{and } \text{adj } A = \begin{bmatrix} 1 & 2 \\ -3 & 5 \end{bmatrix}$$

$$A^{-1} = \frac{1}{11} \begin{bmatrix} 1 & 2 \\ -3 & 5 \end{bmatrix}$$

288 (c)

We know that, if

$$A^n = \begin{bmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{bmatrix} = \text{diag} [d_1 \ d_2 \ d_3]$$

Then,

$$A^n = \begin{bmatrix} d_1^n & 0 & 0 \\ 0 & d_2^n & 0 \\ 0 & 0 & d_3^n \end{bmatrix} = \text{diag} [d_1^n \ d_2^n \ d_3^n]$$

$$\therefore A^5 = \begin{bmatrix} 2^5 & 0 & 0 \\ 0 & 2^5 & 0 \\ 0 & 0 & 2^5 \end{bmatrix} = 16A$$

289 (c)

$$AB = I \Rightarrow B = A^{-1}$$

$$= \frac{1}{1 + \tan^2 \theta} \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix}$$

$$= \frac{1}{\sec^2 \theta} \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix}$$

$$\Rightarrow (\sec^2 \theta) B = \begin{bmatrix} 1 & -\tan \theta \\ \tan \theta & 1 \end{bmatrix} = A(-\theta)$$

290 (b)

It is given that $A = [a_{ij}]$ is a skew-symmetric matrix

$$a_{ij} = -a_{ji} \text{ for all } i, j$$

$$\Rightarrow a_{ii} = -a_{ii} \text{ for all } i$$

$$\Rightarrow 2a_{ii} = 0 \text{ for all } i \Rightarrow a_{ii} = 0 \text{ for all } i$$

291 (c)

We know that, if $A = \text{diag.} (d_1 \ d_2 \ \dots \ d_n)$ is a

diagonal matrix, then for any $k \in \mathbb{N}$

$$A^k = \text{diag}(d_1^k, d_2^k, \dots, d_n^k)$$

Here, $A = \text{diag}(a, a, a)$

$$\therefore A^n = \text{diag}(a^n, a^n, a^n) = \begin{bmatrix} a^n & 0 & 0 \\ 0 & a^n & 0 \\ 0 & 0 & a^n \end{bmatrix}$$

292 (b)

We have,

$$(AB - BA)^T = (AB)^T - (BA)^T \\ = B^T A^T - A^T B^T$$

$$\Rightarrow (AB - BA)^T = BA - AB \quad [\because A^T = A, B^T = B]$$

$$\Rightarrow (AB - BA)^T = -(AB - BA)$$

So, $AB - BA$ is skew-symmetric matrix

293 (b)

Since AB exists

\therefore No. of rows in $B =$ No. of columns in A

\Rightarrow No. of rows in $B = n$

Also, BA exists

\Rightarrow No. of columns in $B =$ No. of rows in A

\Rightarrow No. of columns in $B = m$

Hence, B is of order $n \times m$

294 (c)

We have,

$$(kA)(\text{adj } kA) = |kA| I_n$$

$$\Rightarrow k(A \text{ adj } kA) = k^n |A| I_n \quad [\because |kA| = k^n |A|]$$

$$\Rightarrow A(\text{adj } kA) = k^{n-1} |A| I_n$$

$$\Rightarrow A \text{ adj } (kA) = k^{n-1} A(\text{adj } A) \quad [\because A \text{ adj } A = |A| I_n]$$

$$\Rightarrow A \text{ adj } (kA) = A(k^{n-1} \text{adj } A)$$

$$\Rightarrow A^{-1}(A \text{ adj } (kA)) = A^{-1}(A(k^{n-1} \text{adj } A))$$

$$\Rightarrow (A^{-1}A)(\text{adj } kA) = (A^{-1}A)(k^{n-1} \text{adj } A)$$

$$\Rightarrow I(\text{adj } kA) = I(k^{n-1} \text{adj } A)$$

$$\Rightarrow \text{adj } kA = k^{n-1} (\text{adj } A)$$

295 (c)

It has a non-zero solution if

$$\begin{vmatrix} 1 & k & -1 \\ 3 & -k & -1 \\ 1 & -3 & 1 \end{vmatrix} = 0$$

$$\Rightarrow -6k + 6 = 0$$

$$\Rightarrow k = 1$$

296 (b)

$$(aI + bA)^2 = (aI + bA)(aI + bA)$$

$$= a^2 I^2 + aI(bA) + bA(aI) + (bA)^2$$

Now, $I^2 = I$ and $IA = A$

$$\therefore (aI + bA)^2 = a^2 I + 2abA + b^2(A^2)$$

$$\text{Now, } A^2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} = O$$

$$\therefore (aI + bA)^2 = a^2 I + 2abA$$

297 (b)

Since A is orthogonal matrix

$$\therefore A A^T = I = A^T A$$

$$\Rightarrow |A A^T| = |I| = |A^T A|$$

$$\Rightarrow |A||A^T| = 1 = |A^T||A|$$

$$\Rightarrow |A|^2 = 1 \Rightarrow |A| = \pm 1$$

298 (b)

Since, given system of equations has no solution, $\Delta = 0$ and any one amongst $\Delta x, \Delta y, \Delta z$ is non-zero.

$$\text{Where } \Delta = \begin{vmatrix} 2 & -1 & 2 \\ 1 & -2 & 1 \\ 1 & 1 & \lambda \end{vmatrix} = 0$$

$$\text{And } \Delta z = \begin{vmatrix} 2 & -1 & 2 \\ 1 & -2 & -4 \\ 1 & 1 & \lambda \end{vmatrix} = 6 \neq 0$$

$$\Rightarrow \lambda = 1$$

299 (c)

Since, A is an idempotent matrix, therefore $A^2 = A$

$$\Rightarrow \begin{bmatrix} 2 & -2 & -16 - 4x \\ -1 & 3 & 16 + 4x \\ 4 + x & -8 - 2x & -12 + x^2 \end{bmatrix}$$

$$= \begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & x \end{bmatrix}$$

On comparing, $16 + 4x = 4$

$$\Rightarrow x = -3$$

300 (a)

We have,

$$A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 3 & 0 \\ 0 & 1 & 2 \end{bmatrix} \text{ and } \text{adj } A = \begin{bmatrix} 6 & -2 & -6 \\ -4 & 2 & x \\ y & -1 & 1 \end{bmatrix}$$

Clearly, $|A| = 6 - 8 + 4 = 2$

$$\therefore A(\text{adj } A) = |A| I$$

$$\Rightarrow \begin{bmatrix} 1 & 2 & 2 \\ 2 & 3 & 0 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 6 & -2 & -6 \\ -4 & 2 & x \\ y & -1 & -1 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 2y - 2 & 0 & 2x - 18 \\ 0 & 2 & 3x - 12 \\ 2y - 4 & 0 & x - 2 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

$$\Rightarrow 2y - 2 = 2, 2y - 4 = 0, 2x - 18 = 0, 3x - 12 = 0, x - 2 = 2$$

$$\Rightarrow x = 4, y = 2 \Rightarrow x + y = 6$$

301 (a)

$$\text{Since, } \begin{bmatrix} x + y & 2x + z \\ x - y & 2z + w \end{bmatrix} = \begin{bmatrix} 4 & 7 \\ 0 & 10 \end{bmatrix}$$

$$\Rightarrow x + y = 4 \dots(i)$$

$$x - y = 0 \dots(ii)$$

$$2x + z = 7 \dots(iii)$$

$$\text{and } 2z + w = 10 \dots(iv)$$

On solving these equations, we get

$$x = 2, y = 2, z = 3, w = 4$$

302 (b)

We have,

$$A = \begin{bmatrix} -i & 0 \\ 0 & -i \end{bmatrix}$$

$$\therefore A^2 = \begin{bmatrix} -i & 0 \\ 0 & -i \end{bmatrix} \begin{bmatrix} -i & 0 \\ 0 & -i \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$\Rightarrow A^2 = -I \Rightarrow A^2 + I = O$$

303 (d)

$$m[-3 \ 4] + n[4 \ -3] = [10 \ -11]$$

$$\Rightarrow [-3m + 4n \ 4m - 3n] = [10 \ -11]$$

$$\Rightarrow -3m + 4n = 10 \quad \dots(i)$$

$$\text{and } 4m - 3n = -11 \quad \dots(ii)$$

On solving Eqs. (i) and (ii), we get
 $n = 1, m = -2$
 Now, $3m + 7n = 3(-2) + 7(1) = 1$

304 (b)

We know that
 $A(\text{adj } A) = |A| I$
 If $A = \begin{bmatrix} \cos x & \sin x \\ -\sin x & \cos x \end{bmatrix}$, then $|A| = 1$
 $\therefore A(\text{adj } A) = I \Rightarrow KI = I \Rightarrow k = 1$

305 (a)

$$\because |A| = 3, \text{adj}(A) = \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{3} \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix}$$

$$\Rightarrow (A^{-1})^3 = \frac{1}{27} \begin{bmatrix} 1 & -2 \\ 0 & 3 \end{bmatrix}^3$$

$$= \frac{1}{27} \begin{bmatrix} 1 & -26 \\ 0 & 27 \end{bmatrix}$$

306 (d)

Given, $M = [a_{uv}]_{n \times n}$
 $= [\sin(\theta_u - \theta_v) + i \cos(\theta_u - \theta_v)]$
 $\Rightarrow \bar{M} = [\sin(\theta_u - \theta_v) - i \cos(\theta_u - \theta_v)]$
 $\Rightarrow (\bar{M})^T = [\sin(\theta_v - \theta_u) - i \cos(\theta_v - \theta_u)]$
 $= [-\sin(\theta_u - \theta_v) - i \cos(\theta_u - \theta_v)]$
 $= -[\sin(\theta_u - \theta_v) + i \cos(\theta_u - \theta_v)]$
 $= -M$

307 (a)

If given system of equations have infinitely many solutions, then

$$\begin{vmatrix} 2 & -1 & 1 \\ 1 & -2 & 1 \\ \lambda & -1 & 2 \end{vmatrix} = 0$$

$$\Rightarrow 2(-4 + 1) + 1(2 - \lambda) + 1(-1 + 2\lambda) = 0$$

$$\Rightarrow -6 + 2 - \lambda - 1 + 2\lambda = 0$$

$$\Rightarrow \lambda - 5 = 0$$

$$\Rightarrow \lambda = 5$$

308 (c)

If $AB = O$, then A and B may be equal to O individually. It is not necessary in any condition

309 (b)

$$E(\theta)E(\phi) = \begin{bmatrix} \cos^2\theta & \cos\theta \sin\theta \\ \cos\theta \sin\theta & \sin^2\theta \end{bmatrix}$$

$$\times \begin{bmatrix} \cos^2\phi & \cos\phi \sin\phi \\ \cos\phi \sin\phi & \sin^2\phi \end{bmatrix}$$

$$= \begin{bmatrix} \cos^2\theta \cos^2\phi + \cos\theta \sin\theta \cos\phi \sin\phi \\ \cos\theta \sin\theta \cos^2\phi + \sin^2\theta \cos\phi \sin\phi \\ \cos^2\theta \cos\phi \sin\phi + \cos\theta \sin\theta \sin^2\phi \\ \cos\theta \sin\theta \cos\phi \sin\phi + \sin^2\theta \sin^2\phi \end{bmatrix}$$

$$= \begin{bmatrix} \cos\theta \cos\phi \cos(\theta - \phi) & \cos\theta \sin\phi \cos(\theta - \phi) \\ \cos\phi \sin\theta \cos(\theta - \phi) & \sin\theta \sin\phi \cos(\theta - \phi) \end{bmatrix}$$

$$= \begin{bmatrix} \cos\theta \cos\phi \cos(2n+1)\frac{\pi}{2} \\ \cos\theta \sin\phi \cos(2n+1)\frac{\pi}{2} \\ \cos\theta \sin\phi \cos(2n+1)\frac{\pi}{2} \\ \sin\theta \sin\phi \cos(2n+1)\frac{\pi}{2} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \quad \left[\because \cos(2n+1)\frac{\pi}{2} = 0 \right]$$

310 (c)

Given, $x \sin 3\theta - y + z = 0$
 $x \cos 2\theta + 4y + 3z = 0$ and $2x + 7y + 7z = 0$
 For non-trivial solution.

$$\begin{vmatrix} \sin 3\theta - 1 & 1 \\ \cos 2\theta & 4 & 3 \\ 2 & 7 & 7 \end{vmatrix} = 0$$

$$\Rightarrow \sin 3\theta(28 - 21) - \cos 2\theta(-7 - 7) + 2(-3 - 4) = 0$$

$$\Rightarrow 7 \sin 3\theta + 14 \cos 2\theta - 14 = 0$$

$$\Rightarrow 7(3 \sin \theta - 4 \sin^3 \theta) + 14(1 - 2 \sin^2 \theta) - 14 = 0$$

$$\Rightarrow -28 \sin^3 \theta - 28 \sin^2 \theta + 21 \sin \theta = 0$$

$$\Rightarrow -7 \sin \theta(4 \sin^2 \theta + 4 \sin \theta - 3) = 0$$

$$\Rightarrow \sin \theta(2 \sin \theta + 3)(2 \sin \theta - 1) = 0$$

$$\Rightarrow \sin \theta = 0, \sin \theta = \frac{1}{2} \quad \left(\because \sin \theta \neq -\frac{3}{2} \right)$$

$$\Rightarrow \theta = n\pi, n\pi + (-1)^n \frac{\pi}{6}$$

311 (c)

$$\because kA = \begin{bmatrix} 0 & 3a \\ 2b & 24 \end{bmatrix}$$

$$\Rightarrow k \begin{bmatrix} 0 & 2 \\ 3 & -4 \end{bmatrix} = \begin{bmatrix} 0 & 3a \\ 2b & 24 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 0 & 2k \\ 3k & -4k \end{bmatrix} = \begin{bmatrix} 0 & 3a \\ 2b & 24 \end{bmatrix}$$

$$\Rightarrow 2k = 3a, 3k = 2b, -4k = 24$$

$$\Rightarrow a = \frac{2k}{3}, b = \frac{3k}{2}, k = -6$$

$$\therefore a = -4, b = -9, k = -6$$

312 (d)

$$\because \text{adj}(\text{adj } A) = |A|^{n-2} A$$

Here $n = 3$
 $\Rightarrow \text{adj}(\text{adj } A) = |A| A$

313 (c)

$$\begin{aligned} A^5 &= A^2 A^2 A = (A + I)(A + I)A \\ &= (A^2 + 2AI + I^2)A \\ &= (A + I + 2A + I)A = (3A + 2I)A \\ &= 3A^2 + 2IA = 3(A + I) + 2IA \\ &= 3A + 3I + 2A = 5A + 3I \end{aligned}$$

314 (c)

Matrices $A + B$ and AB are defined only if both A and B are of same order $n \times n$.

315 (a)

$$\begin{aligned} UV + XY &= [2 \ -3 \ 4] \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix} + [0 \ 2 \ 3] \begin{bmatrix} 2 \\ 2 \\ 4 \end{bmatrix} \\ &= [6 \ -6 \ 4] + [0 \ 4 \ 12] = [4] + [16] = [20] \end{aligned}$$

316 (b)

For matrix $A = \begin{bmatrix} a & 2 \\ 2 & 4 \end{bmatrix}$ to be singular,

$$\begin{aligned} \begin{vmatrix} a & 2 \\ 2 & 4 \end{vmatrix} &= 0 \\ \Rightarrow 4a - 4 &= 0 \\ \Rightarrow a &= 1 \end{aligned}$$

317 (a)

$$\begin{aligned} \text{adj}(A) &= \begin{bmatrix} 4y & -x \\ -x^2 & 1 \end{bmatrix} \\ \therefore \text{adj}(A) + B &= \begin{bmatrix} 4y & -x \\ -x^2 & 1 \end{bmatrix} + \begin{bmatrix} -3 & 1 \\ 1 & 0 \end{bmatrix} \\ \Rightarrow \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} &= \begin{bmatrix} 4y - 3 & -x + 1 \\ -x^2 + 1 & 1 + 0 \end{bmatrix} \\ \Rightarrow 4y - 3 &= 1 \Rightarrow y = 1 \\ \text{and } -x + 1 &= 0 \Rightarrow x = 1 \end{aligned}$$

318 (c)

$$\begin{aligned} |A| &= 1. (\cos^2 \alpha + \sin^2 \alpha) = 1 \\ \text{Now, } A^{-1} &= \frac{1}{|A|} \text{adj}(A) = \text{adj}(A) \end{aligned}$$

319 (a)

We have,
 $(A + B)(A - B) = A^2 - AB + BA - B^2$
 So, option (a) is correct.

320 (a)

$$\begin{aligned} \text{Given, } (1 - x)f(x) &= 1 + x \\ \Rightarrow (I - A)f(A) &= (I + A) \quad (\because \text{Put } x = A) \\ \Rightarrow f(A) &= (I - A)^{-1}(I + A) \\ &= \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \right)^{-1} \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix} \right) \\ \Rightarrow f(A) &= \begin{bmatrix} 0 & -2 \\ -2 & 0 \end{bmatrix}^{-1} \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix} \\ &= \frac{\begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}}{-4} \\ &= \frac{\begin{bmatrix} 4 & 4 \\ 4 & 4 \end{bmatrix}}{-4} = \begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix} \end{aligned}$$

321 (b)

We have,
 $AA^{-1} = I$

$$\begin{aligned} \Rightarrow \det(AA^{-1}) &= \det(I) \\ \Rightarrow \det(A) \det(A^{-1}) &= 1 \\ \left[\because \det(AB) &= \det(A) \det(B) \right. \\ &\left. \text{and } \det(I) = 1 \right] \\ \Rightarrow \det(A^{-1}) &= \frac{1}{\det(A)} \end{aligned}$$

322 (b)

$$\begin{aligned} \text{Since } AB &= 3I \\ \Rightarrow A^{-1}AB &= 3IA^{-1} \\ \Rightarrow B &= 3A^{-1} \\ \Rightarrow A^{-1} &= \frac{B}{3} \end{aligned}$$

323 (b)

$$\text{We have, } \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 - x_1 & y_2 - y_1 & 0 \\ x_3 - x_1 & y_3 - y_1 & 0 \end{vmatrix} = 0$$

[using $R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1$]

\therefore The given points $(x_1, y_1), (x_2, y_2)$ and (x_3, y_3) are collinear, therefore the rank of matrix is always greater than 0 and less than 3.

324 (a)

$$\begin{aligned} \because A^2 &= A \cdot A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} \\ &= \begin{bmatrix} 4 + 1 & -2 - 2 \\ -2 - 2 & 1 + 4 \end{bmatrix} \\ &= \begin{bmatrix} 5 & -4 \\ -4 & 5 \end{bmatrix} \\ \text{And } 4A - 3I &= 4 \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix} - 3 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 8 & -4 \\ -4 & 8 \end{bmatrix} - \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 5 & -4 \\ -4 & 5 \end{bmatrix} \\ \therefore A^2 &= 4A - 3I \end{aligned}$$

325 (a)

$$\begin{aligned} \text{Given, } A &= \begin{bmatrix} -2 & 6 \\ -5 & 7 \end{bmatrix} \\ \therefore \text{adj } A &= \begin{bmatrix} 7 & -6 \\ 5 & -2 \end{bmatrix} \end{aligned}$$

326 (c)

$$\begin{aligned} \text{Given, } A &= \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix} \\ \Rightarrow |A| &= 1 \\ \therefore A \text{ adj}(A) &= \begin{bmatrix} 3 & 4 \\ 5 & 7 \end{bmatrix} \begin{bmatrix} 7 & -4 \\ -5 & 3 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = 1 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = |A|I \end{aligned}$$

327 (c)

$$\begin{aligned} AB &= \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} = \begin{bmatrix} a & 2b \\ 3a & 4b \end{bmatrix} \\ \text{And } BA &= \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} a & 2b \\ 3a & 4b \end{bmatrix} \end{aligned}$$

If $AB = BA$, then $a = b$

Hence, $AB = BA$ is possible for infinitely many values of B 's.

328 (b)

We have,

$$2A + 3B = \begin{bmatrix} 2 & -1 & 4 \\ 3 & 2 & 5 \end{bmatrix} \quad \dots(i)$$

$$\text{and } A + 2B = \begin{bmatrix} 5 & 0 & 3 \\ 1 & 6 & 2 \end{bmatrix} \quad \dots(ii)$$

On multiplying Eq.(ii) by 2 and then subtracting Eq.(i) from Eq.(ii), we get

$$B = 2 \begin{bmatrix} 5 & 0 & 3 \\ 1 & 6 & 2 \end{bmatrix} - \begin{bmatrix} 2 & -1 & 4 \\ 3 & 2 & 5 \end{bmatrix} = \begin{bmatrix} 8 & 1 & 2 \\ -1 & 10 & -1 \end{bmatrix}$$

329 **(b)**

Determinant of unit matrix of any order is 1

330 **(b)**

$$AB = \begin{bmatrix} 3 & 5 \\ 2 & 0 \end{bmatrix} \begin{bmatrix} 1 & 17 \\ 0 & -10 \end{bmatrix} = \begin{bmatrix} 3 & 1 \\ 2 & 34 \end{bmatrix}$$

$$\Rightarrow |AB| = 102 - 2 = 100$$

331 **(a)**

$$\text{We have, } A = \begin{bmatrix} 6 & 8 & 5 \\ 4 & 2 & 3 \\ 9 & 7 & 1 \end{bmatrix} \text{ and } A' = \begin{bmatrix} 6 & 4 & 9 \\ 8 & 2 & 7 \\ 5 & 3 & 1 \end{bmatrix}$$

$$\therefore \text{Symmetric matrix, } B = \frac{A+A'}{2}$$

$$= \frac{1}{2} \left\{ \begin{bmatrix} 6 & 8 & 5 \\ 4 & 2 & 3 \\ 9 & 7 & 1 \end{bmatrix} + \begin{bmatrix} 6 & 4 & 9 \\ 8 & 2 & 7 \\ 5 & 3 & 1 \end{bmatrix} \right\}$$

$$= \frac{1}{2} \begin{bmatrix} 12 & 12 & 14 \\ 12 & 4 & 10 \\ 14 & 10 & 2 \end{bmatrix} = \begin{bmatrix} 6 & 6 & 7 \\ 6 & 2 & 5 \\ 7 & 5 & 1 \end{bmatrix}$$

332 **(b)**

Since A is non-singular. Therefore, A^{-1} exists

$$\text{Now, } A(\text{adj } A) = |A|I = (\text{adj } A)A$$

$$\Rightarrow |A||\text{adj } A| = |A|^n = |\text{adj } A||A|$$

$$\Rightarrow |\text{adj } A| = |A|^{n-1} \quad [\because |A| \neq 0]$$

333 **(a)**

$$\therefore A + B = \begin{bmatrix} 1 & -1 \\ 2 & -1 \end{bmatrix} + \begin{bmatrix} 1 & a \\ 4 & b \end{bmatrix} = \begin{bmatrix} 2 & -1 + a \\ 6 & -1 + b \end{bmatrix}$$

$$\Rightarrow (A + B)^2 = \begin{bmatrix} 2 & -1 + a \\ 6 & -1 + b \end{bmatrix} \begin{bmatrix} 2 & -1 + a \\ 6 & -1 + b \end{bmatrix} = \begin{bmatrix} -2 + 6a & -1 + a - b + ab \\ 6 + 6b & -5 + 6a - 2b + b^2 \end{bmatrix}$$

$$\text{and } A^2 = \begin{bmatrix} 1 & -1 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 2 & -1 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$\text{Also, } B^2 = \begin{bmatrix} 1 & a \\ 4 & b \end{bmatrix} \begin{bmatrix} 1 & a \\ 4 & b \end{bmatrix} = \begin{bmatrix} 1 + 4a & a + ab \\ 4 + 4b & 4a + b^2 \end{bmatrix}$$

$$\text{Given, } (A + B)^2 = A^2 + B^2$$

$$\therefore \begin{bmatrix} -2 + 6a & -1 + a - b + ab \\ 6 + 6b & -5 + 6a - 2b + b^2 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} 1 + 4a & a + ab \\ 4 + 4b & 4a + b^2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -2 + 6a & -1 + a - b + ab \\ 6 + 6b & -5 + 6a - 2b + b^2 \end{bmatrix} = \begin{bmatrix} 4a & a + ab \\ 4 + 4b & -1 + 4a + b^2 \end{bmatrix}$$

On comparing both sides, we get

$$-2 + 6a = 4a \quad \text{and} \quad 6 + 6b = 4 + 4b$$

$$\Rightarrow a = 1 \quad \text{and} \quad b = -1$$

334 **(d)**

$$(AB^{-1}C)^{-1} = C^{-1}(B^{-1})^{-1}A^{-1} = C^{-1}BA^{-1}$$