



5. The graph shows the variation of stopping potential with frequency of incident radiation for two photosensitive metals A and B. Which one of the two has higher value of work function? Justify your answer. **All India 2014**




6. The graph shows variation of stopping potential  $V_0$  versus frequency of incident radiation  $\nu$  for two photosensitive metals A and B. Which one of the two metals has higher threshold frequency and why?

**All India 2014**



7. In photoelectric effect, why should the photoelectric current increase as the intensity of monochromatic radiation incident on a photosensitive surface is increased? Explain. **Foreign 2014**

8. The given graph shows the variation of photoelectric current ( $I$ ) versus applied voltage ( $V$ ) for two different photosensitive materials and for two different intensities of the incident radiations. Identify the pairs of curves that



# Previous Years Examination Questions

## ✍ 1 Mark Questions

1. Define the term threshold frequency, in the context of photoelectric emission. **Delhi 2019**
2. Define the term intensity in photon picture of electromagnetic radiation. **Delhi 2019**
3. Draw graphs showing variation of photoelectric current with applied voltage for two incident radiations of equal frequency and different intensities. Mark the graph for the radiation of higher intensity. **Delhi 2018**
4. Define intensity of radiation on the basis of photon picture of light. Write its SI unit. **All India 2014, 2012**

corresponds to different materials but same intensity of incident radiation.

Delhi 2013

9. Show on a plot the nature of variation of photoelectric current with the intensity of radiation incident on a photosensitive surface. Delhi 2013C
10. Why is photoelectric emission not possible at all frequencies? All India 2012C
11. For a given photosensitive material and with a source of constant frequency of incident radiation, how does the photocurrent vary with the intensity of incident light? All India 2011C
12. Define the term stopping potential in relation to photoelectric effect. All India 2011
13. Show the variation of photoelectric current with collector plate potential for different frequencies but same intensity of incident radiation. Foreign 2011, All India 2010
14. Show the variation of photoelectric current with collector plate potential for different intensities but same frequency of incident radiation. Foreign 2011, All India 2010; 2011

## 2 Marks Questions

15. Photoelectric emission occurs when a surface is irradiated with the radiation of frequency (i)  $\nu_1$  and (ii)  $\nu_2$ . The maximum kinetic energy of the electrons emitted in the two cases are  $K$  and  $2K$  respectively. Obtain the expression for the threshold frequency for the surface. CBSE 2022 (Term-II)
16. (i) Name the factors on which photoelectric emission from a surface depends.  
(ii) Define the term threshold frequency for a photosensitive material.  
CBSE 2022 (Term-II)
17. If the frequency of light incident on the cathode of a photo-cell is increased, how will the following be affected? Justify your answer.


- (i) Energy of the photoelectrons.  
(ii) Photocurrent. All India 2020

18. Light of same wavelength is incident on three photo sensitive surfaces  $A$ ,  $B$  and  $C$ . The following observations are recorded.
- (i) From surface  $A$ , photoelectrons are not emitted.  
(ii) From surface  $B$ , photoelectrons are just emitted.  
(iii) From surface  $C$ , photoelectrons with some kinetic energy are emitted.

Compare the threshold frequencies of the three surfaces and justify your answer.

All India 2020

19. Figure shows the stopping potential ( $V_0$ ) for the photoelectron *versus*  $\left(\frac{1}{\lambda}\right)$  graph, for two metals  $A$  and  $B$ ,  $\lambda$  being the wavelength of incident light.



- (i) How is the value of Planck's constant determined from the graph?  
(ii) If the distance between the light source and the surface of metal  $A$  is increased, how will the stopping potential for the electrons emitted from it be effected? Justify your answer. Delhi 2020
20. Why is wave theory of electromagnetic radiation not able to explain photoelectric effect? How does photon picture resolve this problem? Delhi 2019
21. Explain with the help of Einstein's photoelectric equation any two observed features in photoelectric effect which cannot be explained by wave theory.

Delhi 2019

22. (i) Define the terms (a) threshold frequency and (b) stopping potential in photoelectric effect.  
 (ii) Plot a graph of photocurrent *versus* anode potential for a radiation of frequency  $\nu$  and intensities  $I_1$  and  $I_2$  ( $I_1 < I_2$ ). **Delhi 2019**

23. If light of wavelength 412.5 nm is incident on each of the metals given below, which ones will show photoelectric emission and why?

Metal	Work Function (eV)
Na	1.92
K	2.15
Ca	3.20
Mo	4.17


**Delhi 2018**

24. Find the frequency of light which ejects electrons from a metal surface, fully stopped by a retarding potential of 3.3 V. If photoelectric emission begins in this metal at a frequency of  $8 \times 10^{14}$  Hz, calculate the work function (in eV) for this metal. **All India 2018C**

25. Monochromatic light of frequency  $6.0 \times 10^{14}$  Hz is produced by a laser. The power emitted is  $2.0 \times 10^{-3}$  W. Calculate the (i) energy of a photon in the light beam and (ii) number of photons emitted on an average by the source. **All India 2018C**

26. In the wave picture of light, intensity of light is determined by the square of the amplitude of the wave. What determines the intensity in the photon picture of light? **All India 2016**

27. Using the graph shown in the figure for stopping potential *versus* the incident frequency of photons, calculate Planck's constant.



**Delhi 2015**

28. (i) Monochromatic light of frequency  $6.0 \times 10^{14}$  Hz is produced by a laser. The power emitted is  $2.0 \times 10^{-3}$  W. Estimate the number of photons emitted per second on an average by the source.

- (ii) Draw a plot showing the variation of photoelectric current *versus* the intensity of incident radiation on a given photosensitive surface. **Delhi 2014**

- Or (i) Monochromatic light of frequency  $5.0 \times 10^{14}$  Hz is produced by laser. The power emitted is  $3.0 \times 10^{-3}$  W. Estimate the number of photons emitted per second on an average by the source.

- (ii) Draw a plot showing the variation of photoelectric current *versus* the intensity of incident radiation on a given photosensitive surface. **Delhi 2014**

29. Write three basic properties of photons which are used to obtain Einstein's photoelectric equation. Use this equation to draw a plot of maximum kinetic energy of the electrons emitted *versus* frequency of incident radiation.

**All India 2014**

30. (i) Define the term stopping potential.  
 (ii) Plot a graph showing the variation of photoelectric current as a function of anode potential for two light beams of same intensity but of different frequencies  $\nu_1$  and  $\nu_2$  ( $\nu_2 > \nu_1$ ).

**All India 2014C**

31. (i) Define the term threshold frequency as used in photoelectric effect.  
 (ii) Plot a graph showing the variation of photoelectric

current as a function of anode potential for two light beams having the same frequency  $h\nu$  different intensities  $I_1$  and  $I_2(I_1 > I_2)$ .

**All India 2014C**

- 32.** Two monochromatic radiations of frequencies  $\nu_1$  and  $\nu_2(\nu_1 > \nu_2)$  and having the same intensity are in turn, incident on a photosensitive surface to cause photoelectric emission.

Explain giving reason in which case (i) more number of electrons will be emitted and (ii) maximum kinetic energy of the emitted photoelectrons will be more.

**Delhi 2014C**

- 33.** Two monochromatic radiations, blue and violet of the same intensity are incident on a photosensitive surface and cause photoelectric emission. Would (i) the number of electrons emitted per second and (ii) the maximum kinetic energy of the electrons be equal in the two cases? Justify your answer. **Delhi 2010**


- 34.** Write Einstein's photoelectric equation. State clearly the three salient features observed in photoelectric effect which can be explained on the basis of above equation. **All India 2010**

- 35.** Plot a graph showing the variation of stopping potential with the frequency of incident radiation for two different photosensitive materials having work-functions  $\phi_1$  and  $\phi_2$  ( $\phi_1 > \phi_2$ ). On what factors does the
- slope and
  - intercept of the lines depend?

**Delhi 2010**

### 3 Marks Questions


- 36.** State Einstein's photoelectric explaining the symbols used.



Light of frequency  $\nu$  incident is on a photosensitive surface. A graph of the square of the maximum speed of the electrons ( $v_{\max}^2$ ) versus  $\nu$  is obtained as shown in the figure. Using Einstein's photoelectric equation, obtain expressions for (i) Planck's constant and (ii) work function of the given photosensitive material in terms of parameters  $l$ ,  $n$  and mass of the electron  $m$ . **All India 2018C**

- 37.** (i) How does one explain the emission of electrons from a photosensitive surface with the help of Einstein's photoelectric equation?  
 (ii) The work function of the following metals is given as Na = 2.75 eV, K = 2.3 eV, Mo = 4.17 eV and Ni = 5.15 eV. Which of these metals will not cause photoelectric emission for radiation of wavelength 3300 Å from a laser source placed 1 m away from these metals? What happens if the laser source is brought nearer and placed 50 cm away? **Delhi 2017**
- 38.** (i) State two important features of Einstein's photoelectric equation.  
 (ii) Radiation of frequency  $10^{15}$  Hz is incident on two photosensitive surfaces  $P$  and  $Q$ . There is no photoemission from surface  $P$ . Photoemission occurs from surface  $Q$  but photoelectrons have zero kinetic energy. Explain these observations and find the value of work function for surface  $Q$ . **Delhi 2017**

39. In the study of a photoelectric effect, the graph between the stopping potential  $V$  and frequency  $\nu$  of the incident radiation on two different metals  $P$  and  $Q$  is shown below.



- (i) Which one of the two metals has higher threshold frequency?
- (ii) Determine the work function of the metal which has greater value.
- (iii) Find the maximum kinetic energy of electron emitted by light of frequency  $8 \times 10^{14}$  Hz for this metal. **Delhi 2017**
40. Sketch the graphs showing variation of stopping potential with frequency of incident radiations for two photosensitive materials  $A$  and  $B$  having threshold frequencies  $\nu_A > \nu_B$ .
- (i) In which case, is the stopping potential more and why?
- (ii) Does the slope of the graph depend on the nature of the material used? Explain. **All India 2016**
41. Plot a graph showing the variation of photoelectric current with intensity of light. The work function for the following metals is given. Na : 2.75 eV and Mo : 4.175 eV. Which of these will not give photoelectron emission from a radiation of wavelength 3300 Å from a laser beam? What happens if the source of laser beam is brought closer? **Foreign 2016**
42. Define the term “cut-off frequency” in photoelectric emission. The threshold frequency of a metal is  $f$ . When the light of frequency  $2f$  is incident on the metal plate, the maximum velocity of photoelectron is  $v_1$ . When the frequency of the incident radiation is increased to  $5f$ , the maximum velocity of photoelectrons is  $v_2$ . Find the ratio  $v_1 : v_2$ . **Foreign 2016**

43. Write three characteristic features in photoelectric effect which cannot be explained on the basis of wave theory of light, but can be explained only using Einstein's equation. **Delhi 2016**
44. (i) Write the important properties of photons which are used to establish Einstein's photoelectric equation.  
(ii) Use this equation to explain the concept of (a) threshold frequency and (b) stopping potential. **Delhi 2015**
45. (i) Describe briefly three experimentally observed features in the phenomenon of photoelectric effect.  
(ii) Discuss briefly how wave theory of light cannot explain these features. **Delhi 2015**
46. Write Einstein's photoelectric equation and mention which important features in photoelectric effect can be explained with the help of this equation.  
The maximum kinetic energy of the photoelectrons gets doubled when the wavelength of light incident on the surface changes from  $\lambda_1$  to  $\lambda_2$ . Derive the expressions for the threshold wavelength  $\lambda_0$  and work function for the metal surface. **Delhi 2015, All India 2014**
47. A beam of monochromatic radiation is incident on a photosensitive surface. Answer the following questions giving reasons.  
(i) Do the emitted photoelectrons have the same kinetic energy?  
(ii) Does the kinetic energy of the emitted electrons depend on the intensity of incident radiation?  
(iii) On what factors does the number of emitted photoelectrons depend? **Foreign 2015**
48. Write Einstein's photoelectric equation and point out any two characteristic properties of photons on which this equation is based. Briefly explain three observed features which can be explained by this equation. **All India 2013**

- 49.** (i) State three important properties of photons which describe the particle picture of electromagnetic radiation.  
(ii) Use Einstein's photoelectric equation to define the terms :  
(a) Stopping potential and  
(b) Threshold frequency. **Delhi 2013C**
- 50.** (i) Why photoelectric effect cannot be explained on the basis of wave nature of light? Give reasons.  
(ii) Write the basic features of photon picture of electromagnetic radiation on which Einstein's photoelectric equation is based. **Delhi 2013**
- 51.** Draw a graph between the frequency of incident radiation ( $\nu$ ) and the maximum kinetic energy of the electrons emitted from the surface of a photosensitive material. State clearly how this graph can be used to determine  
(i) Planck's constant and  
(ii) work function of the material. **Foreign 2012**
- 52.** Write two characteristic features observed in photoelectric effect which supports the photon pictures of electromagnetic radiation. Draw a graph between the frequency of incident radiation ( $\nu$ ) and the maximum kinetic energy of the electrons emitted from the surface of a photosensitive material. State clearly how this graph can be used to determine the  
(i) Planck constant and  
(ii) work function of the material? **Foreign 2012**
- 53.** Write Einstein's photoelectric equation. State clearly how this equation is obtained using the photon picture of electromagnetic radiation. Write the three salient features observed in photoelectric effect which can be explained using this equation. **Delhi 2012**
- 54.** Define the terms cut-off voltage and threshold frequency in relation to the phenomenon of photoelectric effect. Using Einstein's photoelectric equation show how the cut-off voltage and threshold frequency for a given photosensitive material can be determined with the help of a suitable plot/graph. **All India 2012**
- 55.** Light of wavelength  $2000 \text{ \AA}$  falls on a metal surface of work function  $4.2 \text{ eV}$ . What is the kinetic energy (in eV) of the fastest electrons emitted from the surface?  
(i) What will be the change in the energy of the emitted electrons if the intensity of light with same wavelength is doubled?  
(ii) If the same light falls on another surface of work function  $6.5 \text{ eV}$ , what will be the energy of emitted electrons? **Foreign 2011**
- 56.** Draw a plot showing the variation of photoelectric current with collector plate potential for two different frequencies,  $\nu_2 > \nu_1$  of incident radiation having the same intensity. In which case will the stopping potential be higher? Justify your answer. **All India 2011**
- 57.** (i) Ultraviolet light of wavelength  $2271 \text{ \AA}$  from a  $100 \text{ W}$  mercury source is incident on a photocell made of molybdenum metal. If the stopping potential is  $1.3 \text{ V}$ , estimate the work function of the metal.  
(ii) How would the photocell respond to high intensity ( $10^5 \text{ W/m}^2$ ) red light of wavelength  $6328 \text{ \AA}$  produced by a He-Ne laser? **Delhi 2011C**
- 58.** Define the terms threshold frequency and stopping potential in the study of photoelectric emission. Explain briefly the reasons why wave theory of light is not able to explain the observed features in photoelectric effect? **Foreign 2010**

## Explanations


- Refer to text on page 335 (Laws of Photoelectric Emission). (1)
- The intensity of radiation is defined as the rate of emitted energy from unit surface area in a given interval of time.  
Its SI unit is watt/metre<sup>2</sup>. (1)

# Previous Years

## Examination Questions

### 1 Mark Questions

1. Draw a plot showing the variation of de-Broglie wavelength of electron as a function of its KE. Delhi 2015C
2. Figure shows a plot of  $1/\sqrt{V}$ , where  $V$  is the accelerating potential *versus* the de-Broglie wavelength  $\lambda$  in the case of two particles having same charge  $q$  but different masses  $m_1$  and  $m_2$ . Which line ( $A$  or  $B$ ) represents a particle of large mass? Delhi 2014



3. Write the expression for the de-Broglie wavelength associated with a charged particle having charge  $q$  and mass  $m$ , when it is accelerated by a potential. All India 2013
4. State de-Broglie hypothesis. Delhi 2012
5. A proton and an electron have same kinetic energy. Which one has greater de-Broglie wavelength and why? All India 2012
6. A particle is moving three times as fast as an electron. The ratio of the de-Broglie wavelength of the particle to that of the electron is  $8.3 \times 10^{-4}$ . Calculate the particle's mass and identify the particle. All India 2011C
7. Write the relationship of de-Broglie wavelength  $\lambda$  associated with a particle of mass  $m$  in terms of its kinetic energy  $E$ . Delhi 2011C

8. Show graphically, the variation of de-Broglie wavelength ( $\lambda$ ) with the potential ( $V$ ) through which an electron is accelerated from rest. Delhi 2011

## 2 Marks Questions

9. A proton and a electron have the same de-Broglie wavelength  $\lambda$ . Prove that the energy of the proton is  $(2m\lambda c/h)$  times the kinetic energy of the electron. All India 2019
10. The wavelength of light from the spectral emission line of sodium is 590 nm. Find the kinetic energy at which the electron would have the same de-Broglie wavelength. All India 2019
11. Plot a graph showing variation of de-Broglie wavelength ( $\lambda$ ) associated with a charged particle of mass  $m$  versus  $\frac{1}{\sqrt{V}}$ , where  $V$  is the potential difference through which the particle is accelerated. How does this graph give us the information regarding the magnitude of the charge of the particle? Delhi 2019
12. (i) Plot a graph showing variation of de-Broglie wavelength ( $\lambda$ ) associated with a charged particle of mass  $m$  versus  $\sqrt{V}$ , where  $V$  is the accelerating potential.  
(ii) An electron, a proton and an alpha particle have the same kinetic energy. Which one has the shortest wavelength? Delhi 2019
13. An  $\alpha$ -particle and a proton are accelerated through the same potential difference. Find the ratio of their de-Broglie wavelengths. Delhi 2017; All India 2010
14. The wavelength  $\lambda$  of a photon and the de-Broglie wavelength of an electron have the same value. Show that energy of a photon is  $(2\lambda mc/h)$  times the kinetic energy of electron, where  $m$ ,  $c$  and  $h$  have their usual meaning. Foreign 2016
15. A proton and an  $\alpha$ -particle have the same de-Broglie wavelength. Determine the ratio of (i) their accelerating potentials (ii) their speeds. Delhi 2015
16. A proton and a deuteron are accelerated through the same accelerating potential. Which one of the two has  
(i) greater value of de-Broglie wavelength associated with it and  
(ii) less momentum?  
Give reasons to justify your answer. Delhi 2014
17. A deuteron and an  $\alpha$ -particle are accelerated with the same accelerating potential. Which one of the two has  
(i) greater value of de-Broglie wavelength, associated with it and  
(ii) less kinetic energy? Explain. Delhi 2014
18. X-rays fall on a photosensitive surface to cause photoelectric emission. Assuming that the work-function of the surface can be neglected, find the relation between the de-Broglie wavelength ( $\lambda$ ) of the electrons emitted to the energy ( $E$ ) of the incident photons. Draw of the graph for  $\lambda$  as function of  $E$ . Delhi 2014 C
19. An electron is revolving around the nucleus with a constant speed of  $2.2 \times 10^8$  m/s. Find the de-Broglie wavelength associated with it. All India 2014 C
20. An electron is accelerated through a potential difference of 100 V. What is the de-Broglie wavelength associated with it? To which part of the electromagnetic spectrum does this value of wavelength correspond? Delhi 2010
- Or An electron is accelerated through a potential difference of 144 V. What is the de-Broglie wavelength associated with it? To which part of electromagnetic spectrum does this wavelength correspond? Delhi 2010
- Or An electron is accelerated through a potential difference of 64 V. What is the de-Broglie wavelength associated with it? To which part of the electromagnetic spectrum does this value of wavelength correspond? Delhi 2010




21. Find the ratio of de-Broglie wavelengths associated with

- (i) protons, accelerated through a potential of 128 V and
- (ii)  $\alpha$ -particles, accelerated through a potential of 64 V. **Delhi 2010C**

22. The ratio between the de-Broglie wavelengths associated with protons, accelerated through a potential of 512 V and  $\alpha$ -particles, accelerated through a potential of  $X$  volt is found to be one. Find the value of  $X$ . **Delhi 2010C**

23. The two lines marked  $A$  and  $B$  in the given figure, show a plot of de-Broglie wavelength  $\lambda$  versus  $1/\sqrt{V}$ , where  $V$  is the accelerating potential for two nuclei  ${}^2_1\text{H}$  and  ${}^3_1\text{H}$ .



- (i) What does the slope of the lines represent?
- (ii) Identify which of the lines corresponded to these nuclei.

**All India 2010**

26. (i) Determine the de-Broglie wavelength of a proton whose kinetic energy is equal to the rest mass energy of an electron. Mass of a proton 1836 times that of electron.

- (ii) In which region of electromagnetic spectrum does this wavelength lie? **All India 2011C**

27. (i) The mass of a particle moving with velocity  $5 \times 10^6$  m/s has de-Broglie wavelength associated with it to be 0.135 nm. Calculate its mass.

- (ii) In which region of the electromagnetic spectrum does this wavelength lie? **All India 2011C**

28. (i) A particle is moving three times as fast as an electron. The ratio of the de-Broglie wavelength of the particle to that of the electron is  $1.813 \times 10^{-4}$ . Calculate the particle's mass and identify the particle.

- (ii) An electron and a proton have the same kinetic energy. Which of the two will have larger de-Broglie wavelength? Give reason. **All India 2011C**

29. An electron and a photon each have a wavelength 1nm. Find

- (i) their momenta
- (ii) the energy of the photon and
- (iii) the kinetic energy of electron.

**All India 2011C**

### 3 Marks Questions

24. Find the ratio of the de-Broglie wavelengths associated with an alpha particle and proton, if both

- (i) have the same speeds,
- (ii) have the same kinetic energy,
- (iii) are accelerated through the same potential difference.

25. An electron is accelerated from rest through a potential difference of 100V. Find

- (i) the wavelength associated with
- (ii) the momentum of and
- (iii) the velocity required by the electron